

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

1 of 40

PM 91

275-61

6-16-98

JUN 16 1998

S545425	301
-	17

Ms. Debbie Forrey
 Abbott Laboratories
 1401 Sheridan Road
 D28R, Bldg. A1
 North Chicago, IL

Dear Ms. Forrey:

Subject: ProGibb 4% Plant Growth Regulator Solution
 EPA Reg. No. 275-61
 Your Fax of June 9, 1998

On December 8, 1997, you requested to amend the label of the cited product, primarily to add a use on hybrid rice. The subject fax represents a revised label, sent in response to the Biopesticides and Pollution Prevention Division (BPPD) label review letter of May 5, 1998.

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy of the label is enclosed for your records.


Please note, however, that minor label deficiencies may exist on the subject label. Also, there is an unresolved tolerance/application rate question for the crops lemons, oranges and grapefruit versus the crop group citrus fruits, relative to 40 CFR 180.1098 and 40 CFR 180.224, respectively. The BPPD will address these outstanding issues in a forthcoming letter. Further revision of the label may be required at that time.

CONCURRENCES							
SYMBOL	7511C	7511L					
SURNAME	Greenway	<i>[Signature]</i>					
DATE	6/15/98	6/15/98					

2040

Should you have questions or concerns, please contact me, or telephone Denise Greenway on (703) 308-8263.

Sincerely,



Janet L. Andersen, Ph.D.
Biopesticides and Pollution
Prevention Division (7511C)

DGreenway/308-8263/6-9-98/progibb.ltr/GA3

30440

MASTER LABEL

Primary Product name: ProGibb 4% Plant Growth Regulator Solution

Active Ingredient:

Gibberellic Acid.....4.0% w/w

Inert Ingredients.....96.0% w/w

Total.....100.0% w/w

EPA Registration No. 275-61

Abbott Laboratories
Chemical and Agricultural Products Division
1401 Sheridan Road
North Chicago, IL 60064

ACCEPTED
with COMMENTS
in EPA Letter Dated

JUN 16 1998

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

275-61

4 of 40

SUB LABEL

ProGibb 4% Plant Growth Regulator Solution

For use on grape (table and wine varieties); citrus (orange, lemon, lime, tangerine, grapefruit); fruit (blueberry, cherry, non-bearing peach, Italian prune, strawberry); vegetable (rhubarb, artichoke, celery, lettuce seed, pepper, melon, cucumber, seed potato, spinach); floriculture (pompom chrysanthemum, statice, azalea); turf (golf), hops, cotton

50440

Plant Growth Regulator

PROGIBB® 4%

SOLUTION

Active Ingredient:

Gibberellic Acid.....	4.0% w/w
Inert Ingredients.....	96.0% w/w
Total.....	100.0% w/w

ProGibb 4% liquid contains approximately 1.0 gram active ingredient per fluid ounce of formulated product.

KEEP OUT OF REACH OF CHILDREN
WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES - Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

IF SWALLOWED - Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol.

IF INHALED - Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

IF ON SKIN - Wash with plenty of soap and water. Get medical attention.

EPA Reg. No. 275-61

EPA Est. No.

Abbott Laboratories
Chemical and Agricultural Products Division
1401 Sheridan Road
North Chicago, IL 60064

Net Contents: _____
This container will treat _____ acres at the maximum use rate, as recommended for use on _____.

60440

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING**

Causes substantial but temporary eye injury. Harmful if inhaled or absorbed through skin. Do not get in eyes or on clothing. Avoid breathing vapor or spray mist, and avoid contact with skin. Wear protective eyewear (goggles or face shield). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- * Long sleeved shirt
- * Long pants
- * Chemical resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, and viton
- * Shoes plus socks
- * Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- * Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- * Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning of equipment or disposing of equipment washwaters.

70/40

PHYSICAL OR CHEMICAL HAZARDS**FLAMMABLE!** Keep away from heat and open flame.**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

EXCEPTION: If the product is soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- * Coveralls
- * Chemical resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, and viton
- * Shoes plus socks
- * Protective eyewear

p984%01.ssm/5
rev. 05/04/88

80440

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter without appropriate protective clothing until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE:

Keep containers tightly closed when not in use. Keep away from heat and open flame.

PESTICIDE DISPOSAL:

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL DIRECTIONS FOR USE

Use only as directed. The label should be read thoroughly and understood before making applications. Keep out of reach of children.

Do not apply this product through any type of irrigation system.

Application recommendations:

ProGibb 4% contains gibberellic acid which is an extremely potent plant growth regulator; when applying plant growth regulators, deviations from the label directions in the rates, timings, water volumes, or the adoption of untested spray mixes, may result in undesirable effects. Always consult the Abbott Laboratories agricultural specialist in your area for the spray regimen best suited to your conditions.

-- Do not apply to plants under pest, nutritional, or water stress.

- When a range of rates is indicated, use the concentration and spray volume recommended locally by the Abbott Laboratories agricultural specialist.
- For optimum effectiveness, thorough spray coverage must be achieved; all parts of the plant or crop must receive the spray or desired results will not occur. Prepare solution concentrations by mixing the required amount of product with water in a clean, empty spray tank. Dispose of any unused spray material at the end of each day following local, state or federal law.
- For best results, the water pH should be around neutral, and always below 8.5.
- ProGibb 4% applications made under slow drying conditions (cool to warm temperatures, medium to high relative humidity, and no wind) will increase absorption by the plant, thus optimizing effectiveness. Night-time applications are encouraged when day-time conditions are not conducive to slow drying conditions.
- Product persistence: ProGibb 4% should be re-applied if significant rain occurs within 2 hours of application.
- Compatibility: The ProGibb 4% spray guidelines refer to the use of the product alone. The use of surfactants and other additives has been reported to be beneficial. However, data concerning the compatibility of ProGibb 4% with other agricultural compounds, except DiPel 2X, DiPel DF, or XenTari WDG (registered products of Abbott Laboratories; use as directed on each product label), are not available. Abbott Laboratories does not assume responsibility for unexpected results due to the tank mixing of ProGibb 4% with other products.
- DO NOT apply using ULV application methods. For aerial applications spray volumes must be greater than 2 gallons per acre (10 gallons per acre for tree crops).

ProGibb can be applied up to 7 days before harvest.

ProGibb 4% is a liquid and each fluid ounce contains approximately one (1) gram of active ingredient.

SPRAY GUIDELINES FOR CROP CATEGORIES

GRAPE

For all grapes, application is recommended by ground sprayer. Apply as a concentrate or dilute spray in sufficient water volume to insure thorough wetting. It is important to wet all flower clusters or berries thoroughly. For specific spray rates and timings, by variety, see accompanying tables. Do not exceed maximum rates.

SEEDLESS GRAPE

Do not apply more than 208 grams a.i./acre, per growing season, for all uses.

For cluster elongation and looser cluster forms ("Stretch"). To reduce costs of thinning, allow better air circulation to aid in the control of bunch rot, and increase light penetration to aid in sugar development.

Guide: Apply one to two applications before bloom when flower clusters are 2-to-5 inches long.

For decreased berry set ("Thinning"), reduced hand-thinning costs, and hastened maturity.

Guide: Apply one to three applications during bloom. When the bloom period is extended, subsequent sprays should be made 1-to-7 days after the first application.

NOTE: Higher amounts or multiple applications may cause an excess of shot berries or overthinning, especially in young vines or vines with high vigor.

For larger berries ("Sizing") and larger clusters when used in conjunction with established girdling and thinning practices.

Guide: Apply one to three applications beginning when the average berry size reaches "target" diameter (See Table 1). Timing of the subsequent sprays will be dictated by experience in the vineyard and temperatures occurring between sprays. Potential effect will be reduced if the final spray occurs more than two weeks after the first application.

110240

SEEDLESS GRAPE

(continued)

TABLE 1**APPLICATION RATES (GRAMS A.I./ACRE) FOR SEEDLESS GRAPE,
INCLUDING TARGET BERRY DIAMETERS**

Seedless Grape	Stretch	Thinning	Sizing	
	grams a.i./acre	grams a.i./acre	"Target" Diameter	grams a.i./acre
Perlette	8-to-16	*	4-to-5 mm	32-to-80
Flame	8-to-16	3-to-16	6-to-9 mm	20-to-80
Thompson	8-to-16	8-to-16	3-to-5 mm	32-to-80
Raisin	8-to-16	3-to-12	4-to-5 mm	4-to-12
All Other Seedless Grape	*	*	12-to-14 mm	8-to-48

*No recommendations available for this variety/timing at this time.

NOTE: Do not apply more than 208 grams a.i./acre per growing season for all uses.

SEEDED GRAPE**Emperor Grape**

For reducing berry shrivel. This use can also increase berry size.

Guide: Apply 20 grams a.i./acre as one application approximately two weeks after completion of berry shatter. This timing should correspond to a period when the predominant berry diameter ranges from 10-to-15 mm.

120440

SEEDED GRAPE
(continued)

Black Corinth (Zante Currant) Grape

- For improving berry size.

Guide: Apply 1-to-8 grams a.i./acre as one application 3-to-5 days after full bloom, but before shatter begins.

Wine Varieties

- For looser cluster to reduce incidence of bunch rot.

Guide: Apply one spray when shoots are 15-to-20 inches long. Use 100 gallons of water per acre. Clusters should average 3-to-4 inches in length and may range from 2-to-5 inches in length. Concentrations for registered varieties are shown below (See Table 2).

NOTE: Do not make applications less than four weeks before bloom. IT IS IMPORTANT that the proper rate be used on each variety; if late applications are made or if indicated rates are exceeded, reduction in yield may occur during the year of application and subsequent years.

130/40

TABLE 2

TABLE FOR APPLICATION RATES (GRAMS A.I./ACRE) AND RECOMMENDED WATER VOLUME ON SEEDED WINE VARIETIES.

Variety	PPM	Grams a.i./acre	Gallons/acre
Palomino Sauvignon Blanc Tinta Madeira	1-to-2.5	0.4-to-1	100
Aleatico Carignane Chardonney Chenin Blanc French Colombard Pinot Noir Valdepenas	2.5-to-5	1-to-2	100
Barbera Petite Sirah Zinfandel	5-to-10	2-to-4	100
Green Hungarian	10-to-20	4-to-8	100
Grenache Alicante	20	8	100
Salvadore	20-to-40	8-to-16	100

CITRUS

For all citrus, apply in sprays of sufficient water volumes to insure thorough fruit wetting. Application to trees of low vigor or under stress, (pest, nutritional, or water, etc) may cause severe leaf and/or fruit drop. In most cases some drop of older mature leaves will occur after application. Do not apply in white wash sprays in which lime or other caustic material has produced a high pH in the spray tank.

NAVEL ORANGE

To delay rind aging, reduce physiological disorders (e.g., rind staining, water spotting, sticky or tacky surface, puffy rind and rupture under pressure), and to produce a more orderly harvesting pattern. The delay in rind aging is greatest when an early spray is applied. This spray timing produces the firmest rind possible.

Guide: Apply 16-to-48 grams a.i./acre as a concentrate or dilute spray in sufficient water volume to insure thorough wetting.

EARLY SPRAY: Apply one spray approximately two weeks prior to color break, which normally occurs August through November.

140/40

OR

LATE SPRAY: Apply one spray after marketable color has developed, normally from October through December.

This late application may cause fruit re-greening.

NOTE: Do not apply the early spray to groves that may be harvested early, as fruit coloring will be delayed. Do not apply from January through July, as production may be reduced the following year.

VALENCIA ORANGE

To reduce rind creasing and to delay rind aging and softening.

Guide: Apply a single spray in August to October to trees with a target crop of young fruit. Apply 40-to-80 grams a.i./acre as a concentrate or dilute spray in sufficient water volume to insure thorough wetting.

NOTE: Slower color development should be expected in the target crop. Increased re-greening of mature fruit may occur. After marketable color is achieved, treatment effects may be reduced the longer treated fruit remain on the tree.

OTHER ROUND ORANGES (All States except California)

To reduce rind creasing and to delay aging and softening of the rind.

Guide: Apply a single spray in August to October to trees with a target crop of young fruit. Apply 40-to-80 grams a.i./acre as a concentrate or dilute spray in sufficient water volume to insure thorough wetting.

NOTE: Slower color development should be expected in the target crop. Increased re-greening of mature fruit may occur. After marketable color is achieved, treatment effects may be reduced the longer treated fruit remain on the tree.

LEMON/LIME

To decrease the amount of small ripe fruit and to produce a more desirable production pattern relative to market demand.

Guide: Apply one spray when target crop is 1/2-to-3/4 full size, but still green. Use 10-to-32 grams a.i./acre as a concentrate or dilute spray in sufficient water volume to insure thorough wetting.

When applied two years in a row, an even larger difference in harvest pattern and maturity occurs.

TANGERINE HYBRIDS

To delay disorders associated with rind aging, puffiness, and softening, and to increase peel strength, of tangerine hybrids such as Orlando, Robinson, Minneola and Sunburst.

Guide: Apply 20-to-40 grams a.i./acre approximately two weeks prior to color break. Apply as a dilute spray in sufficient water volume to insure thorough wetting.

NOTE: Do not apply if early harvest is planned. Do not apply after coloring as preharvest rind staining may occur. Application during coloring may cause variation in rind color development.

(All States except California)

To increase fruit set and yields on tangerine hybrids with pollination problems such as the Orlando, Robinson, Minneola and Sunburst.

Guide: Apply 8-to-30 grams a.i./acre during full bloom. Apply as a dilute spray in sufficient water volume to insure thorough wetting.

NOTE: Fruit sizes may be reduced and color development slightly retarded. A slight increase in mature leaf drop may occur in trees under stress.

GRAPEFRUIT

(All states except California)

To delay disorders associated with rind aging (e.g., puffiness, softening, and orange coloration), to prevent preharvest drop of mature fruit, to increase peel strength and reduce water loss during storage, and to produce a more orderly harvesting pattern. The delay in rind aging is greatest when an early spray is applied before color change. This spray timing produces the firmest rind possible.

Guide: Apply 16-to-48 grams a.i./acre in a minimum of 250 gallons per acre.

EARLY SPRAY: Apply one spray approximately two weeks prior to color break, which normally occurs August through September.

OR

LATE SPRAY: Apply one spray after marketable color has developed which is normally

160440

from October through December.

This late application may cause fruit re-greening.

NOTE: Do not apply the early spray to groves that may be harvested early as fruit coloring will be delayed. It is advisable to spot pick heavy crops to aid early marketing and to avoid reduction of yields, which generally follow late held crops. Fully colored fruit to which applications have been made will begin to re-green if allowed to remain on the tree for extended periods.

Application made after December, or when trees begin to break dormancy, may adversely affect new crop. Do not use concentrate sprays. Results may vary from season to season depending on environmental conditions.

STAR RUBY VARIETY
(All States except California)

- To reduce early-season small fruit drop of Star Ruby Variety thereby increasing yields.

Guide: Apply a single spray during the bloom period. Use 25 grams a.i./acre in a minimum of 250 gallons of water per acre.

NOTE: Results may vary from season to season depending on environmental conditions. Maintain a well-balanced fertilization and watering program.

FRUIT CROPS

BLUEBERRY

(All States except California)

To improve fruit set. For natural fruit set problems due to insufficient natural honeybee pollination, adverse weather conditions, or physiological factors.

Highbush blueberry - (for varieties such as Coville, Jersey, Stanley, Earliblue, Weymouth, Walcott, Berkeley, Bluecray, Bluecrop, 1316A, Concord, and others)

Guide: Make one or two applications at 40 grams a.i./acre in 100 gallons of water; the single application should be made at full bloom (when 75% of the flowers are fully open). When 2 applications are made, spray the first one at full bloom, and the second one within 10-14 days of the first one. For Weymouth, application can be delayed up to two weeks after bloom to increase size of "shot" berries.

Rabbiteye blueberry - (for varieties such as Aliceblue, Beckyblue, Bonita, Brightwell, Climax, Delite, Tiftblue, Woodward, and others).

Guide: Make a single application of 40 grams a.i./acre in 100-to-300 gallons of water per acre when most of the flowers are elongated but not yet open (bloom Stage 5).

OR

MULTIPLE APPLICATIONS: Make two to four applications 10-to-14 days apart starting at bloom Stage 5. Spray 40 grams a.i./acre in 50-to-300 gallons of water per application.

SWEET CHERRY

To produce larger, brighter colored, firmer fruit.

Guide: Apply a single spray when the fruit is light green to straw colored. Use 16-to-48 grams a.i./acre in sufficient water volume to insure thorough wetting.

NOTE: Color development and harvest may be slightly delayed.

180/40

RED TART CHERRY
(All states except California)

To maintain and extend high fruiting capacity of bearing tart cherry trees and reduce the occurrence of "blind" nodes. Treatment will cause bud differentiation, which is apparent the year after application. Therefore, changes in shoot, spur, and flower production will not be evident until two or three years after program initiation. Applications must be applied annually to insure vegetative development and subsequent yield improvement year after year.

Guide: Apply one spray 14-to-28 days after bloom. Optimum timing is defined as that stage when 3-to-5 terminal leaves have fully expanded, or, at least 1-to-3 inches of terminal shoot extension has occurred. Use 4-to-18 grams a.i./acre, depending on tree age and vigor (See Table 3). Apply as a concentrate or dilute spray in sufficient water volume to insure thorough wetting.

TABLE 3

**RECOMMENDED APPLICATION RATES (GRAMS A.I./ACRE)
FOR TART CHERRY TREES BY AGE**

Tree Age (years)	Rate (grams a.i./acre)
6-to-10	4-to-6
11-to-15	8-to-10
16-to-20	10-to-14
20 + years	14-to-18

NOTE: Rates are based on expected normal tree vigor at various ages. Adjust rate according to tree vigor. If trees are vigorous, use lowest recommended rates. Lowest rates should also be used on trees that have been heavily pruned or hedged. Use higher rates for trees low in vigor and weak in shoot and spur production. Excessive application rates will increase vegetative growth at the expense of fruit production the following year.

190440

RED TART CHERRY

(continued)

Applications will not improve growth of trees under stress conditions, such as nutritional, moisture, or pest. Best results will be obtained when combined with good cultural practices.

ITALIAN PRUNE

(All states except California)

To reduce internal browning, improve quality, and increase size.

Guide: Apply 4-to-5 weeks before expected harvest. Apply a single spray at 16-to-48 grams a.i./acre in sufficient water volume to insure thorough wetting.

NOTE: Color development and harvest may be slightly delayed. May reduce bloom the following season.

20940

NON-BEARING FRUIT TREES

To reduce flowering and fruiting in young tart and sweet cherry, and peach trees in order to minimize the competitive effect of early fruiting on tree development.

NOTE: DO NOT SPRAY TREES IN THE FIRST YEAR. Treat in the second season for reduction of flowering in the third season, and again in the third season if flower reduction and fruiting is desired in the fourth season. Treat only trees that are in good physiological condition. Discontinue treatment the year before desired harvest.

YOUNG TART AND SWEET CHERRY (All states except California)

Guide: Apply a single spray of 20-to-40 grams a.i./acre, 2-to-4 weeks after bloom. Apply a foliar spray of 25-to-50 gallons per acre, assuming a tree density of 100 trees per acre equivalent. Under conditions of low vigor, two applications are recommended. If two spray applications are made, allow at least a seven-day interval between sprays.

PEACH (All states except California)

Guide: Apply a single spray in the fall after flower buds have been initiated. This corresponds to the period immediately before and at the onset of early leaf drop, typically late September to early October. Apply at the rate of 40-to-80 grams a.i./acre in 50 gallons of water per acre. Best results are obtained when applied with a handgun and tree canopy is wetted thoroughly to the point of run-off.

21 of 40

OTHER FRUIT

**OLYMPUS STRAWBERRY
(All states except California)**

To increase runner production of mother plants of the Olympus cultivar.

Guide: Apply a single spray of 20 grams a.i./acre to mother plants 10-to-30 days after planting. At the time of spraying, plants should have 1-to-6 leaves. Apply 100 gallons/acre to thoroughly wet new foliage to the point of run-off.

NOTE: Not for use on fruiting plants. Treatments may not be effective on plantings set out after mid-May.

VEGETABLE CROPS

RHUBARB

To break dormancy on plants receiving insufficient chilling and to increase marketable yield of forced rhubarb.

Guide: Make a single application of 2 fluid ounces (60 ml) of a solution containing 20 grams a.i. in 10 gallons of water to each cleaned crown, when the rest period is not completely broken. When the rest period is broken by cold weather, apply 2 fluid ounces (60 ml) of a solution containing 10 grams a.i. in 10 gallons of water to each cleaned crown.

NOTE: Keep forcing house temperatures at 40 F-to-50 F for 24 hours after application. If house is warmer than 50 F, the crowns should be covered with plastic. Temperatures in the forcing house above 50 F may lower yields and cause poor stalk color.

ARTICHOKE

To accelerate maturity and shift harvest to an earlier date.

Guide: For perennials, apply one to three applications at bud initiation stage. For annuals, apply one to four applications at 2 week intervals, beginning at the fourth true leaf. Use 10-to-20 grams a.i./acre per application in sufficient water volume to insure thorough wetting of the entire plant (leaves, stems and buds).

CELERY

To increase plant height and yield and overcome stress due to cold weather conditions or saline soils, and to obtain earlier maturity.

Guide: Apply a single spray one to four weeks prior to harvest at a rate of 2.5-to-10 grams a.i./acre. Use 25-to-50 gallons of water per acre by ground application or 5-to-10 gallons of water per acre for aerial application*. Use lower concentrations applying 3-to-4 weeks before harvest and higher concentrations within 1-to-2 weeks before harvest.

*Do not apply by air in California.

NOTE: Do not apply earlier than 4 weeks before harvest as bolting (seed stalk formation) may occur.

LETTUCE FOR SEED

230440

- To obtain uniform bolting and increase seed production.

Guide: Apply one to four applications at 2 week intervals, beginning at the fourth true leaf. Use 1-to-4 grams a.i./acre per application in sufficient water volume to insure thorough wetting.

PEPPER (All states except California)

- To Promote Plant Growth

Guide: Apply one to two sprays of 1-to-3 grams a.i./acre in 25-to-50 gallons of water per acre at two week intervals. Begin sprays 2 weeks after transplanting.

NOTE: This use is recommended for areas with short growing seasons, or when low temperatures slow plant growth.

- To Increase Fruit Set and Promote Fruit Growth

Guide: Apply one to two sprays of 1-to-3 grams a.i./acre in 25-to-50 gallons of water per acre at weekly intervals during the flowering period. The high rate is recommended for areas and/or varieties with pollination and/or fruit set problems.

- To Increase Fruit Size

Guide: Apply 1-to-3 grams a.i./acre in 25-to-50 gallons of water per acre at the beginning of the picking period. The high rate is recommended for plants with heavy fruit loads.

MELON AND CUCUMBER (All states except California)

- To stimulate fruit set during periods of cool temperatures.

Guide: Use 2 grams a.i./acre in sufficient water volume for thorough coverage of exposed foliage. Make one application prior to bloom followed by two additional applications at intervals of 10-to-14 days on cantaloupes and watermelons. On cucumbers, up to four applications may be required.

For maximum benefits, vines must be in good condition, except for reduced rate of growth due to cool temperatures.

SEED POTATO

p984%01.sam/21
rev. 05/04/98

240440

To stimulate uniform sprouting to aid in maximum production, more uniform development, fewer late maturing plants, and to break dormancy of newly harvested potatoes that have not had a full rest period.

Guide: Dip whole or cut seed pieces in a solution containing 0.2-to-0.4 grams a.i. in 100 gallons of water prior to planting.

NOTE: Under high soil temperatures use the minimum concentration for dormant seed. Do not treat rested seed pieces.

SPINACH (All states except California)

To facilitate harvest, increase yield and improve quality of fall and over-winter spinach.

Guide: Apply a single spray 10-to-18 days before each anticipated harvest on fall or over-winter spinach, ideally when daytime temperatures are 40° F-to-70° F and during early morning hours when dew is present on crop. Apply 6-to-8 grams a.i./acre in 10-to-50 gallons of water per acre by ground sprayer or in a minimum of 5-to-10 gallons of water per acre by air. When applied to promote growth of second cutting, wait until some regrowth has started before spraying. Maximum benefit is obtained when below normal temperatures predominate following application and growth would be otherwise slowed in untreated spinach.

NOTE: Since the promotion of bolting may occur, do not apply after the mid-winter period or if temperatures may be expected to exceed 75° F within several days of application. Do not apply on spring plantings.

250/40

FLORICULTURE CROPS

POMPOM CHRYSANTHEMUM (All States except California)

- For elongating peduncles on Pompom chrysanthemum.

Guide: Apply a single spray 4-to-5 weeks after initiation of short day conditions.

Use 0.5-to-1 grams a.i. in 12 gallons of water for application to 1,000 sq. ft. of bed.
Apply with overhead nozzles directing the spray to the flower buds.

NOTE: Overuse or incorrect timing may cause long, spindly, and weak stems.

STATICE

(All States except California)

- To promote earlier flowering and to increase flower yield.

Guide: Apply a single drench spray when plants are more than 10 inches in diameter (approximately 90-to-110 days after normal seeding time). Use 40-to-50 grams a.i. in 25 gallons of water to provide 10 ml solution (4-5 mg a.i.) per plant.

NOTE: Do not exceed specified rates. Do not apply repeated sprays. Accelerated flowering is influenced by extended photoperiod, adequate nutrition, and reduced night temperatures. This treatment reduces the cold requirement and/or the long photoperiod.

AZALEA

(All States except California)

The following recommendations are based on results with common azalea cultivars. Differences in responsiveness may vary from one cultivar to another, or from one set of growing conditions to another, or from one cultural management system to another. Therefore, prior to widespread usage, we recommend testing a small number of plants from each cultivar under a specific set of growing and cultural management conditions to verify desired efficacy.

Spray plants to run-off. The actual spray application rate will vary, depending on plant size and spacing density. Thorough spray coverage is essential for uniform flowering. Do not apply after flower buds show color.

NOTE: A representative spray application rate which has been proven effective for 6 inch potted plants spaced at a density of 1 per square foot is 1 gallon spray solution/200 square feet.

- As a partial replacement of cold treatment to break flower dormancy.

260440

Guide: Apply three sprays of 250-to-500 ppm a.i. (See Table 4) at weekly intervals after 3-to-4 weeks of chilling.

NOTE: Plants should be at Stage 5 of floral development (i.e., style elongated and open) when treatment is initiated. A representative spray schedule would consist of applications made at 3, 10, and 17 days after four weeks of chilling. Flowers will not develop properly if applied prior to Stage 5.

NOTE: On some cultivars (e.g., 'Gloria', 'Prize', and 'Redwing'), a single spray of 1000 ppm a.i. after 3-to-4 weeks of chilling has proven effective in breaking dormancy.

- As a complete substitution of cold treatment to break flower dormancy.

Guide: Apply four to six sprays of 1000 ppm a.i. (See Table 4) at weekly intervals. Plants must be at Stage 5 of floral development (style elongated and open) before first spray is applied.

NOTE: Flowers will not develop properly if applied prior to Stage 5 of floral development.

- To inhibit flower bud initiation during vegetative growth.

Guide: After each pinch, apply two to three sprays of 100-to-750 ppm a.i. (See Table 4) at intervals of 2-to-3 weeks.

Table 4 can be used to convert spray concentrations (ppm of a.i.), to actual number of fluid ounces of ProGibb 4% needed for one gallon of spray solution.

27/2/40

TABLE 4**APPLICATION RATES AND RECOMMENDED WATER VOLUME
FOR AZALEA**

Desired ppm Value	grams a.i./ gallon*	grams a.i./ acre**	Fluid ounces of ProGibb 4%/acre**
100	0.38	87	87
250	0.95	207	207
500	1.90	414	414
750	2.85	610	610
1000	3.80	828	828

*Note: ProGibb 4% is a liquid. Each fluid ounce contains approximately one (1) gram of active ingredient.

**Based on a spray application rate of 1 gallon of spray solution/200 square feet.

280/40

GOLF/TURF**BERMUDAGRASS GOLF TURF
(All States except California)**

- To initiate or maintain growth and prevent color change during periods of cold stress and light frosts on golf course Bermudagrass (e.g., Tifdwarf, Tifgreen, etc.).

Guide: Apply 10 grams a.i./acre weekly or 25 grams a.i./acre biweekly in 25-to-100 gallons of water/acre.

NOTE: Do not exceed specified rates. Do not apply during extended warm period where night temperatures exceed 65 F.

- To maintain or enhance regrowth during summer months.

Guide: Apply 1-to-3 grams a.i./acre weekly in 25-to-100 gallons of water/acre.

NOTE: Maintain adequate moisture and proper fertilization programs recommended in local area. Discontinue treatments if thinning is observed. Do not apply the high rate more frequently than every two weeks. More frequent mowing may be necessary. Do not use on dormant turf.

29 of 40

OTHER CROPS

HOP

(Northwestern U.S. only)

For seeded and seedless Fuggle hop and similar varieties adapted to the Northwestern states.

To increase yield and fruit set.

Guide: Apply a single spray when vine growth is 5-to-8 feet in length. Use 4-to-6 grams a.i./acre in 100-to-150 gallons of water/acre.

COTTON

(All states except California)

To promote early plant growth, increase early seedling vigor, and to overcome stress caused by cool weather.

Guide: Apply 1-to-6 grams a.i./acre via in-furrow application to seed, or as a foliar application from the cotyledon stage through the 5 leaf stage. Repeat applications as needed every 5-to-7 days, to a maximum of 4 applications. Use 5-to-40 gallons of water by ground application or 3-to-10 gallons by air.

NOTE: Use higher rates when temperatures will likely average 75° F or less during the 14 days following the applications. Do not tank mix with herbicides. Do not apply more often than necessary to achieve the desired height, as overdosage may result in excessive growth.

308140

CONVERSION TABLE

GRAMS OF GIBBERELIC ACID PER ACRE	TO	AMOUNT OF PROGIBB 4% FORMULATION PER ACRE
Desired Gibberellic Acid Concentration (Grams Active Ingredient per acre) in Finished Spray		ProGibb 4% Liquid Contains approximately 1.0 Gram Active Ingredient/ Fluid Ounce of Formulated Product
0.5		0.5 oz.
1.0		1 oz.
2.0		2 oz.
4.0		4 oz.
5.0		5 oz.
8.0		8 oz.
10.0		10 oz.
12.0		12 oz.
16.0		16 oz.
20.0		20 oz.
25.0		25 oz.
32.0		32 oz.
40.0		40 oz.
48.0		48 oz.
50.0		50 oz.

NOTICE TO USER:

Seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance with accompanying directions.

©1998

p984%01.ssm/28
rev. 05/04/98

31 of 40

SUB LABEL

RyzUp Plant Growth Regulator Solution

For use on rice (foliar and seed treatments); hybrid rice seed production; cotton; and turf (golf).

32040

RYZUP®

PLANT GROWTH REGULATOR SOLUTION

Active Ingredient:

Gibberellic Acid.....4.0% w/w

Inert Ingredients.....96.0% w/w

Total.....100.0% w/w

RyzUp liquid contains approximately 1.0 gram active ingredient per fluid ounce of formulated product.

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES - Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get medical attention.

IF SWALLOWED - Call a doctor or get medical attention. Do not induce vomiting or give anything by mouth to an unconscious person. Drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol.

IF INHALED - Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

IF ON SKIN - Wash with plenty of soap and water. Get medical attention.

EPA Reg. No. 275-61

EPA Est No:

Abbott Laboratories
Chemical and Agricultural Products Division
1401 Sheridan Road
North Chicago, IL 60064

Net Contents: _____

This container will treat ___ acres at the maximum use rate, as recommended for use on _____.

p97ryz02.ssm/30
rev. 5/4/98

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

WARNING

Causes substantial but temporary eye injury. Harmful if inhaled or absorbed through skin. Do not get in eyes or on clothing. Avoid breathing vapor or spray mist, and avoid contact with skin. Wear protective eyewear (goggles or face shield). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long sleeved shirt.
- Long pants.
- Chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, and viton.
- Shoes plus socks.
- Protective eyewear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or

340440

disposing of equipment washwaters. Exposed treated seed may be hazardous to birds and other wildlife. Dispose of all excess treated seed and seed packaging by burial away from bodies of water.

PHYSICAL OR CHEMICAL HAZARDS

FLAMMABLE! Keep away from heat and open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

EXCEPTION: If the product is soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls.
- Chemical resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride, and viton.
- Shoes plus socks.
- Protective eyewear.

350/40

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter without appropriate protective clothing until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Keep containers tightly closed when not in use. Keep away from heat and open flame.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL INFORMATION

IMPORTANT: Before application, read directions thoroughly and use only as directed.

RyzUp is an extremely potent plant growth regulator. Consult your local experiment station specialist, distributor, or the Abbott agricultural specialist in your area for the spray schedule best suited to your conditions.

RICE**1) FOLIAR APPLICATION**

Early season foliar applications of RyzUp may promote vigorous and more uniform seedling growth of rice prior to permanent flood establishment. This may permit earlier flooding (5 to 10 days earlier) of drill or broadcast-seeded rice and is particularly effective on semi-dwarf varieties. Early flooding may reduce the additional flushing costs associated with a delay in establishing the permanent flood, reduce weed infestations and the number of herbicide applications, and/or promote earlier and more uniform grain maturity.

Late season foliar applications of RyzUp between split-boot and 100% heading may increase panicle height of rice. This may facilitate harvest efficiency in the field by allowing the rice grain to be cut above the leaf canopy at faster combine speeds and at reduced vegetative load. Grain quality and maturity may be advanced with the promotion of tiller panicle development.

p97ryz02.samv33
rev. 5/4/98

360840

Heading applications to the first crop may also accelerate regrowth of second crop rice. This may result in earlier second crop maturity and maximized grain yield.

Timing and Rate Recommendations

Seedling Applications (Early Season)

RyzUp may be applied at a rate of 1 to 3 fl oz (30 to 90 ml) of product per acre to rice between the 1-2 and 4-5 leaf stages of growth. Timing and dosage should be based upon environmental conditions, tank mix combinations with herbicides, and preferred permanent flood practice in relation to rice leaf stage.

For best results, apply RyzUp at a rate of 1 to 2 fl oz (30 to 60 ml) of product per acre using either a non-ionic surfactant known to be non-phytotoxic to rice or in tank mix combination with rice herbicides (see Compatibility with Other Chemicals section). Use higher rates of 1.5 to 3 fl oz (45 to 90 ml) of product with some dry and water-based herbicide formulations, or when temperatures will likely average 75° F or less during 14 days after application.

NOTE: Foliage may temporarily appear lighter green in color due to accelerated growth rates following RyzUp application.

Panicle Extension Applications (Late Season)

RyzUp may be applied at a rate of 3 to 8 fl oz (90 to 240 ml) of product per acre between split-boot and 100% panicle heading to promote main culm and tiller panicle extension. Tank mix with a non-ionic surfactant known to be non-phytotoxic to rice.

Use Precautions

Avoid drift or accidental application to other crops.

Do not apply when rice is subjected to drought stress conditions.

Mixing Instructions

Fill the treatment tank with half of the final tank mix volume. Add the required amount of RyzUp and mix thoroughly while adding water to the desired final volume. Dispose of any unused spray material at the end of the day.

Application Equipment

RyzUp may be applied by aerial or ground spray equipment. As an aerial spray, use a spray system capable of producing a uniform spray pattern of medium to fine spray droplets at 10 gallon per acre (GPA). Apply no less than 3 GPA of total spray volume. Low pressure ground sprayers equipped with boom and flat fan nozzles using 10 to 15 GPA spray volume may be used.

Compatibility with Other Chemicals

RyzUp can be tank mixed with most commonly used rice herbicides and fungicides. When applying RyzUp in mixture with Arrosolo®, Riverside Propanil® 60DF, Stam® M4, Stam® 80EDF, or Wham!® EZ, plus one of their recommended adjuvants, use of an additional surfactant

37/02/40

is not necessary. Do not apply RyzUp with Whip[®] IEC or Whip[®] 360.

2) SEED TREATMENT APPLICATION

RyzUp may be used as a seed treatment for rice. RyzUp stimulates seed germination and promotes faster and more uniform stand establishment. The following table describes RyzUp application and expected benefits.

RYZUP SEED TREATMENT APPLICATION

Crop	RyzUp Use Rates	Important Considerations	Benefits
Rice	0.5 to 2.1 fl oz product in 8-20 fl oz water/100 lbs seed (Equivalent to 15 to 62 ml in 237 to 591 ml water/45 kg seed)	For use with drill or broadcast seeding systems. Do not apply RyzUp prior to a 24 hour presoak or to water used for the presoak. Do not exceed 2.1 fl oz product/100 lbs of seed (or 62 ml product/45 kg seed)	May promote germination and emergence for semi-dwarf and tall varieties. May help increase final stand density and uniformity when seed are planted deeper to receive adequate moisture.

Mixing Instructions

RyzUp may be applied to seed with standard mist treating equipment. For best results, higher treatment volume of 12 to 20 fl oz per 100 pounds of seed (355 to 591 ml/45 kg seed) ensures complete and uniform coverage.

Fill the treatment tank with half of the final tank mix volume. Add the required amount of RyzUp and mix thoroughly while adding water and other co-applied seed treatment products (see Compatibility with Other Chemicals section) to the desired final volume.

An approved dye must be added to distinguish RyzUp treated seed and prevent inadvertent use for food, feed, or oil purposes. Treated seed must be labeled in accordance with the requirements of the Federal Seed Act.

Use Restriction

Do not use treated seed for food, feed or oil purposes.

Compatibility with Other Chemicals

RyzUp is compatible with most commonly used fungicide seed treatments (e.g. Vitavax[®] CT and Dithane[®]), standard dyes, and sticker/binding agents. When preparing tank mixes, ensure adequate physical compatibility and mixing.

p97ryz02.sarr/35
rev. 5/4/98

HYBRID RICE SEED PRODUCTION

RyzUp may be applied during heading to increase panicle height of hybrid rice. This may facilitate pollination and harvest efficiency in the field, thus maximizing potential seed yield.

Timing and Rate Recommendations

For hybrid rice, 1 to 3 applications can be applied during heading to promote main culm and tiller panicle extension, thus helping to maximize flower pollination.

Do not apply more than a total of 100 fl oz (3000 ml) of product per acre.

Use Restrictions

Hybrid seed harvested from RyzUp treated rice may only be used for planting. Do not use the seed for food, feed or oil purposes. An approved dye must be added to distinguish the harvested hybrid seed and prevent inadvertent use for food, feed or oil purposes. All non-seed portions of the female plants and all pollinator (male) rice plants, including male seed, must be destroyed.

Use Precautions

Avoid drift or accidental application to other crops. Higher rates of RyzUp application to hybrid rice plants can result in excessive vegetative growth, thus producing a taller plant that is more prone to lodging.

Mixing Instructions

Fill the treatment tank with half of the final tank mix volume. Add the required amount of RyzUp and mix thoroughly while adding water to the desired final volume. Dispose of any unused spray material at the end of the day.

Application Equipment

Aerial application of RyzUp should be made with spray systems capable of producing a uniform spray pattern of medium to fine spray droplets. Apply no less than 3 gallons per acre (GPA) of total spray volume. Low pressure ground sprayers may be used equipped with boom and flat fan nozzles using 10 to 15 GPA spray volume.

COTTON

Early season foliar applications of RyzUp to young cotton plants may promote growth and increased seedling vigor, thus helping to overcome the vegetative growth "lag" phase. This benefit may reduce the time interval needed to develop optimum leaf area and plant height, thus maximizing the potential for earliness and improved yields.

Timing and Rate Recommendations

Apply 1 to 6 fl oz (30 to 180 ml) of RyzUp per acre via in-furrow application to seed, or as a foliar application from the cotyledon leaf stage through the 7 leaf/node stage. Repeat applications as needed to a maximum of 3 applications.

Use higher rates (within the indicated range) when temperatures will likely average 75°F or less

39 of 40

during the 14 days following application(s).

Use Precautions

Avoid drift or accidental application to other crops.

Do not apply more often than necessary to achieve the desired height, as overdosage may result in excessive vegetative growth.

Do not apply RyzUp to cotton plants that are under drought stress.

If the cotton plants are under continuous stress, application of RyzUp should be delayed until the stress is alleviated and the plants are beginning to recover.

Mixing Instructions

Fill the treatment tank with half of the final tank mix volume. Add the required amount of RyzUp and mix thoroughly while adding water to the desired final volume. Dispose of any unused spray material at the end of the day.

Application Equipment

RyzUp may be applied by aerial or ground spray equipment. As an aerial spray, use a spray system capable of producing a uniform spray pattern of medium to fine spray droplets at 10 gallon per acre (GPA). Apply no less than 3 GPA of total spray volume. Low pressure ground sprayers equipped with boom and flat fan nozzles using 10 to 15 GPA spray volume may be used.

Compatibility with Other Chemicals

Data regarding the compatibility of RyzUp with herbicides used in cotton are not available.

TURF (GOLF) BERMUDAGRASS

1) COOL WEATHER APPLICATION

RyzUp may be used to initiate or maintain growth and prevent color change during periods of cold stress and light frosts on golf course Bermudagrass (e.g., Tifdwarf, Tifgreen, etc.).

Timing and Rate Recommendations

Apply 10 fl oz (300 ml) of product per acre weekly, or 25 fl oz (750 ml) of product per acre biweekly in 25 to 100 gallons per acre (GPA).

Use Precautions

Do not exceed specified rates.

Do not apply these rates during extended warm periods where night temperatures exceed 65° F.

2) HOT WEATHER APPLICATION

RyzUp may also be used to maintain or enhance regrowth of golf course Bermudagrass (e.g., Tifdwarf, Tifgreen, etc.) during summer months:

p97ryz02.sam/37
rev. 5/4/98

40 of 40

Timing and Rate Recommendation

Apply 1 to 3 fl oz (30 to 90 ml) of product per acre weekly in 25 to 100 GPA.

Maintain adequate moisture and proper fertilization programs recommended in local area.

Use Precautions

Do not apply the high rate more frequently than every two weeks.

Do not use on dormant turf.

Discontinue treatments if thinning is observed. More frequent mowing may be necessary.

NOTICE TO USER

Seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. User assumes all risks of use, storage or handling not in strict accordance with accompanying directions.

Arrosolo is a registered trademark of Zeneca Ag.ICI Americas Inc.

Riverside Propanil 60DF is a registered trademark of Terra Inc.

Stam M4, Stam 80EDF and Dithane are registered trademarks of Rohm and Haas Co.

Wham! EZ is a registered trademark of RiceCo.

Whip IEC and Whip 360 are registered trademarks of AgrEvo.

Vitavax CT is a registered trademark of Uniroyal Chemical Company, Inc.

©1998