

ACCEPTED
OCT 17 10 22

PM17 275-67

275-67

NOT REGISTERED IN CALIFORNIA

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
(and Domestic Animals)**

CAUTION

Avoid contact with skin, eyes or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

Beneficial Insects

Honeybees foraging treated areas are not harmed by DIPEL BAF use.

DIPEL BAF does not interrupt the activities of beneficial and predaceous arthropods in pest management programs.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Refer to Package Enclosure for crops, rates and general directions.

**STORAGE AND
DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep containers tightly closed when not in use. At temperatures less than 0°F and greater than 100°F, DIPEL BAF should be stored under cover. Roll or shake the drum before dispensing.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

NOTICE TO USER

Seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning use of this product other than as indicated on the label. User assumes all risk of use, storage or handling not in accordance with accompanying directions.

List No. 5407-04-01

E.P.A. Registration No. 275-67

E.P.A. Est. No. 2-782-1A-1



Chemical and Agriculture
Products Division
Abbott Laboratories
North Chicago, IL 60064

Lot No.

01-3741-2/R2

**DIPEL 8BAF
WORM KILLER**

**AQUEOUS FLOWABLE
BIOLOGICAL INSECTICIDE**

Active Ingredient:

**Bacillus thuringiensis, var.
kurstaki**, 14,500 International Units of
Potency per mg (64 billion
International Units per gallon) 2.9%
Inert Ingredients 97.1%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

NET CONTENTS: 55 GALLONS

**AQUEOUS FLOWABLE
BIOLOGICAL INSECTICIDE**

Active Ingredient:
Bacillus thuringiensis, var. kurstaki,
14,500 International Units of Potency per
mg (64 billion International Units per
gallon) 2.9%
Inert Ingredients 97.1%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

EPA Registration No. 275-67
EPA Est. No. 33762-1A-1

CAUTION

**Avoid contact with skin, eyes or clothing.
In case of contact immediately flush eyes
or skin with plenty of water. Get medical
attention if irritation persists.**

BENEFICIAL INSECTS

*Honeybees foraging treated areas are not
harmed by DIFPEL 8AF use.*

DIFPEL 8AF does not interrupt the activities of beneficial and predatory arthropods in pest management programs.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

STORAGE AND DISPOSAL

See container label.

Days to harvest: There are no restrictions on applying DIFPEL 8AF up to the time of harvest.

DIRECTIONS FOR USE

DIFPEL 8AF is a highly selective insecticide for use against listed caterpillars (larvae) of lepidopterous insects. Close scouting and early attention to infestations is highly recommended. Larvae must eat deposits of DIFPEL 8AF to be affected. Always follow these directions:

- Treat when larvae are young (early instars) before extensive damage has occurred.
- Larvae must be actively feeding on treated exposed plant parts.
- Thorough spray coverage is needed to provide a uniform deposit of DIFPEL 8AF at the site of larval feeding.
- Under heavy pest population pressure, use the higher label rates and/or raise gallonage to improve spray coverage.
- If attempting to control a pest with a single spray, make the treatment when egg hatch is essentially complete, but before extensive crop damage occurs.
- If rain is anticipated, a spreader-sticker which has been approved for use on growing crops may be added to improve weather fastness of the spray deposits.

After eating a lethal dose of DIFPEL 8AF larvae stop feeding within the hour and will die within several days. Dying larvae move slowly, discolor, then shrivel, blacken and die.

DIFPEL 8AF may be applied in conventional ground or aerial equipment with quantities of water sufficient to provide thorough coverage of infested plant parts. The amount of water needed per acre will depend on weather, spray equipment, and local experience. Fill the mix tank or plane hopper with the desired quantity of water. Start the mechanical agitator to provide moderate circulation before adding DIFPEL 8AF. If a sticker is to be used, always add the sticker to the water prior to the addition of DIFPEL 8AF. Add the desired volume of DIFPEL 8AF to the tank or plane hopper and continue circulation. Include agitator water from nozzles. Maintain the agitator on while being sprayed. Do not use more DIFPEL 8AF than can be used in a 24-hour period. **CAUTION:** Rinse and flush spray equipment thoroughly following each use.