Code 663

ACCEPTED

1987 FEB 1

Under the Federal 1. Pungicide, and Roders. as amended, for the publication tedinfered fuger 506. 49 **Net Weight**

Kolodust Extra

Fungicide

EPA Reg. No. 279-42

Active Ingredient:	By Wt.
Sullur	69.0%
Inert Ingredients:	
Sulfur Fineness 99.7% thru 325 mesh.	100%

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS Hazards to Humans (& Domestic Animals)

inhalid of absorbed amough of. Caution
Harmful if swallowed Avoid breathing dust or spray mist.
Avoid contact with eyes, skin and clothing.

Wash thoroughly with soap and water effer handling. Remove contaminated clothing and wash before reuse.

Statement of Practical Treatment

If swallowed: Drink one or two glasses of water and induce vomiting by touching back of throat with linger. Do not induce vomiting or give anything by inouth to an unconscious person.

If In eyes: Flush with plenty of water. Get medical attention if irritation

If on skin: Wash with plenty of soap and water. Great medical

For Emergency Assistance Cali 716-735-3765

See other panels for additional precautionary information

PHYSICAL OR CHEMICAL HAZARDS

Keep away from heat, sparks, or flame. Do not smoke while applying this product.



FMC Corporation Agricultural Chemical Group 2000 Market Street Philadelphia Pennsylvania 19103

7/30/86-R

Page 1 of 3

ENVIRONMENTAL HAZARDS

Do not apply directly to water Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas for 24 hours unless appropriate protective clothing is worn.

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven tabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

CAUTION

Area treated with Kolodust* Extra on (insert date). Do not enter treated areas for 24 hours unless appropriate protective clothing is worn (insert State Department of Agriculture's re-entry interval. if more restrictive).

In case of accidental exposure to pesticide spray or dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.

STORAGE AND DISPOSAL

PestIcide Storage
Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat.
Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not containinal other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC collect: (716) 735-3765.

To confine spill: If liquid, dike surrounding area or absort with sand, cal litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposai

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning, if burned, stay out of smoke.

see page

This product contains Bentonite Sulfur. Bentonite Sulfur is made by the absorption of fused sulfur into Bentonite in the proportion of thirty parts sulfur and seventy parts Bentonite. A portion of the sulfur particles are very minute, some of them colloidal. The process of manufacture and the character of the sulfur particles impart to the product distinctive properties with respect to toxicity to certain fungi, persistence on plants, suspensibility in spray tank and safety to host plants.

Crop	To Control	Remarks
Apples Pear	Scab	Use 40 pounds of dust per acre. Dust thoroughly applying to both upper and lower leaf surfaces as well as the fruit. Make repeated applications frequently enough to keep new growth protected. During periods of protonged well weather, apply Kolodust Extra in the rain.
Peaches	Scab Brown Rot	
Chernes Piums Prunes	Brown Rot	
Strawberries		thin ten days to two weeks of an oil spray. I spraying and dusting materials containing

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user

of excessively high summer temperatures.

Sulfur should not be made just prior to or during periods

agree to the following conditions:
Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing

and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller

and are assumed by buyer at his own risk.
Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice. Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

.

Kolodust Extra Fungicide Code 663

Kolodust Extra is a dust formulation. This product contains bentonite sulfur. Bentonite sulfur is made by adsorption of fused sulfur into bentonite in the proportions of 30 parts sulfur and 70 parts bentonite. A portion of the sulfur particles are very minute, some of them colloidal. The process of manufacture and the character of the sulfur particles impart to the product distinct properties with respect to toxicity to certain fungi, persistence on plants, suspensibility in spray tank and safety to host plants.


Applications of spraying and dusting materials containing sulfur should not be made just prior to or during periods of excessibly high summer temperatures. Do not use within 10 to 14 days of an oil spray. Do not exceed 40 pounds Kolodust Extra per acre.

Apple, Pears: To control Scab -- Apply Kolodust Extra from green-tip through petal fall.

Peaches: To control Scab, Brown Rot -- Apply Kolodust Extra from pink bud through petal fall and again preharvest if conditions favor fruit rot.

Cherries, Plums, Prunes: To control Brown Rot -- Apply Kolodust Extra from popcorn stage through petal fall and again preharvest if conditions favor fruit rot.

Apply when rain begins or high humidity occurs prior to infection. Dust early applying to both upper and lower leaf surfaces as well as the fruit. Make repeated applications frequently enough to keep new growth protected. During period of prolonged wet weather, apply Kolodust Extra in the rain.



(