

Code 10512

Net Weight

Kolospray® Fungicide-Insecticide

For Agricultural or Commercial Use Only

EPA Reg. No. 279-47

EPA Est., 279-

Active Ingredient:	By Wt.
Sulfur.....	81.25%
Inert Ingredients:	18.75%
	100.0%

The product contains bentonite sulfur. Sulfur fineness 99.7% thru 325 mesh.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS Hazards to Humans (and Domestic Animals)

Caution
Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT


If **swallowed**: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If in **eyes**: Flush with plenty of water. Get medical attention if irritation persists.

If on **skin**: Wash with plenty of soap and water. Get medical attention.

See other panels for additional precautionary information.

For Emergency Assistance Call: (716) 735-3765



FMC Corporation
Agricultural Chemical Group
Philadelphia PA 19103

1/6/87-A
4/87

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Keep away from heat, sparks, or flame. Do not smoke while applying this product.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas for 24 hours unless appropriate protective clothing is worn.

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. An example of such information is given on written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

CAUTION

Area treated with Kolospray® fungicide-insecticide on (insert date). Do not enter treated area for 24 hours unless appropriate protective clothing is worn (insert State Department of Agriculture's re-entry interval, if more restrictive).

In case of accidental exposure to pesticide spray or dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.

STORAGE AND DISPOSAL

Pesticide Storage
Keep away from fire and sparks.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC collect: (716) 735-3765.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal
Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal
Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Application of dusting and spraying materials containing sulfur should not be made just prior to or during periods of excessively high temperatures. Do not make applications within two weeks of an oil spray.

Do not use in undiluted form.

Fill the spray tanks about two-thirds full of water. Start the engine so that the agitator is in motion and sift the Kolospray into the tank while the tank is being filled. Add other spray materials that are required after the Kolospray has been dispersed in the spray tank. Observe all cautions and limitations on labeling of all products used in mixtures.

Almonds: Brown Rot, Scab—Use 6 pounds per 100 gallons of water in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

Apples, Pears: To control Scab and Powdery Mildew—Use 6 to 8 pounds of Kolospray or 4 to 6 pounds of Kolospray plus 2 to 3 pounds of Polysulfide Compounds per 100 gallons of water. Spray in the delayed dormant (silver tipped), early pre-pink (green tip) and early cover stages of growth. Do not apply on pears after the fruit begins to turn down. Sulfur may injure certain apple varieties. Check plant tolerance prior to use. Repeat at 7 to 10 day intervals or as necessary to maintain control.

Apples Only: To control Powdery Mildew—Use 1 to 3 pounds per 100 gallons of dilute spray. Apply 3 pounds per 100 gallons at delayed dormant (silver tipped) through the early pre-pink (green tip) stages and 2 to 3 pounds per 100 gallons during full bloom and 1 to 3 pounds per 100 gallons from petal fall through period when new growth ceases. Sulfur may injure certain apple varieties. Check plant tolerance prior to use. Repeat at 7 to 10 day intervals or as necessary to maintain control.

Pears Only: As a fall clean-up spray for control of Blister Mite, European Red Mite and Twospotted Spider Mite—Use 6 pounds of Kolospray plus 6 pounds of Polysulfide Compound or 5 gallons of Liquid Lime Sulfur Solution per 100 gallons of dilute spray.

Beans: Powdery Mildew, Leaf Spot, Rust—Use 4 to 7 pounds per 100 gallons of water. Begin applications when first true leaves form or at first sign of disease. Repeat at 7 to 10 day intervals. Sulfur may injure certain varieties of pears.

Blackberries, Currants, Dewberries, Gooseberries, Huckleberries, Raspberries, Strawberries: To control Powdery Mildew—Use 2 pounds of Kolospray per 100 gallons of water. Apply when disease first appears and repeat at 7 to 10 day intervals or as necessary to maintain control.

Citrus: Citrus Flat Mite, Yuma Spider Mite, Sixspotted Mite, and Citrus Rust Mite—Use 4 to 5 pounds per 100 gallons of water (a minimum of 60 pounds per acre). Apply when mites appear and repeat as necessary. Do not apply sulfur within 60 days of an oil application. Do not apply sulfur just prior to or during periods of excessively hot weather.

Grapes: Powdery Mildew—Use 2 to 5 pounds per 100 gallons. Begin applications when shoots are 6 to 8 inches long and repeat at 7 to 10 day intervals. Do not apply to Concord or other sulfur sensitive varieties.

Peaches, Plums, Prunes, Nectarines, Cherries: Powdery Mildew, Cherry Leaf Spot, Brown Rot, Prune Rust—Use 6 to 10 pounds per 100 gallons of water. Peach Silver Mite—Use 6 pounds per 100 gallons of water. (By ground, apply a minimum of 200 gallons of water per acre.) Apply in the pre-bloom, bloom and post-bloom periods. Apply when mites appear and when diseases threaten during rain periods and repeat at 7 to 10 day intervals.

Pears: Powdery Mildew, Rust—Use 4 to 7 pounds per 100 gallons of water. Begin early in season before disease appears. Repeat at 7 to 10 day intervals.

Potatoes: Powdery Mildew—Use 3 to 4 pounds of Kolospray in sufficient water to cover one acre. Applying by aircraft, use a minimum of 8 gallons of spray per acre. With ground equipment, use a minimum of 10 gallons of spray per acre. Apply at the first sign of infection and repeat at three week intervals.

Tomatoes: Russet Mite—Use 7 pounds per acre in a minimum of 25 gallons of water by ground applicator or in a minimum of 5 gallons of water by aircraft. Apply when mites first appear and repeat as necessary.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.