

CODE 323

NET WEIGHT
6 POUNDS

KOLOFOG

FUNGICIDE - MITICIDE

EPA REG. NO. 279-49 ZA

ACTIVE INGREDIENT	BY WT
Sulfur	30.00%
INERT INGREDIENTS	70.00%
	<hr/>
	100.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

Applications of dusting and spraying materials containing sulfur should not be made just prior to, or during, periods of excessively high summer temperatures. Do not apply sulfur within 10 days to 2 weeks of an oil spray.

ENVIRONMENTAL AND DISPOSAL CAUTIONS

Do not apply when weather conditions favor drift from the areas treated. Do not apply where run-off is likely to occur. Apply this product only as specified on this label.

DISPOSE OF WASTE by burying in non-crop land away from water supplies. Empty bags should be buried with wastes or burned. Keep Out Of Smoke.

For Emergency Assistance Call: 716-735-3765

Dealer Should Sell in Original Packages Only.

WARRANTY: FMC warrants that this product at the time of sale by FMC, (1) conforms to the agreement statement on the label, and (2) is reasonably fit for the purpose set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY, EXPRESS OR IMPLIED.

0111

MANUFACTURED BY FMC

AGRI-CHEMICALS DIVISION

1000 W. 10TH ST. ST. LOUIS, MO 63102

KOLOFOG

This product contains Bentonite Sulfur which is made by the adsorption of fused sulfur into bentonite in the proportion of thirty parts sulfur and seventy parts bentonite. A portion of the sulfur particles are very minute, some of them colloidal. The process of manufacture and the character of the sulfur particles impart to the product distinctive properties with respect to toxicity to certain fungi, persistence on plants, suspensibility in spray tank.

DIRECTIONS FOR USE

(WESTERN UNITED STATES)

Fill the spray tank about two-thirds full of water. Start the engine so that the agitator is in motion, and then slowly add the proper amount of Kolofog while the tank is being filled. Keep the agitator running. Thorough agitation is necessary. Introduce other materials into tank after Kolofog. Apply immediately. DO NOT attempt to wash Kolofog through the screen, or to make it into a paste before adding it to the tank.

Observe all cautions and limitations on labeling of all products used in mixtures.

ALMONDS: Brown Rot, Shothole Fungus and Brown Mites — Use 6 pounds to each 100 gallons of water in pre-blossom and petal fall applications.

APPLES, PEARS: Scab - Use 6 pounds, or 4 pounds plus 1 1/2 to 2 1/2 pounds Polysulphide Compound or 1 1/2 gallons of a lime sulfur solution, to each 100 gallons of water in the preblossom and calyx sprays. In the cover sprays, where a sulfur fungicide is desired, use 6 pounds to each 100 gallons of water and apply at 10 to 14 day intervals as required to maintain control. **Powdery Mildew** - Use 3 to 4 pounds plus 1/4 pound of dry spreader to each 100 gallons of water. Apply in preblossom, calyx and first cover sprays. (This recommendation is made for sections of Pacific Coast areas where scab is not a factor.) **Pacific Spider Mites** - Use 2 pounds plus 1/4 pound of dry spreader to each 100 gallons of water in the calyx and first cover sprays, timing the applications not more than seven or eight days apart. **Rust Mites** - Use 2 pounds plus 1/4 pound of dry spreader to each 100 gallons of water when Mites first appear. (This recommendation also applies for spraying Prunes.) **Blister Mites** - Use 4 to 6 pounds with 1 1/2 to 2 gallons of a light medium summer emulsion, or with 3 pounds Polysulphide Compound, or 2 gallons of a lime sulfur solution to each 100 gallons of water in early fall when crop has been picked.

CHERRIES: Cherry Leaf Spot and Brown Rot — Use 4 to 6 pounds to each 100 gallons of water for the usual pre-blossom and petal fall applications following dormancy. For Cherry Leaf Spot, apply again as a postharvest spray. Two-spotted Spider Mites, Pacific Spider Mites and Rust Mites — Use 2 pounds plus 1/4 pound of dry spreader to each 100 gallons of water when Mites first appear.

GRAPES: Powdery Mildew — Use 4 pounds in 20 gallons of water per acre. Apply as a spray beginning when shoots are 6 to 8 inches long and repeat at 7 to 10 day intervals until blossom opens. Concord and other labrusca type grapes may be injured by sulfur.

PEACHES, PLUMS, PRUNES: Brown Rot — Use 4 to 6 pounds to each 100 gallons of water for pre-blossom and petal fall applications. Late fall applications are recommended for control of late blight on peaches.