

Code 30282

Net Contents

Dormant Quik-Mix Heavy

Insecticide

EPA Reg. No. 279-61 ZA

EPA Est., 279-

Active Ingredient:	By Wt.
Petroleum Oil	98.00%
Inert Ingredients:	2.00%
	100.00%

Classification--Dormant Heavy
Unsulphonated Residue--78% Minimum

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

Caution

Harmful if swallowed. Avoid inhalation of spray mist.

Avoid application of oil sprays just prior to or during hot weather, rainy weather or drying winds. Apply when temperature is above 40° F. Do not apply just prior to freezing weather. Use only when trees are in healthy condition, and when moisture conditions are sufficient. Avoid overspraying.

Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

For Emergency Assistance Call 1-800-725-3755

See Other Panels for Additional Precautionary Information.

FMC

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

PA 351

Environmental Hazards

Keep out of lakes, ponds, or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. To dispose of container, triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill, or bury in a safe place. Do not cut or weld this container. Do not reuse empty 28 gallon drum. Return it to drum reconditioner or destroy it by perforating or crushing. Consult Federal, State or local disposal authorities for approved alternative procedures.

Directions for Mixing

Use only in equipment with mechanical or paddle agitation. Be sure spray tank is clean. With enough water in tank to cover agitator shaft, add desired amount of this material to the tank with agitator in motion. Pump this mixture through the overflow and back into tank for about two minutes. If this mixture turns white, it indicates good emulsification. Fill the tank with water, adding other desired materials as tank is filling. Keep agitator running at all times.

Do not use if material does not emulsify. Do not use in undiluted form.

Apply the following recommended rates in sufficient water to thoroughly cover one acre. With ground application equipment on Apples and Pears use a minimum of 300 gallons. On stone fruits and nuts use a minimum of 200 gallons. On grapes for cane pruned vines use a minimum of 200 gallons. For cordon pruned or arbor trained vines use a minimum of 300 gallons of diluted spray per acre. When using concentrate spray equipment use a minimum of 50 to 90 gallons of diluted spray per acre.

Apples and Pears: San Jose Scale, Oyster Shell Scale, Brown Apricot Scale, Black Scale and aid in control of European Red Mite--Use 2 to 4 gallons per 100 gallons of water. (Concentrate spray 6 to 8 gallons per acre) or 2 to 3 gallons of Dormant Quik Mix Heavy in combination with 6 to 7 pounds of Polysulphide Compound or 5 to 6 gallons of Lime Sulphur Solution per 100 gallons of water. (Concentrate spray use 6 to 8 gallons of Dormant Quik Mix Heavy plus 6 gallons of Lime Sulphur Solution per acre. Apply in the dormant period.

EPA Reg. No. 279-61
Page 2 of 2

Apricots, Cherries, Nectarines, Peaches and Almonds: San Jose Scale, Oyster Shell Scale, Brown Apricot Scale, Black Scale and aid in control of European Red Mite—Use 3 gallons per 100 gallons of water (Concentrate spray use 6 gallons per acre). Apply in the fall, dormant, or delayed dormant period.

As an Adhesive: For sticking copper on Apricots, Cherries, Nectarines, Peaches, Prunes and Almonds—Use 1½ to 2 gallons per 100 gallons of water. Apply in fall, dormant, or delayed dormant periods. **Olive**—Use 2 gallons per 100 gallons of water. Apply at post-harvest and before new growth appears.

Prunes: San Jose Scale, Oyster Shell Scale, Brown Apricot Scale and aid in control of European Red Mite Eggs—Use 3 gallons of Dormant Quik Mix Heavy in combination with 6 to 7 pounds Polysulphide Compound or 5 to 6 gallons of Lime Sulphur Solution per 100 gallons of water (Concentrate spray use 6 gallons of Dormant Quik Mix Heavy plus 6 gallons of Lime Sulphur Solution per acre). Apply in the dormant or delayed dormant periods (Sacramento and Suisun Valley areas). Application preferred in pre-bloom when fruit buds show green tips. Certain varieties of prunes may be injured by oil sprays. Check tolerance to oil prior to treatment.

Grapes: Grape Mealybug—Use 4 gallons per acre in 200 to 400 gallons of spray combined with 2½ quarts of Parathion 4 Emulsifiable. Apply in the dormant period. Observe the warning directions on the Parathion 4 Emulsifiable label. Do not combine this mixture with sodium arsenite.

Walnuts: Frosted Scale, European Fruit Lecanium Scale, Calico Scale and aid in control of European Red Mite—Use 3 gallons per 100 gallons of water. Apply in the delayed dormant period. This treatment should only be used on trees that have not suffered from lack of moisture at anytime during the growing season. Further, oil should not be applied to orchards when soil moisture is low.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.