

CODE: 30466

50 POUNDS NET WEIGHT

DUSTING SULPHUR

DIRECTIONS FOR USE

On **APPLES, PEARS** for control of Scab, on **ALMONDS, CHERRIES, PEACHES, NECTARINES, PLUMS AND PRUNES** for the control of Brown Rot, Cherry Leaf Spot, Silver Mite and Prune Rust and on **APPLES, NECTARINES** and **PEACHES** for the control of Powdery Mildew use at the rate of 40 to 50 pounds per acre in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals. On Pears do not apply after the fruit begins to turn down. Sulphur may injure certain varieties of apples, check tolerance prior to using.

On **GRAPES** for control of Powdery Mildew use 8 to 15 pounds per acre. Begin applications when shoots are 6 to 8 inches long and repeat as necessary. On **COTTON**: For control of Strawberry Mite (Atlantic Mite) apply 35 to 40 lbs. per acre when mites first appear and repeat as necessary.

On **STRAWBERRIES AND BUSHBERRIES**: For control of Powdery Mildew use 10 to 20 pounds per acre. Apply when disease first appears and repeat as necessary.

On **CITRUS**: For control of Citrus Thrips, Citrus Flat Mite, Yuma Spider Mite, Six Spotted Mite and Citrus Rust Mite, use at the rate of 100 pounds per acre. Apply when pests first appear and repeat as necessary.

VEGETABLE AND FIELD CROPS: Apply recommended rate when pests or disease first appear and repeat as necessary. For disease control keep new growth protected.

FUNGICIDE

U.S.D.A. REGISTRATION NO. 270-72

ACTIVE INGREDIENT:	By Wt.
Sulphur.....	98.00%
INERT INGREDIENTS.....	2.00%
	100.00%

The material in this bag is guaranteed to pass 95% through a 325 mesh screen.

CAUTION

KEEP OUT OF REACH OF CHILDREN

270-72

W/W

On **BEANS**: Powdery Mildew, Rust, Thrips, Spider Mites—Apply 15 to 30 pounds per acre. For rust control start applications as soon as the first true leaves form and continue at weekly intervals until blossom time. May cause injury to some varieties. Check tolerance prior to application.

On **CARROTS**: Petrobia Mite — Apply 30 to 35 pounds per acre.

On **ONIONS**: Petrobia Mite — Apply 20 to 30 pounds per acre.

On **PEAS**: Powdery Mildew — Apply 15 to 20 pounds per acre.

On **TOMATOES**: Tomato Russet Mite — Apply 30 to 40 pounds per acre.

On **TURNIPS AND RUTABAGAS**: Powdery Mildew — Apply 15 to 20 pounds per acre.

On **ORNAMENTALS**: Rust, Powdery Mildew—Dust for complete coverage.

CAUTION

Application of dusting and spraying materials containing sulphur should not be made just prior to or during periods of excessively high temperatures. Do not make applications within two weeks of an oil spray.

On Citrus do not make an oil application within 60 days prior to or after a sulphur application.

Do not use on or allow this material to drift onto melons, squash, cucumbers, apricots or other plants which will not tolerate sulphur.

