

Code: 229

[279-78]

50 Pounds Net Weight

KOLOKILL[®]

DIRECTIONS FOR USE

May be applied with ground or airplane application equipment. Observe days interval between last application and harvest indicated by number in () following the crop.

APPLES (30), PEARS (30): Apple Maggot, Codling Moth, Plum Curculio, Powdery Mildew, Red-banded Leaf Roller, Scab. Use 40-60 pounds per acre. Make first application at petal fall and repeat at 7 to 10 day intervals as required to maintain control. If additional scab and mildew control is required within 30 days of harvest, use a suitable fungicide. Remove excess residues at time of harvest. Do not graze livestock on treated areas.

CHERRIES (2): Brown Rot, Cherry Fruit Fly, Leaf Spot. Use 40-60 pounds per acre. Apply when flies first emerge and repeat at weekly intervals. For disease control, this material must be applied before the disease appears.

FUNGICIDE - INSECTICIDE

EPA REGISTRATION NO. 279-78

ACTIVE INGREDIENTS:	BY WT.
Sulphur	68.40%
Lead Arsenate	15.00%
INERT INGREDIENTS:	16.60%
	100.00%

CONTAINS STANDARD LEAD ARSENATE
 Arsenic expressed as metallic 3.24
 Arsenic in water soluble form expressed as metallic 0.50
 Lead expressed as metallic 8.95

WARNING

KEEP OUT OF REACH OF CHILDREN

May be fatal if Swallowed. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using. Avoid breathing dust or spray mist.

This product contains 13.5 per cent bentonite sulfur. Bentonite sulfur is made by the adsorption of fused sulfur into bentonite in the proportions of thirty parts sulfur and seventy parts bentonite. A portion of the sulfur particles are very minute, some of them colloidal. The process of manufacture and the character of the sulfur particles impart to the product distinctive properties with respect to toxicity to certain fungi, persistence on plants, susceptibility in spray tank and safety to host plants.

USAGE CAUTIONS

WARNING - POISON

This material contains ARSENIC or combinations thereof for agricultural purposes only. Do not keep or use in a room where food is being manufactured or prepared or where powdered food is stored.

This product is toxic to wildlife. Birds and other wildlife in treated areas may be harmed. Keep out of lakes, ponds and streams. Do not pollute the water by cleaning of equipment or disposal of wastes. Apply to plants only as specified on the label.

Avoid contact with dusting and spraying materials containing Sulfur. Do not be made just prior to or during periods of excessively high temperatures or within 10 days to 2 weeks of air oil spray.


Do not allow this material to drift onto sulfur sensitive crops.

Do not apply when weather conditions favor drift from the area treated. Do not apply when runoff is likely to occur.


Not for use or storage in or around the home.

DISPOSE OF WASTES by burying at least 18 inches deep in non-crop lands away from water supplies. Do not reuse or burn container or burn with wastes.

For Emergency Assistance Call
716-735-3765



ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



LABEL

FOR INFORMATION OF THE USER, THE FOLLOWING INFORMATION IS
CONTAINED ON THE BAGS TO WHICH THIS LABEL WILL BE ATTACHED.
NUTRITIONAL INFORMATION: VITAMIN C (AS ASCORBIC ACID) 1000 MG.
PER 100 GMS. OF VITA-GUM. VITAMIN E (AS TOCOPHEROL) 1000 IU.
PER 100 GMS. OF VITA-GUM. VITAMIN K (AS MENAQUINONE) 1000
MCU. PER 100 GMS. OF VITA-GUM.



REINFORCED WITH METAL WIRE
FOR EXTRA STRENGTH
AND DURABILITY

