

Code: 30453

50 Pounds

CR 50 DUST

INSECTICIDE

U.S.D.A. REGISTRATION NO. 279-83

Niagara CR 50 Dust is a dust containing 50% Cryolite for use in the control of many insects affecting fruits and vegetables.

DIRECTIONS FOR USE

Use at the rate of 20-30 pounds per acre. Apply when insects first appear and repeat as necessary. May be applied with ground or airplane application equipment.

ON LETTUCE to control Alfalfa Looper, Corn Earworm and Cutworms. Do not apply after the seedling stage.

ON BEANS to control Diabrotica and Flea Beetles. Remove excessive residues by washing, brushing, or other effective means.

ON CUCUMBERS to control Striped Cucumber Beetle, Flea Beetle and Diabrotica Beetle. Remove excessive residues by washing, brushing or other effective means.

ON CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS and BROCCOLI to control Cabbage Worms, Diabrotica and Flea Beetles. On Broccoli, Brussels Sprouts, and Cabbage do not apply after the edible parts or the heads start to form. On Cauliflower do not apply after the seedling stage.

ON ASPARAGUS to control Asparagus Beetle. Do not apply during the cutting season.

ON MELONS to control Cucumber and Diabrotica Beetles. Remove excessive residues by washing, brushing, or other effective means.

ACTIVE INGREDIENT:	BY WT.
Sodium Fluoaluminate - - - - -	50.00%
INERT INGREDIENTS - - - - -	50.00%
	100.00%
Fluorine expressed as the element	27.63

USAGE CAUTIONS

This product is toxic to fish and ponds and streams. Apply this product according to label.

DISPOSE OF WASTES by burying in non-productive areas. Empty bags should be buried. Keep out of smoke.

This product is highly toxic to bees or residues on crops. Protective instructions on your Cooperative Agricultural Extension Service label.

CAUTION
KEEP OUT OF REACH OF CHILDREN

ACCEPTED

AUG 11 1969

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENT CIDE ACT
ON 8/11/69 10:51 AM REGISTERED
UNDER NO. 279-83 SUBJECT
TO ATTACHED COMMENTS

CR 50 DUST

50 Pounds Net Weight

INSECTICIDE

U.S.D.A. REGISTRATION NO. 279-83

ACTIVE INGREDIENT:	BY WT.
Sodium Fluoaluminate - - - - -	50.00%
INERT INGREDIENTS - - - - -	50.00%
	100.00%
Fluorine expressed as the element	27.6%

CAUTION
KEEP OUT OF REACH OF CHILDREN

USAGE CAUTION

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Apply this product only as specified on the label.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

ACCEPTED
AUG 11 1969
FUNGICIDE AND INSECTICIDE ACT
REG. NO. 279-83
LD UNDER NO. 279-83 SUBJECT
TO ATTACHED COMMENTS

Printed in U.S.A.

Code: 30453

CR 50

Niagara CR 50 Dust is a dust containing 50% Cryolite for use in the control of many insects affecting fruits and vegetables.

DIRECTIONS FOR USE

Use at the rate of 20-30 pounds per acre. Apply when insects first appear and repeat as necessary. May be applied with ground or airplane application equipment.

ON LETTUCE to control Alfalfa Looper, Corn Earworm and Cutworms. Do not apply after the seedling stage.

ON BEANS to control Diabrotica and Flea Beetles. Remove excessive residues by washing, brushing, or other effective means.

ON CUCUMBERS to control Striped Cucumber Beetle, Flea Beetle and Diabrotica Beetle. Remove excessive residues by washing, brushing or other effective means.

ON CABBAGE, CAULIFLOWER, BRUSSELS SPROUTS and BROCCOLI to control Cabbage Worms, Diabrotica and Flea Beetles. On Broccoli, Brussels Sprouts, and Cabbage do not apply after the edible parts or the heads start to form. On Cauliflower do not apply after the seedling stage.

ON ASPARAGUS to control Asparagus Beetle. Do not apply during the cutting season.

ON MELONS to control Cucumber and Diabrotica Beetles. Remove excessive residues by washing, brushing, or other effective means.

INSE

U.S.D.A. REGIST

ACTIVE INGREDIENT:

Sodium Fluoaluminate - - -


INERT INGREDIENTS - - - -

Fluorine expressed as th

CAUT

KEEP OUT OF REA

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ON TO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE OF THIS PRODUCT ARE SUBJECT TO THE TERMS OF THE LABEL.

WARRANTY - Niagara Chemical Division warrants that the product is as described on the label. Niagara Chemical Division does not warrant the results of use of this product. NIAGARA CHEMICAL DIVISION WARRANTS THE PRODUCT IS AS DESCRIBED ON THE LABEL. NO OTHER WARRANTY EXPRESSES OR IMPLIES.

DAMAGE - Damage to crops or other property caused by the use of this product is the responsibility of the user. Niagara Chemical Division does not warrant the results of use of this product. NIAGARA CHEMICAL DIVISION does not warrant the results of use of this product. NIAGARA CHEMICAL DIVISION does not warrant the results of use of this product.

EX USE FROM NON-REGISTERED AREAS - NIAGARA CHEMICAL DIVISION does not warrant the results of use of this product in non-registered areas. NIAGARA CHEMICAL DIVISION does not warrant the results of use of this product in non-registered areas. NIAGARA CHEMICAL DIVISION does not warrant the results of use of this product in non-registered areas.

MANUFACTURED IN U.S.A. BY



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, NEW YORK

