

Code: 216

# KOLODUST®

## DIRECTIONS FOR USE

Kolodust may be applied by aerial or ground equipment, unless otherwise noted. Thorough coverage, at recommended rate, is necessary for control.

**ALMONDS, CHERRIES, PEACHES, NECTARINES, PLUMS, PRUNES:** Brown Rot, Cherry Leaf Spot, Silver Mite, Prune Rust, Powdery Mildew (Nectarines and Peaches only). Use 40 to 50 pounds per acre. Apply in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

**APPLES, PEARS:** Scab, Powdery Mildew — Use 40 to 50 pounds per acre. Apply in pre-bloom, bloom and post-bloom periods. Check tolerance prior to using on Apples as sulphur may injure certain varieties. Do not apply to Pears after the fruit begins to turn down.

**CITRUS:** Citrus Thrips, Citrus Flat Mite, Yuma Spider Mite, Sixspotted Mite, Citrus Rust Mite — Use 100 pounds per acre. Apply when pests first appear and repeat as necessary to maintain control.

**BLACKBERRIES, BOYSENBERRIES, GOOSEBERRIES, LOGANBERRIES, RASPBERRIES, STRAWBERRIES:** Powdery Mildew — Use 10 to 20 pounds per acre. Apply when disease appears and repeat as necessary to maintain control.

**GRAPES:** Powdery Mildew — Use 8 to 15 pounds per acre. Begin applications when shoots are 6 to 8 inches long and repeat as necessary. Ground application equipment preferred.

## FUNGICIDE-INSECTICIDE

EPA REGISTRATION NO. 279-126 AA

### ACTIVE INGREDIENT:

Sulphur

### INERT INGREDIENTS

BE

### BY WT

84.00%

16.00%

100.00%

AVAILABLE

## CAUTION

### KEEP OUT OF REACH OF CHILDREN

### ENVIRONMENTAL AND DISPOSAL CAUTIONS

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Applications of sulfur should not be made first prior to or during periods of excessively high summer temperature. Do not apply within 2 weeks (60 days on Citrus) following an application of oil.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty bags should be buried with wastes or burned. Keep Out Of Smoke.

### VEGETABLES AND FIELD CROPS:

BEANS: Powdery Mildew, Bean Rust per acre. For Rust control start and continue at weekly intervals until check tolerance prior to using.

CARROTS: Petrobis Mite — Use 30 lb

COTTON: Strawberry Mite — Use 30 lb

ONIONS: Petrobis Mite — Use 30 lb

PEAS, BUTTERNUTS, TURNIPS: Powdery

TOMATOES: Tomato Russet Mite — Use

ORNAMENTALS (Asters, Hollyhock, Roses)

### DEALER SHOULD

TERMS OF SALE OR USE: On purchase to the following conditions:

WARRANTY: FMC warrants that the forms to the ingredient statement of set forth in the Directions for Use. SOLD AS IS. FMC MAKES NO

DAMAGES: Buyer's or user's negligence shall be limited to direct and shall not include incidental or be reported to FMC within 10 days or user waives any claim for such

EXCUSE FROM NON-CONTROLLABLE created or incurred by buyer or as a condition of purchase or liability arising out of the handling. No person is authorized to vary

0074W Revised

MANUFACTURED IN USA BY

**FMC CORPORATION**

AGRICULTURAL CHEMICAL DIVISION

MIDDLEPORT NY  
FRESNO CA

YAKIMA WA  
RICHMOND CA

**ACCEPTED**  
**31 JUL 1975**  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 279-126

**FMC** Chemicals

# KOLODUST®

Net Weight  
50 Pounds

## FUNGICIDE-INSECTICIDE

EPA REGISTRATION NO. 279-126 AA

### ACTIVE INGREDIENT:

Sulphur

### INERT INGREDIENTS

### BY WT

84.00%

16.00%

100.00%

## CAUTION

KEEP OUT OF REACH OF CHILDREN

### ENVIRONMENTAL AND DISPOSAL CAUTIONS

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Applications of sulfur should not be made first prior to or during periods of excessively high summer temperature. Do not apply within 2 weeks (60 days on Citrus) following an application of oil.

DISPOSE OF WASTES by burying in noncrop land away from water supplies. Empty bags should be buried with wastes or burned. Keep Out Of Smoke.

**VEGETABLES AND FIELD CROPS** Apply recommended rate when pests or disease first appear and repeat as necessary. For disease control keep new growth protected.

**BEANS:** Powdery Mildew, Bean Rust, Thrips, Spider Mites - Use 15 to 30 pounds per acre. For Rust control start applications as soon as first true leaves form and continue at weekly intervals until blossom time. May cause injury to some varieties. Check tolerance prior to using.

**CARROTS:** Petroselin Mite - Use 30 to 35 pounds per acre.

**COTTON:** Strawberry Mite - Use 35 to 40 pounds per acre.

**ONIONS:** Petroselin Mite - Use 20 to 30 pounds per acre.

**PEAS, RUTABAGAS, TURNIPS:** Powdery Mildew - Use 15 to 20 pounds per acre.

**TOMATOES:** Tomato Russet Mite - Use 30 to 40 pounds per acre.

**ORNAMENTALS (Asters, Carnations, Chrysanthemums, Hollyhock, ROSES):** Rust, Powdery Mildew - Dust for complete coverage.

### DEALER SHOULD SELL IN ORIGINAL PACKAGES ONLY

**TERMS OF SALE OR USE:** On purchase or use of this product buyer and user agree to the following conditions:

**WARRANTY:** FMC warrants that this product, as of the time of sale by FMC (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

**DAMAGES:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damages.

**EXCUSE FROM NON-CONTROLLABLE CONDITIONS:** Because of certain critical conditions created or incurred by buyer or user over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

**FMC Chemicals**

### USE

Apply in serial or ground application as noted. Thorough coverage is necessary.

**APPLES, PEACHES:** Brown Rot, Cherry Powdery Mildew (Peaches and Peaches only) - Apply during bloom and post-bloom periods. Repeat at 7 to 10 day intervals.

Use 20 to 30 pounds per acre. Apply in advance prior to using on Apples to Peers after the fruit begins to

Apply to Mites, Spider Mite, Citrus Rust first appear and repeat as neces-

**LEMONS, RASPBERRIES,** 20 pounds per acre. Apply when control.

per acre. Begin applications when dry. Ground application equipment

ACCEPTED

31 JUL 1975

THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
ECONOMIC POISON REGISTER  
UNDER NO. 279-126

Code: 216

# KOLOD

### DIRECTIONS FOR USE

FUNGICIDE-INSECTICIDE

Kolodust may be applied by aerial or ground equipment, unless otherwise noted. For coverage, at recommended rate, 15-20% for control.

**ALMONDS, CHERRIES, PEACHES, NECTARINES, PLUMS, PRUNES** Brown Rot, Leaf Spot, Silver Mite, Prune Rust, Powdery Mildew (Nectarines and Peaches) Use 40 to 50 pounds per acre. Apply in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

**APPLES, PEARS** Scab, Powdery Mildew - Use 40 to 50 pounds per acre. Apply in pre-bloom, bloom and post-bloom periods. Check tolerance prior to using on Apples as sulphur may injure certain varieties. Do not apply to Pears after the fruit has turned down.

**CITRUS** Citrus Thrips, Citrus Flat Mite, Yuma Spider Mite, Spotted Mite, Citrus Rust Mite - Use 100 pounds per acre. Apply when pests first appear and repeat as necessary to maintain control.

**BLACKBERRIES, BOYSENBERRIES, GOOSEBERRIES, LOGANBERRIES, RASPBERRIES, STRAWBERRIES** Powdery Mildew - Use 10 to 20 pounds per acre. Apply when disease appears and repeat as necessary to maintain control.

**GRAPES** Powdery Mildew - Use 8 to 15 pounds per acre. Begin applications when shoots are 6 to 8 inches long and repeat as necessary. Ground application equipment preferred.

ACTIVE INGREDIENT  
SULFUR  
INERT INGREDIENT

## CAUTION

### KEEP OUT OF REACH

#### ENVIRONMENTAL AND DISPOSAL

Do not contaminate water, soil, or equipment, or disposal of this product only as specified on label.

Applications of sulfur first prior to or during excessively high summer temperatures (not apply within 2 weeks of citrus) following an application of sulfur.

DISPOSE OF WASTES by burying in non-crop area. Empty bags should be buried with wastes or burned.

FMC CORPORATION

AGRI-CHEMICALS DIVISION

**ACCEPTED**  
**31 JUL 1975**  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 279126