

Code 216

Net Weight \_\_\_\_\_ Pounds

# Kolodust®

## FUNGICIDE-INSECTICIDE

EPA REG. NO. 279-126 ZA      EPA Est., 279-

Active Ingredient:	By Wt.
Sulphur .....	84.00%
Inert Ingredients: .....	16.00%
	<hr/> 100.00%

Kolodust® — FMC Trademark

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans (and Domestic Animals)

Caution

#### Environmental Hazards

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, State or local procedures. Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

**KEEP OUT OF REACH OF CHILDREN**


**CAUTION**

For Emergency Assistance Call 716—735-3765.



FMC Corporation  
Agricultural Chemical Group  
2000 Market Street  
Philadelphia Pennsylvania 19103

PA 3 82



Applications of sulphur should not be made just prior to or during periods of excessively high summer temperatures. Do not apply within two weeks (60 days on Citrus) following an application of oil.


Kolodust may be applied by aerial or ground equipment, unless otherwise noted. Thorough coverage, at recommended rate, is necessary for control.

**Almonds, Cherries, Peaches, Nectarines, Plums, Prunes:** Brown Rot, Cherry Leaf Spot, Peach Silver Mite, Prune Rust, Powdery Mildew (Nectarines and Peaches only)—Use 40 to 50 pounds per acre. Apply in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

**Apples, Pears:** Scab, Powdery Mildew—Use 40 to 50 pounds per acre. Apply in pre-bloom, bloom and post-bloom periods. Check tolerance of variety. Apples as sulphur may injure certain varieties. Do not apply to Pears after the fruit begins to turn down.

**Citrus:** Citrus Thrips, Citrus Flat Mite, Yuma Spider Mite, Sixspotted Mite, Citrus Rust Mite—Use 100 pounds per acre. Apply when pests first appear and repeat as necessary to maintain control.

**Blackberries, Boysenberries, Gooseberries, Loganberries, Raspberries, Strawberries:** Powdery Mildew—Use 10 to 20 pounds per acre. Apply when disease appears and repeat as necessary to maintain control.



**Grapes:** Powdery Mildew—Use 8 to 15 pounds per acre. Begin applications when shoots are 6 to 8 inches long and repeat as necessary. Ground application equipment preferred.

**Vegetables and Field Crops:** Apply recommended rate when pests or disease first appear and repeat as necessary. For disease control keep new growth protected.

**Beans:** Powdery Mildew, Bean Rust, Thrips, Spider Mites—Use 15 to 30 pounds per acre. For Rust control start applications as soon as first true leaves form and continue at weekly intervals until blossom time. May cause injury to some varieties, check tolerance prior to using.

**Carrots:** Petrobia Mite—Use 30 to 35 pounds per acre.

**Cotton:** Strawberry Mite—Use 35 to 40 pounds per acre.

**Onions:** Petrobia Mite—Use 20 to 30 pounds per acre.

**Peas, Rutabagas, Turnips:** Powdery Mildew—Use 15 to 20 pounds per acre.

**Tomatoes:** Tomato Russet Mite—Use 30 to 40 pounds per acre.

**Ornamentals (Asters, Carnations, Chrysanthemums, Hollyhocks, Roses):** Rust, Powdery Mildew—Dust for complete coverage.

### Dealers Should Sell in Original Packages Only.

**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions.

**Warranty:** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

(216-PA-3/82)