

16F3  
Pina 1  
279-126

ACCEPTED  
FEB 2 1987  
Under the Federal Insecticide, Fungicide, and Herbicide Act, as amended, for the pesticide registered under EPA Reg. No. 279-126

Net Weight

Code 216

**Kolodust**<sup>®</sup>

FUNGICIDE-INSECTICIDE

EPA REG. NO. 279-126 ZA      EPA Est., 279-

Active Ingredient:	By Wt.
Sulfur.....	84.00%
Inert Ingredients.....	16.00%
	100.00%

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas for 24 hours unless appropriate protective clothing is worn.

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

**CAUTION**

Area treated with Kolodust on (insert date). Do not enter treated areas for 24 hours unless appropriate protective clothing is worn (insert State Department of Agriculture's re-entry interval, if more restrictive).

In case of accidental exposure to pesticide spray or dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

**PRECAUTIONARY STATEMENTS**  
Hazards to Humans (& Domestic Animals)

**Caution** Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**Statement of Practical Treatment**

If swallowed: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

If on skin: Wash with plenty of soap and water. *Get medical attention*

For Emergency Assistance Call 716-735-3765

See other panels for additional precautionary information.

**PHYSICAL OR CHEMICAL HAZARDS**

Keep away from heat, sparks, or flame. Do not smoke while applying this product.

**FMC**

FMC Corporation  
Agricultural Chemical Group  
2000 Market Street  
Philadelphia Pennsylvania 19103

7/30/86 - R

**STORAGE AND DISPOSAL**

**Pesticide Storage**  
Keep away from fire and sparks.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and tie back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC collect: (716) 735-3765.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

**Pesticide Disposal**  
Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal**  
Completely empty bag into application equipment. Treat or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

(All page 3)

**Almonds, Cherries, Peaches, Nectarines, Plums, Prunes:** Brown Rot, Cherry Leaf Spot, Peach Silver Mite, Prune Rust, Powdery Mildew (Nectarines and Peaches only)—Use 40 to 50 pounds per acre. Apply in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

**Apples, Pears:** Scab, Powdery Mildew—use 40 to 50 pounds per acre. Apply in pre-bloom, bloom and post-bloom periods. Check tolerance prior to using on Apples as sulfur may injure certain varieties. Do not apply to Pears after the fruit begins to turn down.

**Citrus:** Citrus Thrips, Citrus Flat Mite, Yuma Spider Mite, Sixspotted Mite, Citrus Rust Mite—Use 100 pounds per acre. Apply when pests first appear and repeat as necessary to maintain control.

**Blackberries, Boysenberries, Gooseberries, Loganberries, Raspberries, Strawberries:** Powdery Mildew—Use 10 to 20 pounds per acre. Apply when disease appears and repeat as necessary to maintain control.

**Grapes:** Powdery Mildew—Use 8 to 15 pounds per acre. Begin applications when shoots are 6 to 8 inches long and repeat as necessary. Ground application equipment preferred.

**Vegetables and Field Crops:** Apply recommended rate when pests or disease first appear and repeat as necessary. For disease control keep new growth protected.

**Beans:** Powdery Mildew, Bean Rust, Thrips, Spider Mites—Use 15 to 30 pounds per acre. For Rust control start applications as soon as first true leaves form and continue at weekly intervals until blossom time. May cause injury to some varieties; check tolerance prior to using.

**Carrots:** Petrobia Mite—Use 30 to 35 pounds per acre.

**Cotton:** Strawberry Mite—Use 35 to 40 pounds per acre.

**Onions:** Petrobia Mite—Use 20 to 30 pounds per acre.

**Peas, Rutabagas, Turnips:** Powdery Mildew—Use 15 to 20 pounds per acre.

**Tomatoes:** Tomato Russet Mite—Use 30 to 40 pounds per acre.

**Ornamentals (Asters, Carnations, Chrysanthemums, Hollyhocks, Roses):** Rust, Powdery Mildew—Dust for complete coverage.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

**Dealers Should Sell in Original Packages Only.**

**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions:

**Warranty:** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Kolodust and FMC—FMC Trademarks

279-126  
Page 2 of 3

Kolodust Fungicide-Insecticide  
Code 216

Applications of sulfur should not be made just prior to or during periods of excessively high summer temperatures. Cannot apply within 14 days (60 days on citrus) following an oil application. Kolodust may be applied by aerial or ground equipment unless otherwise noted. Thorough coverage at recommended rates are necessary for control. Make applications frequently enough to keep new growth protected or during the early season when disease threatens because of rains or high humidity. Post-bloom application should be made at 7 to 10 day intervals or as necessary to maintain control.

ies

279-126  
Page 3 of 3