

**CODE**  
**49**

# ZIRAM SPRAY

## FUNGICIDE

**NET WEIGHT**  
**30 LBS.**

### DIRECTIONS FOR USE

To mix, fill the spray tank about 2/3 full of water, and with agitator running, slowly sift in the required amount of Z-C Spray. Finish filling the tank with water and let mixing continue in the tank until thoroughly mixed before applying. Continue agitation and apply immediately.

**ALMONDS:** Brown Rot, Shot-hole Fungus, Leaf Blight - Use 2 pounds per 100 gallons of water in pre-bloom, blossom and petal fall periods. Scab - Use 2 pounds per 100 gallons of water at petal fall and 5 weeks after petal fall. Do not use later than 5 weeks after petal fall.

**APPLES:** Scab - Use 1 to 1½ pounds per 100 gallons of water in pre-blossom, blossom and early cover sprays. Bull's Eye Rot (Oregon & Washington) - Use 1½ to 2 pounds per 100 gallons of water beginning 12 to 14 days after calyx stage and repeat as necessary in summer sprays. Perennial Canker (Oregon) - Use 1 to 1½ pounds per 100 gallons of water in pre-blossom and calyx sprays.

**APRICOTS:** Brown Rot, Shot-hole Fungus - Use 2 pounds per 100 gallons of water in pre-bloom, blossom and petal fall thru early cover sprays.

**CHERRIES:** Brown Rot, Blossom Blight - Use 1½ pounds per 100 gallons of water in popcorn, full blossom and petal fall stages. Leaf Spot - Use 1½ pounds per 100 gallons of water in petal fall and shuck fall stages and approximately 2 weeks after shuck fall. Do not apply within 7 days of harvest.

EPA Reg. No. 279-268

### ACTIVE INGREDIENT:

Ziram (zinc dimethyl dithiocarbamate) . . . 76.0%

INERT INGREDIENTS: . . . . . 24.0%

100.0%

Zinc expressed as metallic . . . 16.25%

## CAUTION

### KEEP OUT OF REACH OF CHILDREN.

May be harmful if swallowed or inhaled! May cause irritation of eyes, nose, throat and skin. Avoid contact with eyes, skin or clothing. Avoid breathing dust or spray mist. In case of contact, flush with plenty of water; for eyes, get medical attention.

For Emergency Assistance Call 716-735-3765

Keep away from fire and sparks.

### ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish and wildlife. Keep out of lakes, ponds or streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

**PEACHES:** Brown Rot, Peach Blight (Coryneum) Use 2 pounds in 100 gallons of water in pre-bloom, blossom, petal fall and cover sprays. Leaf Curl - Use 2 to 3 pounds in 100 gallons of water as a dormant application prior to bud swell. Remove excess residue by washing or wiping.

**PEARS:** Scab - Use 1 to 1½ pounds per 100 gallons of water in pre-blossom, blossom and early cover sprays. Storage Rots (Oregon) - Use 1 to 1½ pounds per 100 gallons of water in pre-harvest spray.

**CUCUMBERS, MELONS:** Scab, Anthracnose - Use 2 pounds per 100 gallons of water per acre at 7 day intervals when plants are small. Increase rate to 3 pounds per 100 gallons of water per acre after plants vine.

**BEANS (SNAP, LIMA):** Anthracnose - Use 2 pounds per acre in sufficient water for thorough coverage when plants are small. Increase to 3 pounds per acre on large plants. Make applications at 7 day intervals starting when true leaves appear. Do not apply to snap beans later than 4 days before picking.

**TOMATOES:** Anthracnose, Early Blight - Use 2 to 3 pounds per 100 gallons of water per acre at 7 to 10 day intervals starting 2 to 4 weeks after plants are set. Under severe disease conditions apply 3 pounds in 180 to 200 gallons per acre at 7 day intervals.

ACCEPTED

**ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.**



**Dealers Should Sell in Original Packages Only.**

**DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.**

**TERMS OF SALE OR USE:** On purchase or use of this product buyer and user agree to the following conditions:

**WARRANTY:** FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.**

**DAMAGES:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

**EXCUSE FROM NON-CONTROLLABLE CONDITIONS:** Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured in U.S.A. by:  
FMC CORPORATION  
AGRICULTURAL CHEMICAL DIVISION  
Middleport, N.Y. 14105

**FMC**  
Chemicals