

Code 411

PRECAUTIONARY STATEMENTS

Environmental Hazards

This product is toxic to fish and wildlife. Keep out of lakes, ponds or streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Physical or Chemical Hazards

Keep away from fire and sparks.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of container in an incinerator or landfill approved for pesticide containers or bury in a safe place away from water supplies. Consult Federal, State or local disposal authorities for approved alternative procedures.

To mix, fill the spray tank about 2/3 full of water and with agitator running, slowly sift in the required amount of Ziram Spray. Finish filling the tank with water and let mixing continue in the tank until thoroughly mixed before applying. Continue agitation and apply immediately.

Almonds: Brown Rot, Shot-hole Fungus, Leaf Blight - Use 2 pounds per 100 gallons of water in pre-bloom, blossom and petal fall periods. Scab - Use 2 pounds per 100 gallons of water at petal fall and 5 weeks after petal fall. Do not use later than 5 weeks after petal fall.

Apples: Scab - Use 1 to 1 1/2 pounds per 100 gallons of water in pre-blossom, blossom and early cover sprays. Bulls Eye Rot (Oregon & Washington) - Use 1 1/2 to 2 pounds per 100 gallons of water beginning 12 to 14 days after calyx stage and repeat as necessary in summer sprays. Perennial Canker (Oregon) - Use 1 to 1 1/2 pounds per 100 gallons of water in pre-blossom and calyx sprays.

Apricots: Brown Rot, Shot-hole Fungus - Use 2 pounds per 100 gallons of water in pre-bloom, blossom and petal fall thru early cover sprays.

Cherries: Brown Rot, Blossom Blight - Use 1 1/2 pounds per 100 gallons of water in popcorn, full blossom and petal fall stages. Leaf Spot - Use 1 1/2 pounds per 100 gallons of water in petal fall and shuck fall stages and approximately 2 weeks after shuck fall. Do not apply within 7 days of harvest.

Ziram Spray

Fungicide

EPA Reg. No. 279-268 EPA Est. 274 NY-1

Active Ingredient		
Ziram (zinc dimethyl dithiocarbamate)	26.0%	
Inert Ingredients	24.0%	
		100.0%
Zinc expressed as metallic	16.25%	

KEEP OUT OF REACH OF CHILDREN

CAUTION

May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, throat and skin. Avoid contact with eyes, skin or clothing. Avoid breathing dust or spray mist. In case of contact, flush with plenty of water. For eyes, get medical attention.

For Emergency Assistance Call 716-735-3765.

See Side Panel For Additional Precautionary Statements.

FMC

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia, Pennsylvania 19103

PA 9.80

Net Weight

Nectarines, Peaches: Peach Blight (Coryneum) and Leaf Curl - Use 2 to 3 pounds per 100 gallons of water in the fall before heavy rains. For Leaf Curl alone use 2 to 3 pounds in 100 gallons of water as a dormant application prior to bud swell. Brown Rot and Peach Blight (Coryneum) - Use 2 pounds in 100 gallons of water in pre-bloom, blossom and petal fall sprays. May be used in early cover and before harvest. Remove excess residue by washing or wiping.

Pears: Scab - Use 1 to 1 1/2 pounds per 100 gallons of water in pre-blossom, blossom and early cover sprays. Storage Rot (Oregon) - Use 1 to 1 1/2 pounds per 100 gallons of water in pre-harvest spray.

Cucumbers, Melons: Scab, Anthracnose - Use 2 pounds per 100 gallons of water per acre at 7 day intervals when plants are small. Increase rate to 3 pounds per 100 gallons of water per acre after plants vine.

Beans (Snap, Lima): Anthracnose - Use 2 pounds per acre in sufficient water for thorough coverage when plants are small. Increase to 3 pounds per acre on large plants. Make applications at 7 day intervals starting when true leaves appear. Do not apply to snap beans later than 4 days before picking.

Tomatoes: Anthracnose, Early Blight - Use 2 to 3 pounds per 100 gallons of water per acre at 7 to 10 day intervals starting 2 to 4 weeks after plants are set. Under severe disease conditions apply 3 pounds in 180 to 200 gallons per acre at 7 day intervals.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

