



Code 135

Pounds Net Weight

Polysulphide Compound

(in dry form)

Fungicide-Insecticide

EPA Reg. No. 279-385

EPA Est., 279-

Active Ingredients:	By Wt.
Sodium Polysulphide	84.0%
Sodium Thiosulphate	8.0%
Inert Ingredients:	8.0%
	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

For Emergency Assistance Call 716-735-3765.

See Other Panels for Additional Precautionary Information.

ACCEPTED

JAN 9 1981

Federal Insecticide, Fungicide, and Rodenticide Act

279-385

FMC

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

PA 11/80

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

Avoid contact with skin. Keep away from domestic animals.

Environmental Hazards

Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Store in a cool, dry place. Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or residue that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State or local disposal authorities. To dispose of container, triple rinse (or equivalent) and dispose of in an approved landfill, or bury in a safe place away from water supplies. Consult Federal, State or local disposal authorities.

Do not use this material in combination with Standard or Basic Lead Arsenate. Application of materials containing sulphur should not be made just prior to or during periods of excessively high temperature.

Polysulphide Compound is a dry product which is soluble in water.

Fill the spray tank about two-thirds full of water. Then add Polysulphide Compound, sifting it in gradually. Keep the agitator in motion while the remainder of the tank is being filled and the powder will dissolve very quickly in water.

Apply at the rate of 12 to 14 pounds per 100 gallons of water in delayed dormant stage. Use on Apples and Pears to prevent Apple or Pear Scab infections, and to aid in the control of hatching eggs and young nymphs of Green Apple Aphid, Rosy Aphid, and Pear Psylla.

At the above rate, use on Apples, Pears, Plums and Prunes to control San Jose Scale.

Apply at the rate of 5 to 6 lbs. of Polysulphide Compound plus 3 to 3½ gallons of dormant oil emulsion per 100 gallons of water. Use on Apple, Pear, Plum, Prunes, and Cherries for the control of San Jose Scale, European Red Mite, and Brown Almond Mite in the dormant or delayed dormant.