



Code 135

Pounds Net Weight

# Polysulphide Compound

(in dry form)

## Fungicide-Insecticide

EPA Reg No 279-385      EPA Est 279-

Active ingredients	By Wt
Sodium Polysulphide .....	84.0%
Sodium Thiosulphate .....	14.0%
Inert ingredients .....	2.0%
	100.0%

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

For Emergency Assistance Call 716-735-3765.

See Other Panels for Additional Precautionary Information.

# FMC

FMC Corporation  
Agricultural Chemical Group  
2000 Market Street  
Philadelphia Pennsylvania 19103

PA 8 61

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans (& Domestic Animals)

Avoid contact with skin. Keep away from domestic animals.

#### Environmental Hazards

Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### Storage and Disposal

Store in a cool, dry place. Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of according to procedures approved by Federal, State or local disposal authorities. To dispose of container, triple rinse (or equivalent) and dispose of in an approved landfill or bury in a safe place away from water supplies. Consult Federal, State or local disposal authorities.

Do not use this material in combination with Standard or Basic Lead Arsenate. Application of materials containing sulphur should not be made just prior to or during periods of excessively high temperatures.

Polysulphide Compound is a dry product which is soluble in water.

Fill the spray tank about two-thirds full of water. Then add Polysulphide Compound, sifting it in gradually. Keep the agitator in motion while the remainder of the tank is being filled and the powder will dissolve very quickly in water.

Apply at the rate of 12 to 14 pounds per 100 gallons of water in delayed dormant stage. Use on Apples and Pears to prevent Apple or Pear Scab infections, and to aid in the control of hatching eggs and young nymphs of Green Apple Aphid, Rosy Aphid, and Pear Psylla.

At the above rate, use on Apples, Pears, Plums and Prunes to control San Jose Scale.

Apply at the rate of 5 to 6 lbs. of Polysulphide Compound plus 3 to 3.5 gallons of dormant oil emulsion per 100 gallons of water. Use on Apple, Pear, Plum, Prunes, and Cherries for the control of San Jose Scale, European Red Mite, and Brown Almond Mite in the dormant or delayed dormant.

On Peaches in the Colder Areas of the Northwest where the trees go completely dormant, treat in the dormant period for the control of the above pests: Leaf Curl and Peach Blight.

Apply at the rate of 6 lbs. per 100 gallons of water in combination with 2 gallons summer flowable emulsion or 6 lbs. of Wettable Sulphur. Use on Pear to control Pear Blister Mite at Post harvest.

Apply at the rate of 1.5 to 2 lbs. Polysulphide Compound per 100 gallons of water. Use on Peaches in pre-blossom sprays for the control of Brown Rot and Peach Blight. On Peaches to control Brown Rot and Powdery Mildew use 1 to 3 pounds of Polysulphide Compound and 5 pounds of Kainspray or Wettable Sulphur per 100 gallons of water. Apply when conditions are optimum for disease development. Keep new growth covered throughout these periods.

Apply at the rate of 2 to 3 lbs. Polysulphide Compound plus 4 to 5 lbs. of Wettable Sulphur per 100 gallons of water. On Apples and Bartlett Pears in pre-blossom sprays to control Apple and Pear Scab and Powdery Mildew.

**Dealers Should Sell in Original Packages Only.**

**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions:

**Warranty:** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and of handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to the amount not exceeding the purchase price paid and shall not include incidental or consequential damages.

135-8 81-PA

