

Code: 30042

WETTABLE SULPHUR

DIRECTIONS FOR USE

MIXING DIRECTIONS: Fill the spray tank with water until the agitator blades are covered. With the agitator running sift in the required amount of Wettable Sulphur. Continue filling and keep agitator running.

ALMONDS: Brown Rot, Scab — Use 6 pounds per 100 gallons of water in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

APPLES AND PEARS: For Scab or Powdery Mildew control, use as a spray in Pre-blossom, Calyx and Early cover sprays. Use 6 to 8 pounds of Wettable Sulphur or 4 to 6 pounds of Wettable Sulphur plus 2 to 3 pounds Polysulphide Compound or 2 gallons of Liquid Lime Sulphur Solution per 100 gallons of dilute spray. Sulphur may injure certain varieties of APPLES. Check specific varietal tolerance to sulphur prior to using. **Pears only:** Do not apply after fruit begins to turn down. **Fall Clean-Up Spray:** Blister Mites, European Red Mites and Twospotted Spider Mites — Use 6 pounds of Wettable Sulphur plus 6 pounds of Polysulphide Compound or 5 gallons of Liquid Lime Sulphur Solution to each 100 gallons of dilute spray.

BLACKBERRIES, BOYSENBERRIES, GOOSEBERRIES, LOGANBERRIES, RASPBERRIES, STRAWBERRIES: Powdery Mildew — Use 2 pounds per 100 gallons of water. Apply when disease first appears and repeat as necessary.

CITRUS: Citrus Flat Mite, Yuma Spider Mite, Sixspotted Mite and Citrus Rust Mite — Use 5 to 12 pounds per 100 gallons of water (75 to 125 pounds per acre). Apply when mites appear and repeat as necessary.

GRAPES: Powdery Mildew — Use 2 to 5 pounds per 100 gallons of dilute spray. Use as a spray beginning when shoots are 6 to 8 inches long and repeat at 7 to 10 day intervals. Ground application preferred. Concord and other Labrusa type grapes may be injured by sulphur.

FUNGICIDE-INSECTICIDE

EPA REGISTRATION NO. 279-387 AA

| | |
|--------------------------|----------------|
| ACTIVE INGREDIENT: | BY WT. |
| Sulphur | 92.00% |
| INERT INGREDIENTS: | 8.00% |
| | <u>100.00%</u> |

This material in this bag is guaranteed to pass 95% through a 325 mesh screen.

PLUMS, PRUNES, PEACHES, NECTARINES: Cherry Leaf Spot, Prune Rust — Use 2 to 4 pounds per 100 gallons of water. (200 gallons of water per acre). Apply during post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

TOMATOES: Tomato Russet Mite — Apply with ground equipment at 2 to 4 gallons of water per acre. Use only applicator type agitation. Continuous and controlled spray.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Applications of materials containing sulphur should not be made prior to or during periods of excessively high summer temperatures or within 2 weeks of an oil spray. On citrus do not make an oil application within 60 days prior to or after a sulphur application.

Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, or other undesirable results may occur.

**For Emergency Assistance Call
716-735-3765**

ENVIRONMENTAL AND

Do not apply when weather conditions are unfavorable. Do not apply where runoff may occur. Do not apply to water supplies. Empty bags should be disposed of by burying or burning. Keep Out Of Smoke.

ACCEPTED

DEC 19 1974

UNDER THE FEDERAL INSECTICIDE AND FUNGICIDE ACT AND RELATED ACTS FOR ECONOMIC PEST CONTROL REGISTERED UNDER NO. 279-387 AA SUBJECT TO ATTACHED COMMENTS.

WETTABLE SULPHUR

Net Weight
50 Pounds

CONDITIONS FOR USE

Spray tank with water until the agitator running sift in the sulphur. Continue filling and keep

Use 6 pounds per 100 gallons of water during pre-bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

For Powdery Mildew control, use as a pre-bloom spray. Use 6 to 8 pounds of Wettable Sulphur plus 1 pound of Liquid Lime per 100 gallons of dilute spray. Sulphur may injure plants with low tolerance to sulphur. Check specific varietal tolerance to sulphur. Do not apply after fruit begins to set. Blister Mites, European Red Mites, Spider Mite, Sixspotted Mite and others. Use 6 pounds of Wettable Sulphur plus 1 pound of Liquid Lime per 100 gallons of dilute spray.

GRAPES, GOOSEBERRIES, LOGANBERRIES, PEACHES, PRUNES: Powdery Mildew — Use 2 pounds per 100 gallons of dilute spray. Apply when disease first appears and repeat at 7 to 10 day intervals.

Use 2 to 5 pounds per 100 gallons of water beginning when shoots are 6 to 8 inches long. Repeat application at 7 to 10 day intervals. Ground application on Labrusca type grapes may be injured.

FUNGICIDE-INSECTICIDE

EPA REGISTRATION NO. 279-387 AA

| | |
|--------------------------|----------------|
| ACTIVE INGREDIENT: | BY WT. |
| Sulphur | 92.00% |
| INERT INGREDIENTS: | 8.00% |
| | <u>100.00%</u> |

This material in this bag is guaranteed to pass 95% through a 325 mesh screen.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Applications of materials containing sulphur should not be made prior to or during periods of excessively high summer temperatures or within 2 weeks of an oil spray. On citrus do not make an oil application within 60 days prior to or after a sulphur application.

Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, or other undesirable results may occur.

For Emergency Assistance Call
716-735-3765



PLUMS, PRUNES, PEACHES, NECTARINES, CHERRIES: Powdery Mildew, Cherry Leaf Spot, Prune Rust and Brown Rot — Use 5 to 10 pounds per 100 gallons of water (By ground apply a minimum of 200 gallons of water per acre). Apply in the pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

TOMATOES: Tomato Russet Mite — Use 12 to 25 pounds per acre. Apply with ground equipment at the rate of 60 to 250 gallons of water per acre. Use only application equipment which has paddle type agitation. Continuous and complete agitation is necessary.


ENVIRONMENTAL AND DISPOSAL CAUTIONS

Do not apply when weather conditions favor drift from the areas treated. Do not apply where run-off is likely to occur.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty bags should be buried with wastes or burned. Keep Out Of Smoke.



ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions.

WARRANTY: FMC warrants that this product, as of the time of sale by FMC (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured in U.S.A.
FMC CORPORATION
AGRICULTURAL CHEMICALS DIVISION
Middleport, N.Y. 14105

FMC
Chemicals