



ACCEPTED  
COMMENTS  
EPA Letter Dated

Code 30042

JAN 5 1987

Pounds Net Weight

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, this pesticide is registered under EPA Reg. No. 279-387

PHYSICAL OR CHEMICAL HAZARDS

Keep away from heat, sparks, or flame. Do not smoke while applying this product.

# Wettable Sulfur

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

## Fungicide-Insecticide

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

EPA Reg. No 279-387AA

EPA Est., 279-

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

ACTIVE INGREDIENT:

BY WT.

Sulfur.....	92.0%
Inert Ingredients:.....	8.0%
	100.0%

The material in this bag is guaranteed to pass 95% through a 325 mesh screen.

Do not enter treated areas for 24 hours unless appropriate protective clothing is worn.

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

KEEP OUT OF REACH OF CHILDREN

### CAUTION

CAUTION

Area treated with Wettable Sulfur on (insert date). Do not enter treated areas for 24 hours unless appropriate protective clothing is worn (insert State Department of Agriculture's re-entry interval, if more restrictive).

### PRECAUTIONARY STATEMENTS Hazards to Humans (& Domestic Animals)

Caution: Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

In case of accidental exposure to pesticide spray or dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.

### Statement of Practical Treatment

If swallowed: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If in eyes: Flush with plenty of water. Get medical attention if irritation persists.

If on skin: Wash with plenty of soap and water. Get medical attention.

For Emergency Assistance Call 716-735-3765

See other panels for additional precautionary information.

### STORAGE AND DISPOSAL

Pesticide Storage

Keep away from fire and sparks

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold back roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Contain spills. Call FMC Collect: (716) 735-3765.

To contain spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.



FMC Corporation  
Agricultural Chemical Group  
2000 Market Street  
Philadelphia Pennsylvania 19103

7/30/86-R

Page 1 of 3

Fill the spray tank with water until the agitator blades are covered with the agitator running. Sift in the required amount of Wettable Sulfur. Continue filling and keep agitator running.

**Almonds:** Brown Rot Scab—Use 6 pounds per 100 gallons of water in pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

**Apples and Pears:** For Scab or Powdery Mildew control, use as a spray in pre-blossom, Calyx and Early cover sprays. Use 6 to 8 pounds of Wettable Sulfur or 4 to 6 pounds of Wettable Sulfur plus 2 to 3 pounds Polysulphide Compound or 2 gallons of Liquid Lime Sulfur Solution per 100 gallons of dilute spray. Sulfur may injure certain varieties of Apples. Check specific variety tolerance to sulfur prior to using. **Pears only:** Do not apply after fruit begins to turn down. **Fall Clean-Up Spray:** Bister Mites, European Red Mites and Twospotted Spider Mites—Use 6 pounds of Wettable Sulfur plus 6 pounds of Polysulphide Compound or 5 gallons of Liquid Lime Sulfur Solution to each 100 gallons of dilute spray.

**Citrus:** Citrus Flat Mite, Yuma Spider Mite, Sixspotted Mite and Citrus Rust Mite—Use 5 to 12 pounds per 100 gallons of water (75 to 125 pounds per acre). Apply when mites appear and repeat as necessary. Do not make an oil application within 60 days prior to or after a sulfur application.

**Grapes:** Powdery Mildew—Use 2 to 5 pounds per 100 gallons of dilute spray. Use as a spray beginning when shoots are 6 to 8 inches long and repeat at 7 to 10 day intervals. Ground application preferred. Concord and other Labrusca type grapes may be injured by sulfur.

**Plums, Prunes, Peaches, Nectarines, Cherries:** Powdery Mildew, Silver Mite, Cherry Leaf Spot, Prune Rust and Brown Rot—Use 5 to 10 pounds per 100 gallons of water. (By ground apply a minimum of 200 gallons of water per acre.) Apply in the pre-bloom, bloom and post-bloom periods. Apply when disease threatens during rain periods and repeat at 7 to 10 day intervals.

**Tomatoes:** Tomato Russet Mite—Use 12 to 25 pounds per acre. Apply with ground equipment at the rate of 60 to 250 gallons of water per acre. Use only application equipment which has paddle type agitation. Continuous and complete agitation is necessary.

### Dealers Should Sell in Original Packages Only.


**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions:

**Warranty:** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.



Wettable Sulfur Fungicide-Insecticide  
Code 30042


Applications of material containing sulfur should not be made prior to or during periods of excessively high summer temperatures or within 2 weeks of an oil spray.

To mix Wettable Sulfur, fill the spray tank with water until agitator blades are covered. With agitator running, sift in the required amount of Wettable Sulfur. Continue filling and keep agitator running during spraying.

Blackberries, Boysenberries, Gooseberries, Loganberries, Raspberries, Strawberries: To control Powdery Mildew -- Use 2 pounds Wettable Sulfur per 100 gallons of water. Apply when disease first appears and repeat at 7-10 day intervals or as necessary to maintain control.

*Pistachios: Citrus Flat Mite - Use 50 pounds per acre. Apply when mites first appear and repeat as necessary. May be applied by ground or air.*

10rb  
012



279-581

Page 3 of 3