

CODE  
93

NET WEIGHT  
25 LBS.

ACCEPTED  
July 31, 1973

# CARBAMATE

## FUNGICIDE

### INSTRUCTIONS

To mix this material, slowly sift the desired amount into the partially filled tank of water with the agitator running and when thoroughly mixed apply at once with continuous agitation. Where desired this material may first be made to a thick paste by adding a little water. This should then be diluted and added to the tank while it is filling.

### DIRECTIONS FOR USE

Where this material is used with Lead Arsenate, no Lime need be used as a safener, since this material is itself a good safener for Lead Arsenate. Observe all cautions and limitations on labeling of all products used in mixtures.

OBSERVE DAYS INTERVAL BETWEEN LAST APPLICATION AND HARVEST INDICATED BY NUMBER IN ( ) FOLLOWING CROP.

**APPLES (7):** For early-season control of Scab, Cedar Rust, Blotch - Use 1 to 1½ pounds of this material, or 1 pound plus 3 pounds Kolospray or Kolofog, per 100 gallons of water in pre-bloom, calyx and early cover sprays. For rust control make an additional bloom application using ½ pound per 100 gallons of water.

For late-season control of Scab, Black Rot, Bitter Rot, Sooty Blotch, Fly Speck, and Brooks' Spot - Use ½ pound per 100 gallons of water. Where bitter rot and Brooks' spot are severe problems increase dosage to 1½ to 2 pounds per 100 gallons of water.

On Golden Delicious, Jonathan, Baldwin and other varieties of apples which are especially apt to russet some injury may result from the use of this type of material if used in the early season or bloom applications. The best time for use is in the cover sprays.

When used in combination with sulfur, do not apply within 10 days to 2 weeks of an oil spray or just prior to or during periods of excessively high temperatures.

**PEARS (7):** Scab, Sooty Blotch - Use 1½ pounds per 100 gallons of water in pink, calyx, first and second cover sprays, and 1 pound in summer sprays.

**PEACHES (21):** Leaf Curl - Use 1½ pounds per 100 gallons of water. Apply in the fall after leaves drop or in the spring before buds begin to swell.

**CHERRIES (9):** Brown Rot, Leaf Spot - Use 1½ pounds per 100 gallons of water in petal-fall and cover sprays.

**GRAPES (7):** Black Rot - Use 2 pounds per 100 gallons of water, applying just before bloom, just after bloom and two weeks later.

### CONTAINS FERBAM

EPA REG. NO. 279-388-AA

ACTIVE INGREDIENT:  
Ferbam (Ferric dimethyl dithiocarbamate) 76.0%  
INERT INGREDIENTS: 24.0%  
100.0%

## CAUTION

### KEEP OUT OF REACH OF CHILDREN.

For Emergency Assistance Call  
716-735-3765

May cause irritation of eyes, nose, throat and skin! Avoid breathing dust or spray mist. Avoid contact with eyes, skin or clothing. In case of contact, flush with plenty of water; for eyes, get medical attention.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Keep out of tidal marshes and estuaries. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Rinse equipment and DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Do not reuse container. Destroy when empty.

Keep packages or dust away from sparks, flames or hot surfaces.

**CITRUS (0)** including Grapefruit, Lemons, Limes, Oranges, Tangerines, Tangelos: Scab - Use 1½ pounds per 100 gallons of water. Do not apply more than 1000 gallons of spray per acre. Apply during pre-bloom period and when 2/3 of petals have fallen. May be applied in late summer or early fall if a heavy flush of growth appears.

**CRANBERRIES: Fruit Rots** - Use 2 pounds per 100 gallons of water. Begin applications early in blossoming period and repeat at two week intervals for a total of 5 applications. Do not apply within 28 days after mid-bloom.

**FAIRY RING:** Use 9 pounds per 100 gallons of water and apply 1 gallon of this mixture to a one foot square area. Treat an area 3 feet beyond the advancing line of dead vines and 2 feet within this line. Apply in the fall immediately after harvest.

**CABBAGE (Plant Beds Only): Downy Mildew** - Use 1½ pounds per 100 gallons of water, making 2 or 3 applications each week.

**TOBACCO (Plant Beds): Blue Mold** - Use 2 to 3 pounds per 100 gallons of water. Apply at the rate of 3 gallons per 100 square yards when plants are small, increasing to 6 gallons when plants are ready for transplanting. Begin applications when plants are the size of a dime, or when blue mold is reported in the area, and repeat twice weekly until plants are transplanted.

**TOMATOES: Gray Mold (Botrytis)** - Use 3 pounds per 100 gallons of water. Spray for complete coverage. Make seed bed application just before seedlings break the soil surface and repeat at 7 day intervals. Make field application when disease threatens and repeat at 7 to 10 day intervals as necessary to maintain control.

**ORNAMENTALS: Black Spot and Rust on roses; Rust on snapdragons and carnations; Leaf Spot on Chrysanthemums; Botrytis Blight on peonies and tulips** - Use 1 to 1½ pounds per 100 gallons of water, applying at 7 to 10 day intervals during growing season.

**PINES: Fusarium Rust of Seedlings:** Use 1 to 2 pounds per 100 gallons of water. Apply at 3 to 7 day intervals as needed, beginning when plants emerge. Southern Cane Rust in slash and long leaf pines: Use 2 pounds per 100 gallons of water plus 4 to 8 ounces of suitable spreader-sticker. Begin at time of flowering when strobili are emerging from bud scales. Repeat at 5 day intervals until conelet stage. Direct spray to flower-bearing portions of tree crowns.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



**Dealers Should Sell in Original Packages Only.**  
**USAGE CAUTION:** DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROP OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS, BECAUSE PLANT INJURY, EXCESSIVE RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

**TERMS OF SALE OR USE:** On purchase or use of this product buyer and user agree to the following conditions:

**WARRANTY:** Niagara warrants that this product, as of the time of sale by Niagara, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

**DAMAGES:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to Niagara within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

**EXCUSE FROM NON-CONTROLLABLE CONDITIONS:** Because of certain critical conditions created or incurred by buyer or user or over which Niagara has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release Niagara from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

MANUFACTURED IN U. S. A. BY



NIAGARA CHEMICAL DIVISION  
FMC CORPORATION  
MIDDLEPORT, NEW YORK



FRESNO, CALIF. - RICHMOND, CALIF.  
RIVERSIDE, CALIF. - YAKIMA, WASHINGTON