

CODE  
162

# TOXAKIL MISCIBLE

NET CONTENTS  
5 Gallons

## INSECTICIDE

E.P.A. Reg. No. 279-405 E.P.A. Est. No. 279-FL-1

### ACTIVE INGREDIENTS:

*Toxaphene -----	58.0%
Petroleum Solvent -----	36.0%
INERT INGREDIENTS: -----	6.0%
	100.0%

\* Technical Chlorinated Camphene, 67 to 69% Chlorine  
CONTAINS 6 POUNDS TOXAPHENE PER GALLON.

## WARNING KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed! Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Do not store near feed and food products.

For Emergency Assistance Call 716-735-3765

### ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Rinse equipment and containers and DISPOSE OF WASTES by burying at least 18 inches deep in non-crop lands away from water supplies. Do not reuse container; smash or perforate and bury with wastes.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

DO NOT STORE AT TEMPERATURES BELOW 0° F.

**FMC**  
FMC Corporation  
Agricultural Chemical Division  
Middletown New York 14135  
036-S

ACC  
APR  
Under the  
EPA Reg. No. 279-405

### DIRECTIONS FOR USE (CROPS)

Observe use limitations. Apply listed PINTS PER ACRE, unless otherwise designated, in sufficient water for thorough coverage. When applying this material by aircraft apply the recommended rate in a minimum of 1 to 1 1/2 gallons of water per acre. For use with ground equipment apply specified rate in a minimum of 10 to 15 gallons of water per acre. Do not use in undiluted form.

ALFALFA (Grown for seed): Alfalfa Weevil, Armyworm, Webworm, Pea Aphid - Use 2-2/3 to 4 pints. Grasshopper, Lygus Bug, Climbing Cutworms - Use 2 to 2-2/3 pints. Do not feed treated forage to dairy animals or animals being finished for slaughter.

BEANS (Green, Dry, Lima): Armyworm, Cutworm - Use 2-2/3 pints. Grasshopper - Use 2-2/3 to 4 pints. Bean Beetle - Use 2-2/3 to 5-1/3 pints. Do not apply to green or snap beans after pods begin to form. Do not feed treated vines to dairy animals or animals being finished for slaughter. No limitations on use of shelled beans as human food.

CABBAGE: Corn Earworm, Diamondback Moth - Use 2 to 5-1/3 pints. Fall Armyworm, Green Peach Aphid - Use 2 to 2-2/3 pints. Do not apply after heads start to form.

CARROTS: Grasshoppers - Use 2-2/3 to 4 pints per acre. May be applied up to day of harvest.

CORN: Armyworm, Fall Armyworm - Use 2 to 4 pints per acre. If larvae are young. Climbing Cutworms, Budworm, Chinch Bug - Use 2-2/3 to 4 pints. European Corn Borer - Use 2-1/3 to 3 pints per acre. Hopper - Use 2 to 2-2/3 pints. Do not feed treated forage to dairy animals or animals being finished for slaughter. No limitations on use of grain.

BARLEY, OAT, RYE, WHEAT: Armyworm, Cutworm, Grasshopper, Chinch Bug - Use 2-2/3 pints. Do not feed treated forage to dairy animals or animals being finished for slaughter.

SOYBEANS: Armyworm, Bean Leaf Beetle, Grasshopper - Use 2 to 2-2/3 pints. Flea Beetle - Use 2 pints. Do not apply within 21 days of harvest. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals, or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

COTTON: For Texas, Oklahoma and New Mexico early-season control of thrips and fleahoppers, use 1 1/2 to 3 pints of this material per acre. Applications should be repeated as required, beginning at the four-leaf stage or earlier.

For the control of cutworms and armyworms use 3 to 4 pints of this material per acre. Apply when needed. Do not apply after bolls open. For late-season control of fleahoppers and lygus and other plant bugs, use 3 to 4 pints of this material per acre. Repeat applications as required to maintain effective control. Do not apply after bolls open. Do not graze or feed gin waste to dairy animals or animals being finished for slaughter.

ORNAMENTAL TURF: Armyworm - Use 2-2/3 to 4 pints. Chigger - Use 1-1/3 to 2-2/3 pints. Grasshoppers - Use 2 to 2-2/3 pints. Do not graze treated areas.

ORNAMENTALS: Boxelder Bug - Use 4 pints. Cankerworm, Walnut Caterpillar - Use 2-2/3 to 4 pints. Fall Armyworm - Use 2-2/3 pints. Fall Webworm - Use 2 to 4 pints. Saw Fly, Tent Caterpillar - Use 2 pints.

PEARS: Pear Psylla - Use 1 pint per 100 gallons of water. Do not apply within 40 days of harvest or after second cover spray.

PRE-PLANT SOIL APPLICATION (Beans, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Celery, Collards, Corn, Cucumbers, Eggplant, Lettuce, Peas, Peppers, Tomatoes): Japanese Beetle Grub, May Beetle Grub - Use 3-1/3 gallons per acre in sufficient water to cover. Use as a single pre-planting soil treatment or before edible parts begin to form. Do not apply to land to be planted in crops with edible roots.

### DIRECTIONS FOR USE (LIVESTOCK)

Use only as a spray or in a backrubber device. Do not use as a dip. Use only on animals more than 6 months of age. Do not use on dairy animals. Do not spray overheated animals or when weather is stormy or chilly. Do not spray animals or allow animals access to backrubbers within 28 days of slaughter.

BEEF CATTLE (SPRAY APPLICATION): Horn Flies - Add 1 gallon to 150 gallons of water. Apply to backs of animals every 3 weeks or as needed. Do not apply more than 2 quarts of spray per animal. Lice, Ticks - Add 1 gallon to 150 gallons of water. Apply enough spray to wet animal to the skin. For control of spinose ear tick, make certain to treat the ear and ear zone. Repeat as required.

BEEF CATTLE (BACKRUBBER APPLICATION): Horn Flies - Dilute 1 gallon with 15 gallons of diesel oil. One gallon of this solution will treat 20 linear feet of cable. Re-treat the burlap every 3 to 4 weeks.

SHEEP: Lice, Sheep Ticks (Keds) - Add 1/2 gallon to 150 gallons of water. Apply enough spray to wet animals to the skin. Re-treat after 2 to 3 weeks if required.

SWINE: Lice - Add 1 gallon to 150 gallons of water. Apply enough spray to wet the skin of the animals. Re-treat after 2 to 3 weeks if required. Do not treat sows within 2 weeks of farrowing nor at least 3 weeks thereafter.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROP OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUE OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

### DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Printed in U.S.A.