

Code: 30061

KOLODUST[®] C-O-C-S[®] 50-6 DUST

50 Pounds Net Weight

Maybe applied with airplane or ground application equipment.

DIRECTIONS FOR USE

ON ALMONDS to control Brown Rot and Shot Hole Fungus apply 40-50 pounds per acre. Apply in pre-blossom and petal fall periods.

ON PEACHES and NECTARINES to control Peach Blight, Powdery Mildew, Brown Rot and Peach Leaf Curl use 40-50 pounds per acre. Apply as a fall treatment and in spring at bud swell. To control Brown Rot and Peach Blight use 40-50 pounds per acre. Apply in pre-blossom and petal fall periods.

ON PRUNES AND PLUMS to control Brown Rot use 40-50 pounds per acre. Apply in the popcorn and full bloom period.

ON CHERRIES to control Brown Rot and Leaf Spot use 40-50 pounds per acre. Apply in popcorn, bloom and petal fall periods.

ON APPLES and PEARS in pre-blossom and blossom periods for the control of Fire Blight and Scab use 15-30 pounds per acre. All blossoms must be protected ahead of periods of high humidity. Number of applications to be determined by length of the bloom period. For scab control after the blossom period use a suitable fungicide which does not contain copper. Do not use on Anjou or Conice varieties of pears.

ON CITRUS (California) for prevention of Brown Rot use 50 pounds per acre. Apply during the winter months. Treat the ground around the tree and the skirts of the trees to a height of 3 to 4 feet. Give particular attention to covering the fruit. Treat inside as well as outside, covering the trunks and skirts thoroughly. Retreatment may be necessary. Do not disturb soil after treatment. In areas with a history of severe copper damage spray the skirts no more than 2 feet high.

ON FILBERTS to control Bacterial Blight use 50 pounds per acre. Two applications are necessary. Apply first application in August, the second when half of the leaves have dropped. If the fall is unusually wet, apply a third application when most of the leaves have dropped. Apply during when trees are wet with dew.

ON GRAPES to control Powdery Mildew and Black Rot use at the rate of 20-30 pounds per acre. Apply at the time berries begin to set and thereafter as needed. These applications will also assist in the control of Powdery Mildew. For disease control the material must be applied before the disease appears. Apply with ground application equipment only.

FUNGICIDE
U.S.D.A. REGISTRATION NO. 279-453

Active Ingredients	By Weight
Copper sulphate as acetate	6.00%
Sulphur	42.00%
Inert Ingredients	52.00%
	100.00%

The copper present is in the form of Basic Sulphate and Chlorides.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if Swallowed.
Avoid inhaling dust.
Avoid excessive skin contact.
Keep away from domestic animals, feed and feedstuffs.

USAGE CAUTION

Application of materials containing sulphur should not be applied prior to or during periods of excessively high temperatures.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from treated areas.

Dispose of wastes by burying in non-crop land away from water supplies. Empty bags should be treated with wastes or buried. Keep out of smoke.

FOR DISEASE CONTROL ON THE FOLLOWING CROPS USE 25-35 POUNDS PER ACRE AT 7 TO 10 DAY INTERVALS TO PROTECT NEW GROWTH THROUGHOUT THE GROWING SEASON.

CABBAGES: For control of Leaf Blight.

EGGPLANT: For control of Late Blight.

PEPPERS: For control of Late Blight.

LETTUCE: For control of Downy Mildew.

POTATOES: For control of Early and Late Blight.

SUGAR BEETS: For control of Leaf Spot.

TOMATOES: For control of late blight, early blight and Septoria Leaf Spot.

CELERY: For control of Celery Blight.

HOPS: For control of Downy Mildew. As a crown treatment for early downy mildew control apply in the early spring.


ACCEPTED

Jan 15, 1968

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
ED UNDER NO. 279-453 SUBJECT
TO ATTACHED COMMENTS

279-453

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.
USAGE CAUTION: DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROP OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS, BECAUSE PLANT INJURY, EXHAUSTIVE RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE. On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY. Niagara warrants that this product, as sold, (1) is of the quality of sale by Niagara, (2) conforms to the ingredient statement on the label, and (3) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES. Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to Niagara within 90 days of purchase by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS. Because of certain critical conditions created or incurred by buyer or user or over which Niagara has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release Niagara from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

MANUFACTURED IN U.S.A. BY



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, NEW YORK



FRESNO, CALIF. — RICHMOND, CALIF.
RIVERSIDE, CALIF. — YAKIMA, WASHINGTON