

Code: 30060

Net Weight  
50 Pounds

# PHOSKIL® 2 DUST

## DIRECTIONS FOR USE

Apply when pests first appear and repeat as required. May be applied by ground or air. Observe days interval between last application and harvest indicated by number in ( ) following crop.

**ARTICHOKES (7):** Artichoke Plume Moth — Use 25 pounds per acre.

**BEANS (7):** Aphids, Whitefly, Leafhopper and Mexican Bean Beetle — Use 25 pounds per acre.

**BROCCOLI (7), BRUSSELS SPROUTS (7), CABBAGE (7), CAULIFLOWER (7), KOHLRABI (7):** Aphids, Vegetable Weevil, Armyworms, Diamondback Moth Larvae, and Imported Cabbageworm — Use 15 to 20 pounds per acre. Climbing Cutworms, Cabbage Loopers — Use 25 pounds per acre.

**CARROTS (15):** Aphids, Vegetable Weevil and Petrobia Mite — Use 25 pounds per acre. Do not use treated carrot tops for food or feed.

**CELERY:** Aphids, Celery Leaf-tier, Climbing Cutworms — Use 20 to 25 pounds per acre. Leafhoppers — Use 20 pounds per acre. Do not apply within 21 days of harvest. Leafminers, Loopers, Whiteflies — Use 35 pounds per acre. Do not apply within 30 days of harvest.

**COLLARDS (10), KALE (10), MUSTARD GREENS (10):** Aphids, Vegetable Weevil, Cabbage Looper, Armyworms, Climbing Cutworms, Diamondback Moth Larvae, Imported Cabbageworm — Use 25 pounds per acre.

**CORN (12):** Spider Mites — Use 25 pounds per acre. Do not apply within 12 days of cutting for forage.

**CUCUMBERS (15) and SQUASH (15):** Aphid, Climbing Cutworms, Cucumber Beetle, Squash Bug, Leafminer, Thrips and Flea Beetle — Use 25 pounds per acre. Do not apply before vining on cucumbers.

**EGGPLANT (15):** Aphid, Whitefly, and Leafminer — Use 25 pounds per acre.

**GRAPES (14):** Thrips, Hoplia Beetle, Grape Leaf-roller — Use 25 to 33 pounds per acre. Grape Mealybug — Use 30 to 35 pounds per acre. Ground application preferred.

**HOPS (15):** Aphids — Use 25 to 40 pounds per acre.

**LETTUCE (Head 7, Leaf 21):** Aphids, Leafhoppers, Leafminer, Armyworms, Cabbage Looper, Imported Cabbageworm — Use 25 pounds per acre.

**MELONS (7):** Aphids, Leafhopper, Leafminer, Thrips, Petrobia Mite and Cucumber Beetles — Use 25 pounds per acre.

**ONIONS (15):** Leafminer, Thrips and Petrobia Mite (California) — Use 25 pounds per acre.

**PEACHES, NECTARINES:** Oriental Fruit Moth, European Red Mite, Two-Spotted Spider Mites and Brown Mite — Use 50 pounds per acre. Apply by ground or air. Apply when insects are in damaging numbers. In California; do not apply within 21 days of harvest. Do not apply more than once after bloom. Do not apply more than 5 pounds actual parathion per acre between January 1 and harvest. In areas other than California; do not apply within 14 days of harvest. Do not apply more than 5 pounds actual parathion per acre per year.

## INSECTICIDE

EPA REGISTRATION NO. 279-464 ZA

ACTIVE INGREDIENTS:

Parathion; 0,0-Diethyl 0-p-nitrophenyl thiophosphate ..... 2.00%

INERT INGREDIENTS: ..... 98.00%

BY WT.

2.00%

98.00%

100.00%

**PEAS (10):** Aphid, Pea Weevil, Thrips, Climbing Cutworms, Armyworms and Leafminer — Use 25 pounds per acre.

**PEPPERS (15):** Aphids — Use 10 to 15 pounds per acre.

**PUMPKINS (10):** Aphids, Cucumber Beetle — Use 20 pounds per acre. Climbing Cutworms and Squash Bugs — Use 25 pounds per acre.

**SMALL GRAINS (Barley, Oats, Wheat) (15):** Petrobia (Brown Wheat) Mite, Black Grass Bug and Stinkbugs — Use 25 to 37 pounds per acre. Aphids — Use 25 pounds per acre.

**SPINACH (14):** Aphids, Leafminer, Cabbage Looper, Armyworm — Use 25 pounds per acre. California: Crown Mites and Lepidoptera Caterpillars — Use 25 pounds per acre.

**STRAWBERRIES (14):** Aphids, Lygus Bugs, Leafhoppers, Strawberry Leafroller and Whitefly — Use 25 pounds per acre.

**TOMATOES (10):** Aphid, Leafminer — Use 25 pounds per acre. Russet Mites and Stink Bugs — Use 12 to 22 pounds per acre.



### DANGER - POISON



### KEEP OUT OF REACH OF CHILDREN

**DANGER:** Poisonous if swallowed, inhaled or absorbed through skin! Rapidly absorbed through skin! Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Bureau of Mines for Parathion protection. Keep all unprotected persons and beneficial animals out of operating areas or vicinity where there may danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

#### FIRST AID

Call physician immediately. If swallowed give tablespoon of table salt in a glass of warm water. Repeat until vomiting occurs and vomitus is clear. Do not induce vomiting in an unconscious person. If in eyes, flush eyes with lots of water for 15 minutes. If on skin or clothing remove contaminated clothing or shoes immediately. Bathe with soap and water.

#### ANTIDOTE

NOTE TO PHYSICIANS: Active ingredient is parathion. Patient should be under observation for 48 hours. Parathion is a cholinesterase inhibitor. Atropine is antidotal. Give 4 mg. intramuscularly or intravenously and repeat every 15 minutes until patient is fully atropinized. If cyanotic give artificial respiration. Pralidoxime chloride (2-PAM) should be used in treatment.


#### ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where run-off is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSE OF WASTES by burying at least 18 inches deep in non-crop land away from water supplies. Empty bags should be buried with wastes. Do not burn.

For Emergency Assistance Call  
716-735-3765



**ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.**



**Dealers Should Sell in Original Packages Only.**


**DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.**

**TERMS OF SALE OR USE:** On purchase or use of this product buyer and user agree to the following conditions:

**WARRANTY:** FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.**

**DAMAGES:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

**EXCUSE FROM NON-CONTROLLABLE CONDITIONS:** Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from, liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.



Manufactured in U.S.A. by:  
FMC CORPORATION  
AGRICULTURAL CHEMICAL DIVISION  
Middleport, N.Y. 14105

**FMC**  
Chemicals