

Code: 237



Gallons Net Contents

CHLORDANE 8.0 MISCIBLE

DIRECTIONS FOR USE

Use the required amount of Chlordane 8.0 Miscible in sufficient water to give thorough coverage. Do not use in undiluted form. Unless otherwise noted, for Corn Rootworm, Root Maggots, Wireworms and White Grub control, apply as a PRE-PLANT broadcast treatment to the soil and immediately work it into the top few inches prior to planting. For band and furrow applications reduce dosages in proportion to the area treated. When applying this material by aircraft to listed vegetable and field crops, mix the recommended amount with sufficient water to provide a minimum of 2 gallons of finished spray per acre. Care should be taken that this material is not allowed to drift onto neighboring crop or non-crop areas.

BROCCOLI, CABBAGE, CAULIFLOWER: Wireworms, White Grubs — Use 3 pints per acre. Root Maggot (seed bed treatment) — Use 8 pints per acre.

CORN: Wireworms, White Grubs — Use 3 pints per acre. Corn Rootworm — Use 1 pint per acre as a furrow treatment at time of planting.

BEETS, PEPPERS: Wireworms — Use 3 pints per acre.

POTATOES: Wireworms — Use 5 to 6 pints per acre.

RADISH: Root Maggot — Apply 2 pints per acre to the soil surface in 200 to 400 gallons of water. Do not apply after edible parts start to form.

TURNIPS: Wireworm, Root Maggot — Use 2 pints per acre. For root maggot apply to the soil surface in 200 to 400 gallons of water. Do not apply after edible parts begin to form.

LAWNS, TURF: Wireworms — 3 to 6 pints per acre. For 2 to 3 years control use 14 pints per acre (1/3 pint per 1000 sq. ft.) in sufficient water. Wash insecticide 4 to 6 inches into soil with water. For non-seeded areas, cultivate insecticide into soil to a depth of 4 to 6 inches immediately after applying. Do not allow children or animals on treated areas until the insecticide has been washed into the soil by drenching and is completely dry.

NOTE: Birds feeding over treated areas may be harmed.

ORNAMENTALS (non-crop areas): Grasshoppers — 1/4 pints per acre. Apply when insects appear and repeat as required.

GRASSHOPPER CONTROL: On uncultivated land, field crops not used for food or feed, wheat, oats, rye, barley and corn, use 1 to 2 pints per acre. Do not apply to grain crops after heads begin to form. Do not graze animals in treated fields. Do not feed treated forage to dairy or meat animals.

SHADE TREES, ORNAMENTALS, PERENNIALS, ANNUALS: Armyworms, Cutworms, Fall Armyworms, Japanese Beetle (adults), June Beetle (adults), May Beetles — Apply 1 pint per 100 gallons of water. Apply when insects appear and repeat as necessary.

ANTS, FALSE CHINCH BUGS, EARWIGS and SOW BUGS along fences, trash piles, at the base of trees and in other hiding places, apply this material at the rate of 1 to 2 quarts per 100 gallons of water. In spraying in orchards and vineyards, direct the spray to the soil around the tree trunks. DO NOT APPLY TO THE FRUIT. Do not graze livestock in treated orchards or vineyards. Apply when insects appear and repeat as necessary.

GRAPES: For the control of false chinch bugs in Grape Vineyards use 2-4 pints in sufficient water to thoroughly cover one acre. (By ground apply a minimum of 200 gallons of finished spray per acre). Apply when insect first appears and repeat as

INSECTICIDE

EPA REGISTRATION NO. 279-538 AA

ACTIVE INGREDIENTS:	BY WT.
*Technical Chlordane	72.00%
Petroleum Distillate	23.00%
INERT INGREDIENTS:	5.00%
	100.00%

*Equivalent to 43.2% Octochlor 4,7 methano tetrahydroindane and 28.8% of related compounds. Contains 8 lbs. of Technical Chlordane per gallon.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Contact with skin causes toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.

USAGE CAUTIONS

Do not apply this material on livestock or household pets.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes and containers. Apply this product only as specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Rinse equipment and containers and DISPOSE OF WASTES by burying in non-crop land away from water supplies. Punch holes in containers and bury in non-crop land away from water supplies.

WARNING — FLAMMABLE

Keep away from heat and open flame. Keep containers closed. Leaking packages should be removed to a safe place. Do not store below 10° F.

NIAGARA CHEMICAL DIVISION
FMC CORPORATION



MIDDLEBURY, N.Y.
FRESNO, CALIFORNIA
RICHMOND, CALIF. WAKIMA, WASHINGTON



required. Do not apply after edible parts start to form. When pests are a problem after edible parts start to form apply to ground and weed growth around the vine yards outside the spread of border vines. Do not contaminate the edible fruit.

TERMITE CONTROL IN BUILDINGS: Mix 1 pint Niagara Chlordane 8 Miscible per 1 gallon of water for the following use:

SLAB-TYPE CONSTRUCTION:

(a) — Apply 1 gallon of finished spray per 10 sq. ft. as an overall pre-construction treatment under the slab and under attached porches.

(b) — Apply 2 gallons of finished spray per 5 lineal feet to critical areas under the slab as pre-construction treatment, such as along the inside of foundation walls around utility entrances and interior partition foundation walls.

(c) — Apply 2 gallons of finished spray per 5 lineal feet along the outside of the foundation using a trench to the depth of the foundation.


CRAWL-SPACE HOUSES AND HOMES WITH BASEMENT:

(a) — Apply 2 gallons of finished spray per 5 lineal feet as a trenching application in critical areas under the house, such as along the inside of foundation walls and around piers and utility entrances. If trench is deeper than 15-18 inches, apply 4 gallons per 5 lineal feet.

(b) — Apply 2 gallons of finished spray per 5 lineal feet in a trench along the outside of foundation walls, including entrance platforms and porches where the foundation is not more than 15-18 inches deep. If over 15-18 inches deep but not over 30 inches deep, apply 4 gallons per 5 lineal feet of trench.

(c) — Apply 1 gallon of finished spray per 10 square feet of soil surface as a pre-construction treatment where the attached porches, entrance platforms, utility entrance etc., have covering slab on fill of ground.

For Emergency Assistance Call
716-735-3765



Do not apply when weather conditions favor drift from the areas treated. Do not apply when runoff is likely to occur. Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

Dealer Should Sell in Original Packages Only.

WARRANTY: Niagara warrants that this product, as of the time of sale by Niagara, conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

042W

Printed in U.S.A.

*Amendment to be made
additional use
CHLORDANE 8.0 MISCIBLE*

*DIRECTIONS: To control wireworms
apply 4 to 5 pints per acre.*



Gallons Net Contents

CHLORDANE 8.0 MISCIBLE

DIRECTIONS FOR USE

Amount of Chlordane 8.0 Miscible in sufficient water to give thorough use in undiluted form. Unless otherwise noted, for Corn Rootworm, Wireworms and White Grub control, apply as a PRE-PLANT broadcast soil and immediately work it into the top few inches prior to planting. Row applications reduce dosages in proportion to the area treated. When applied by aircraft to listed vegetable and field crops, mix the recommended amount of sufficient water to provide a minimum of 2 gallons of finished spray per acre. Do not be taken that this material is not allowed to drift onto neighboring areas.

CABBAGE, CAULIFLOWER: Wireworms, White Grubs — Use 3 pints per acre (seed bed treatment) — Use 8 pints per acre.

CORN: White Grubs — Use 3 pints per acre. Corn Rootworm — Use 1 pint per acre. Furrow treatment at time of planting.

WIREWORMS: — Use 3 pints per acre.

WHITE GRUBS: — Use 5 to 6 pints per acre.

ROOT MAGGOT: — Apply 2 pints per acre to the soil surface in 200 to 400 gallons of water. Do not apply after edible parts start to form.

CORN ROOTWORM: — Use 2 pints per acre. For root maggot apply 2 pints per acre in 200 to 400 gallons of water. Do not apply after edible parts start to form.

WIREWORMS: — 3 to 6 pints per acre. For 2 to 3 years control use 14 pints per acre (at per 1000 sq. ft.) in sufficient water. Wash insecticide 4 to 6 inches deep. For non-seeded areas, cultivate insecticide into soil to a depth of 4 to 6 inches immediately after applying. Do not allow children or animals on treated areas until insecticide has been washed into the soil by drenching and is completely dry.

PLANTS: Growing over treated areas may be harmed.

NON-CROP AREAS: Grasshoppers — 1/4 pints per acre. Apply when needed repeat as required.

CONTROL: On uncultivated land, field crops not used for food or feed, such as alfalfa, barley and corn, use 1 to 2 pints per acre. Do not apply to grain crops until they are in the seedling stage. Do not graze animals in treated fields. Do not feed treated areas to meat animals.

ORNAMENTALS, PERENNIALS, ANNUALS: Armyworms, Cutworms, Fall Armyworm, Japanese Beetle (adults), June Beetle (adults), May Beetles — Apply 1 pint per acre. Apply when insects appear and repeat as necessary.

WITCH GRASS, EARWIGS and SOW BUGS along fences, trash piles, at the base of trees and in other hiding places, apply this material at the rate of 1 to 2 quarts per acre. In spraying in orchards and vineyards, direct the spray to the trunk of the tree. DO NOT APPLY TO THE FRUIT. Do not graze livestock in orchards or vineyards. Apply when insects appear and repeat as necessary.

CONTROL OF FALSE CHINCH BUGS in Grape Vineyards use 2-4 pints in 200 to 400 gallons of water to thoroughly cover one acre. (By ground apply a minimum of 200 gallons of water per acre). Apply when insect first appears and repeat as necessary.

INSECTICIDE

EPA REGISTRATION NO. 279-538 AA

ACTIVE INGREDIENTS:	BY WT.
*Technical Chlordane	72.00%
Petroleum Distillate	23.00%
INERT INGREDIENTS:	5.00%
	100.00%

*Equivalent to 43.2% Octochlor 4,7 methano tetrahydroindane and 28.8% of related compounds. Contains 8 lbs. of Technical Chlordane per gallon.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Contact with skin causes toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.

USAGE CAUTIONS

Do not apply this material on livestock or household pets.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes and containers. Apply this product only as specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Rinse equipment and containers and DISPOSE OF WASTES by burying in non-crop land away from water supplies. Punch holes in containers and bury in non-crop land away from water supplies.

WARNING — FLAMMABLE

Keep away from heat and open flame. Keep containers closed. Leaking packages should be removed to a safe place. Do not store below 10° F.

NIAGARA CHEMICAL DIVISION
FMC CORPORATION



MIDDLETOWN, N.Y.
PRESHOP, CALIFORNIA

ST. LOUIS, MO. • WASHINGTON, D.C.

required. Do not apply after edible parts start to form. When pests are a problem after edible parts start to form apply to ground and weed growth around the vineyards outside the spread of border vines. Do not contaminate the edible fruit.

TERMITE CONTROL IN BUILDINGS: Mix 1 pint Niagara Chlordane 8 Miscible per 12 gallons of water for the following use:

SLAB-TYPE CONSTRUCTION:

(a) — Apply 1 gallon of finished spray per 10 sq. ft. as an overall pre-construction treatment under the slab and under attached porches.

(b) — Apply 2 gallons of finished spray per 5 lineal feet to critical areas under the slab as pre-construction treatment, such as along the inside of foundation walls and around utility entrances and interior partition foundation walls.

(c) — Apply 2 gallons of finished spray per 5 lineal feet along the outside of the foundation using a trench to the depth of the foundation.


CRAWL-SPACE HOUSES AND HOMES WITH BASEMENT:

(a) — Apply 2 gallons of finished spray per 5 lineal feet as a trenching application to critical areas under the house, such as along the inside of foundation walls and around piers and utility entrances. If trench is deeper than 15-18 inches, apply 4 gallons per 5 lineal feet.

(b) — Apply 2 gallons of finished spray per 5 lineal feet in a trench along the outside of foundation walls, including entrance platforms and porches where the foundation is not more than 15-18 inches deep. If over 15-18 inches deep but not over 30 inches deep, apply 4 gallons per 5 lineal feet of trench.

(c) — Apply 1 gallon of finished spray per 10 square feet of soil surface as a pre-construction treatment where the attached porches, entrance platforms, utility entrances, etc., have covering slab on fill of ground.

For Emergency Assistance Call
716-735-3765



Do not apply when weather conditions favor drift from the areas treated. Do not apply when runoff is likely to occur. Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

Dealer Should Sell in Original Packages Only.

WARRANTY: Niagara warrants that this product, as of the time of sale by Niagara, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

Printed in U.S.A.

*Amendment to be made
additional use*

CHLORDANE 8.0 MISCIBLE

WATERMELONS: To control wireworms apply 4 to 5 pints per acre.