


Code 20

Kolodust Phos Kil 1.5 Dust

50 Lbs. Net Weight



Fungicide - Insecticide

EPA Reg. No. 279-712 EPA Est. 279-NJ-1

Active Ingredients:	
Sulfur	77.0%
1 Parathion	1.5%
Inert Ingredients:	21.5%
100.0%	

Active Ingredients: Sulfur, Ortho-phosphorothioic acid, triphenyl phosphorothioate

Inert Ingredients: Silica, talc, iron pyrites, iron sulfide

WARNING

KEEP OUT OF REACH OF CHILDREN

Keep out of reach of children. Do not swallow. Do not get in eyes, on face or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator if necessary as required by the U.S. Bureau of Mines for Parathion and its fumes. Keep all non-treated persons and beneficial animals out of the treating area. Do not enter areas where there may be a danger of drift. Ventilate areas where you have been treated. Do not get on your hands and face until you have washed. Do not contaminate food or feed stuff. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash up and change your clothes with soap and hot water before sleeping.

FIRST AID

If Swallowed: Give tablespoon of table salt in a glass of warm water. Repeat until vomiting occurs and vomitus is clear. Never induce vomiting in an unconscious person. Call a physician immediately.

If In Eye: Flush eye for 15 minutes with plenty of water. Call a physician immediately.

If On Skin or Clothing: Remove contaminated clothing and shoes immediately. Bathe with soap and water. Call a physician immediately.

ANTIDOTE

Note to Physician: Active ingredient is a Cholinesterase inhibitor. Patient should be kept under observation for 48 hours. Atropine is antidote. Give 4 mg. intramuscularly or intravenously. Repeat every 15 minutes until patient is fully atropinized. If necessary give artificial respiration. Pralidoxime chloride (2-PAM) should be used in treatment. For additional assistance call 1-800-368-6868.

Environmental and Usage Cautions

This product is toxic to fish, birds and mammals. Do not feed or treat animals that have been exposed to lakes, ponds and streams. Do not apply around fruit trees, to cut, Do not apply when weather conditions favor drift from areas treated. Do not use machinery or equipment or discharge dust or waste. Apply this product only as specified on the label.

This product is highly toxic to bees exposed to direct treatment or to a dust on a leaf. For further information may be obtained from your supplier or Agricultural Extension Service.

In case of waste, by burying it at least 18 inches deep in non-crop lands away from water supplies. Do not reuse containers. Bury with wastes.

Directions for Use

NOT FOR USE OR STORAGE IN OR AROUND THE HOME

Do not prune trees or thin fruit in orchards until a reasonable time (at least two weeks) has been allowed for dissipation of vapors from residues. Applications of this material should not be made while crops are in bloom. Applications of Dusting and Spraying materials containing sulfur should not be made just prior to or during periods of excessively high summer temperature.


Apple Curculio, Codling Moth, Mites, Scab - Use at the rate of 1 1/2 to 2 pounds per tree depending upon the size. Apply at petal-fall and repeat at intervals of 5 to 7 days or as often as required to maintain effective control. Do not apply within 14 days of harvest. Do not apply just prior to or during periods of excessively high summer temperatures or within 10 to 14 days of an oil spray. Applications of this product may cause fruit and foliage injury to Golden Delicious, Jonathan, McIntosh and related varieties.

Beans Mexican Bean Beetle, Leafhoppers, Aphids, Red Spiders - Use at the rate of 30 pounds of dust per acre. Apply when insects or mites first appear and repeat as required to maintain effective control. Dust coverage of both upper and lower leaf surfaces is essential for good control. Do not apply within 7 days of harvest.

Peach (areas other than California) Curculio, Leafhopper, Pear, Oriental Fruit Moth, Mites, Scab, Brown Spot - Use at the rate of 12 1/2 to 15 pounds of dust per tree depending on the size. Make first application at the "chuck-sprout" stage and repeat at 5 to 7 day intervals or as often as required to maintain effective control. Do not apply within 14 days of harvest. If it is necessary to treat for brown rot or peach scab during this 14 day period Kolodust may be applied. Do not apply more than 5 pounds of active ingredient per acre per year.

Strawberries Red Spiders, Aphids - Use at the rate of 15 pounds of dust per acre. Apply when insects or mites first appear and repeat at 5 to 10 day intervals as needed to maintain effective control. Dust plants thoroughly. Do not apply within 14 days of harvest.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions.

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.**

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.