

Code 30279

Net Contents  
Gallons

# Triona\* Light Medium Emulsion

## Insecticide-Miticide

EPA Reg. No 279-847

EPA Est., 279-

Active Ingredient:	By Wt.
Petroleum Oil .....	80.00%
Inert Ingredients: .....	20.00%
	100.00%

Classification—Light Medium  
Unsulphonated Residue—92% Minimum  
\*T.M. Shell Chemical Corp

**KEEP OUT OF REACH OF CHILDREN**

### CAUTION

#### PRECAUTIONARY STATEMENTS

##### Hazards to Humans (& Domestic Animals)

###### Caution

Harmful if swallowed. Do not induce vomiting. Call a Physician immediately.

For Emergency Assistance Call 718-735-3765

See Other Panels for Additional Precautionary Information

FMC Corporation  
Agricultural Chemical Group  
2000 Market Street  
Philadelphia Pennsylvania 19103



#### Environmental Hazards

Keep out of lakes, ponds, or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur. Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### Storage and Disposal

Do not store below 10° F (-12° C). Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. To dispose of container, triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill, or bury in a safe place. Do not cut or weld this container. Empty 28 gallon drums should be returned to your pesticide supplier. Do not reuse this container. Consult Federal, State or local disposal authorities for approved alternative procedures.

#### Usage Cautions

Do not make an oil spray application within 2 weeks prior to or after an application of sulfur. On Citrus do not make an oil application within 60 days prior to or after an application of sulfur.

Avoid application of oil sprays just prior to or during hot weather or during rainy weather or dry hot winds. Apply during dormancy when temperature is above 40° F. Avoid double spraying. Plants receiving oil sprays should be in healthy condition and proper moisture should be present. Use efficient equipment.

#### Mixing Directions

Before using this product be sure the spray tank is clean and contains just enough water to cover the agitator shaft. With the agitator in motion (120 R.P.M. or more) add the amount of spray oil required for a full tank. Pump this mixture through the overflow and back into the tank for a minute or two. The mixture should turn white, indicating good emulsification. Then fill the tank with water. Keep agitator running at all times.

If an injector is used first add 25 to 50 gallons of water to tank with engine running to provide good pressure. Then put injector suction tube into the measured amount of spray oil, opening injector valve to allow the oil to be sucked into the tank. Then fill the tank with water.

Do not use if material does not emulsify  
not use in undiluted form.

#### Directions for Use

Apply when pests first appear and repeat as necessary. Do not undiluted form. Apply the following recommended rates in surface water to thoroughly cover one acre:

**Citrus (Oranges, Lemons, Grapefruit):** Use 1½ to 2 gallons material per 100 gallons of water for control of Yellow Scale, Citrus Scale, Red Scale and Soft Brown Scale.

**Olive:** Use 1½ to 2½ gallons of this material per 100 gallons of water in dormant, delayed dormant or summer periods for the control of Olive Scale, Black Scale and Oleander Scale.

**Deciduous Trees (Apples, Pears, Peaches, Plums, Prunes, Cit Almonds):** Use 1 to 1½ gallons per 100 gallons of water in dormant or summer periods for the control of Spider Mites. (Some varieties of plums and prunes may be injured by oil sprays. Test tolerance to oil prior to treatment.)

**Pears:** After harvest for the control of Blister Mite, Bud Mite and Spider Mites use at the rate of 1½ to 2 gallons in combination with 6 pounds of FMC Kolofog per 100 gallons of dilute spray.

#### Dealers Should Sell in Original Packages (Terms of Sale or Use)

On purchase of this product buyer agrees to the following conditions:

**Warranty:** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as warranted, the product is sold as is. Buyer and user assume all liability for use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully and method of application, weather and crop conditions. Mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer and are assumed by buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such use or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for which no warranty or negligence shall be limited to direct damages exceeding the purchase price paid and shall not include incidental or consequential damages.

30279

Net Contents  
Gallons

# Imona\* Light Medium Emulsion

## Insecticide-Miticide

No. 279-847 EPA Est., 279-

Ingredient:	By Wt.
Mineral Oil	80.00%
Active Ingredients:	20.00%
	100.00%

Emulsion—Light Medium  
Residual Residue—92% Minimum  
FMC Chemical Corp.

KEEP OUT OF REACH OF CHILDREN

### CAUTION

#### PRECAUTIONARY STATEMENTS

Hazardous to Humans (& Domestic Animals)

If swallowed. Do not induce vomiting. Call a Physician immediately.

Emergency Assistance Call 716-735-3766

See Back Panel for Additional Precautionary Information



FMC Corporation  
Agricultural Chemical Group  
1000 Market Street  
Philadelphia Pennsylvania 19103

#### Environmental Hazards

Keep out of lakes, ponds, or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur. Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### Storage and Disposal

Do not store below 10° F (-12° C). Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. To dispose of container, triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill, or bury in a safe place. Do not cut or weld this container. Empty 28 gallon drums should be returned to your pesticide supplier. Do not reuse this container. Consult Federal, State or local disposal authorities for approved alternative procedures.

#### Usage Cautions

Do not make an oil spray application within 2 weeks prior to or after an application of sulfur. On Citrus do not make an oil application within 60 days prior to or after an application of sulfur.

Avoid application of oil sprays just prior to or during hot weather or during rainy weather or dry, hot winds. Apply during dormancy when temperature is above 40° F. Avoid double spraying. Plants receiving oil sprays should be in healthy condition and proper moisture should be present. Use efficient equipment.

#### Mixing Directions

Before using this product, be sure the spray tank is clean and contains just enough water to cover the agitator shaft. With the agitator in motion (120 R P M or more), add the amount of spray oil required for a full tank. Pump this mixture through the overflow and back into the tank for a minute or two. The mixture should turn white, indicating good emulsification. Then fill the tank with water. Keep agitator running at all times.

If an injector is used, first add 25 to 50 gallons of water to tank with engine running to provide good pressure. Then put injector suction tube into the measured amount of spray oil, opening injector valve to allow the oil to be sucked into the tank. Then fill the tank with water.

Do not use if material does not emulsify. Do not use in undiluted form.

#### Directions for Use

Apply when pests first appear and repeat as necessary. Do not use in undiluted form. Apply the following recommended rates in sufficient water to thoroughly cover one acre:

**Citrus (Oranges, Lemons, Grapefruit):** Use 1½ to 2 gallons of this material per 100 gallons of water for control of Yellow Scale, Citricola Scale, Red Scale and Soft Brown Scale.

**Olive:** Use 1½ to 2½ gallons of this material per 100 gallons of water in dormant, delayed dormant or summer periods for the control of Olive Scale, Black Scale and Oleander Scale.

**Deciduous Trees (Apples, Pears, Peaches, Plums, Prunes, Cherries, Almonds):** Use 1 to 1½ gallons per 100 gallons of water in delayed dormant or summer periods for the control of Spider Mites. Certain varieties of plums and prunes may be injured by oil sprays. Check tolerance to oil prior to treatment.

**Pears:** After harvest for the control of Blister Mite, Bud Mite and Red Spider Mites use at the rate of 1½ to 2 gallons in combination with 4 to 6 pounds of FMC Kolotog per 100 gallons of dilute spray.

#### Dealers Should Sell in Original Packages Only.

**Terms of Sale or Use:** On purchase of this product buyer and user agree to the following conditions:

**Warranty:** FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

**Directions and Recommendations:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

**Use of Product:** FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

**Damages:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.