

ACCEPTED
June 5, 1970
 UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 279-909 SUBJECT TO ATTACHED COMMENTS 30337



Gallons
 Net Contents

PHOSKIL® 4.0 MISCIBLE

INSECTICIDE
 U.S.D.A. REGISTRATION NO. 279-909

ACTIVE INGREDIENTS:	BY WT.
†Parathion	47.00%
Petroleum Distillate	46.00%
INERT INGREDIENTS:	7.00%
	100.00%

†0,0-Diethyl 0-p-nitrophenyl thiophosphate
 Contains 4.0 Pounds Parathion Per Gallon

⚠ DANGER - POISON ⚠
KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL WARNING STATEMENTS

FIRST AID TREATMENT
 ANTIDOTE: If swallowed, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. CALL A PHYSICIAN IMMEDIATELY! If on skin, in case of contact, remove contaminated clothing and immediately wash skin with soap and water. Parathion is a cholinesterase inhibitor and treatment of poisoning should include the use of atropine.

ATROPINE IS ANTIDOTAL
 Keep on hand an emergency supply of 20 tablets, grains 1/100. In case of poisoning take 3 tablets, grains (1/100) and CALL A PHYSICIAN AT ONCE!

PHYSICIAN: Enough atropine must be given in the shortest practical time to produce dilation of pupils. Patient must be under constant supervision and additional atropine may be necessary without delay. Clear chest by postural drainage. Use artificial respiration and oxygen if patient becomes comatose. Use of airway may be necessary.

DIRECTIONS FOR USE:
 Apply when pests first appear and repeat as necessary. USE IN UNDILUTED FORM. Use the following directions in sufficient water to thoroughly cover one acre. (By ground apply a minimum of 200 gallons per acre on stone fruits and nuts; a minimum of 300 gallons per acre on pome fruits; a minimum of 200 gallons per acre on bush and vine fruits and a minimum of 10 gallons per acre on vegetable and row crops. By air, apply a minimum of 2 gallons per acre on vegetable and field crops and a minimum of 20 gallons per acre on small fruits and tree crops). Observe days interval between last application and harvest indicated by number in () following the crop.

APPLES (14), PEARS (14): Codling Moth, Orange Tortrix, San Jose, Parlatoria and Erios Scales, Woolly Apple Aphid, Mealy Bug, Bud Moth, Fruit Tree Leafroller, Leafhoppers, Rosy Apple Aphid, Green Apple Aphid — Use 1/2 - 2/3 pint per 100 gallons. By airplane, 3 pints per acre. **ON PEARS:** Blister Mite, Pear Psylla — Use 1/2 - 2/3 pint per 100 gallons in pre-blossom and blossom sprays. By airplane, 2-3 pints per acre. May cause injury to fruit and foliage of McIntosh and related varieties, to Golden Delicious and to Bosc Pears.

CHERRY (14): Cherry Fruitworm, Oriental Fruit Moth, Bud Moth, Orange Tortrix, Cankerworm, Fruit Fly, San Jose Scale Crawlers, Aphids — Use 1/2 pint per 100 gallons. By airplane use 2 pints per acre.

FIG (30): Fig Scale — Use 1/2 pint per 100 gallons.

PLUMS (14) PRUNES (14): Peach Twig Borer, Orange Tortrix, Leafhopper, Codling Moth, Tussock Moth, Bud Moth, Mealy Plum Aphid, Thrips, Fruit Tree Leafroller, San Jose Scale — Use 1/2-2/3 pint per 100 gallons. By airplane use 2-3 pints per acre.

NECTARINES PEACHES: Oriental Fruit Moth, Peach Twig Borer, Aphids, Fruit Tree Leafroller, Cat Facing Insects — Use 1 pint per 100 gallons (apply a minimum of 2 quarts per acre). Thrips — Use 1/2-1 pint per 100 gallons (apply a minimum of 2 pints per acre). **CALIFORNIA:** Do not apply more than 5 pounds of actual Parathion per acre between January 1 and harvest. Do not apply more than once after bloom. Do not apply within 21 days of harvest. **AREAS OTHER THAN CALIFORNIA:** Do not apply more than 5 pounds actual Parathion per acre per year. Do not apply within 14 days of harvest.

GRAPES (14): Thrips, Hoplia Beetle, Mealy Bug, Leafroller — Use 1/2-2/3 pint per 100 gallons (200 gallons per acre). Grape Mealy Bug: Dormant application use 2 1/2 pints plus 2 gallons Dormant Oil Emulsion per 100 gallons. Apply 200 gallons per acre as a dormant spray. Apply only before fruit is size of buckshot or after harvest. Ground application preferred.

STRAWBERRIES (14): Aphids, Lygus Bugs, Leafhoppers, Strawberry Leafroller, Root Weevil, Whitefly — Use 1/2-2/3 pint per 100 gallons. By airplane use 1 pint per acre.

BEANS (15): Corn Earworm, Diabrotica Beetles, Leafhoppers, Looper, Aphids, Leafminer, Whitefly, Armyworm — Use 1-1 1/2 pints per acre.

LETTUCE (Head 7) (Leaf 21): Aphids, Leafhoppers, Leafminer, Loopers, Armyworms — Use 1 pint per acre.

SPINACH (14): Aphids, Leafminer, Seed Corn Maggot in Crowns — Use 1 pint per acre.

WARNING: Poisonous If Swallowed, Inhaled, or Absorbed Through Skin! Rapidly absorbed through skin. Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U.S. Department of Agriculture for parathion protection. Keep all unprotected persons and beneficial animals out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

USAGE CAUTIONS

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Birds feeding on treated areas may be killed. Apply this product only as specified on the label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Rinse equipment and containers and **DISPOSE OF WASTES** by burying at least 18 inches deep in non-crop lands away from water supplies. Empty containers should be smashed and buried with wastes. Do not reuse.

Do not allow this material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

Dealer Should Sell in Original Packages Only.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: Niagara warrants that this product, as of the time of sale by Niagara, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to Niagara within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which Niagara has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release Niagara from, liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

NIAGARA CHEMICAL DIVISION

FMC CORPORATION

