

[279-1061]

CODE
1237

MALATHION 5 DUST

NET WEIGHT
50 LBS.

INSECTICIDE

DIRECTIONS FOR USE

Make Thorough, full coverage applications beginning when insects first appear and repeat as necessary to maintain control. May be applied with airplane or ground application equipment. Observe use limitations. Observe days interval between last application and harvest as indicated by number in parenthesis () following crop.

BEANS (1): Aphids, Cucumber Beetles, Leafhoppers, Mexican Bean Beetle - Use 30 to 35 lbs. per acre.

BROCCOLI (3), BRUSSELS SPROUTS (7), CABBAGE (7), CAULIFLOWER (7), COLLARDS (7), KALE (7), MUSTARD (7), TURNIP (3): Aphids, Imported Cabbageworm, Cabbage Looper, Flea Beetles - Use 40 to 50 lbs. per acre.

CUCUMBERS (1), MELONS (1), PUMPKINS (3), SQUASH (1): Aphids, Cucumber Beetle, Leafhoppers, Spider Mites, Squash Vine Borer - Use 35 lbs. per acre. For Vine Borer control apply weekly to stems and vines at base of plants. Do not apply unless plants are dry.

LETTUCE (Head-7), (Leaf - 14): Aphids, Cabbage Looper, Leafhoppers, Mites - Use 30 to 40 lbs. per acre.

ONIONS (3): Thrips, Onion Maggots - Use 30 to 40 lbs. per acre.

PEAS (3): Pea Aphid, Pea Weevil - Use 25 lbs. per acre. Do not apply within 7 days of harvest if vines are to be used for feed.

PEPPERS (3): Pepper Maggots - Use 30 lbs. per acre.

POTATOES (0): Aphids, False Chinch Bug, Leafhoppers, Mealybug - Use 25 to 30 lbs. per acre.

SPINACH (7): Aphids - Use 25 to 30 lbs. per acre.

TOMATOES (1): Aphids, Drosophila, Spider Mites, Russet Mite - Use 30 to 40 lbs. per acre.

APPLES (3): Aphids, Codling Moth, Curculio, Leaf Rollers, Mealybug, Mites - Use 1/2 to 2 lbs. per tree. May injure foliage on some varieties.

PEACHES (7): Curculio, Oriental Fruit Moth, Mites - Use 1/2 to 1 lb. per tree.

E. P. A. Reg. No. 279-1061

ACTIVE INGREDIENTS:

*Malathion 5.0%

INERT INGREDIENTS: 95.0%

100.0%

*O, O-Dimethyl dithiophosphate of diethyl mercaptosuccinate.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs.

NOTE TO PHYSICIAN: Malathion, may cause cholinesterase inhibition. Atropine is antiaotal.


For Emergency Assistance Call 716-735-3765

USAGE CAUTIONS

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.



COTTON: Aphids; Boll Weevil, Cotton Leafworm, Cotton Leaf Perforator, Fleahoppers, Fall Armyworms, Garden Webworm, Grasshoppers, Leafhoppers, Lygus Bugs, Mites, Thrips, Whitefly - Use 25 to 50 lbs. per acre.

PASTURE AND RANGE GRASS, GRASS AND GRASS HAY: Aphids, Armyworms, Leafhopper - Use 30 to 40 lbs. per acre.

PECANS: Pecan Nut Casebearer - Use 80 lbs. per acre.

Apply when first brood larvae begin to enter nuts and repeat after 7 days if necessary. Pecan Bud Moth - Apply 8 lbs. per acre by airplane. Apply 4 to 5 times, 2 to 3 weeks apart when insects appear.

ORNAMENTALS: Aphid. Make thorough full coverage application. Injury may occur on certain ferns and on some species of crassula. Do not use on canaerti juniper.

BEEF AND DAIRY CATTLE: Hornfly - Using a handful (1.5 oz.) of dust per animal, whisk the dust over the animals back and neck. Repeat at 14 day intervals as required to maintain control. Do not apply to cows within 5 hours before milking or during milking. Avoid contamination of water trough, feedstuff and milking equipment.

BEEF AND NON-MILKING CATTLE: Lice - Dust animals thoroughly. Repeat in 10 days and thereafter as needed. Do not treat calves under one month of age. Do not apply to lactating dairy animals, or non-lactating dairy animals within 2 weeks of freshening.

HOGS: Lice - Dust animal thoroughly. Repeat in 10 days and thereafter as needed.

POULTRY HOUSES: Chicken Red Mite, Northern Fowl Mite, Poultry Lice - Use 1 lb. per 50 to 60 square feet. Apply dust to litter, floor space and to nests, roosts and adjacent areas.

ACCEPTED