CODE 20350

MANGASULFUR 10 TOXAKIL 5 DUST

DIRECTIONS FOR USE

BEANS (Including Dry, Green and Lima): Rust, Powdery Mildew, Armyworms, Corn Earworm, Climbing Cutworm, Thrips and to correct deficiencies of manganese.

Use as a dust at the rate of 40 to 50 pounds per acre depending upon the size of the plants. Begin applications when diseases or insects first appear and repeat as necessary for control (10 day intervals).

For best results make applications with a dusting machine that will give uniform coverage of all above-ground surfaces. Do not apply to green or snap beans within 7 days of harvest. Do not feed treated vines to dairy animals or animals being finished for slaughter. For disease control within this period the use of a suitable fungicide alone, such as Niagara Kolozim Bean Dust is recommended.

Applications of spraying or dusting materials containing sulfur should not be made just prior to or during periods of excessively high temperatures.

FUNGICIDE-INSECTICIDE-NUTRITIONAL


U. S. D. A. Reg. No. 279–1095 ACTIVE INGREDIENTS:	
Sulfur73.70%	
*Toxaphene 5.00%	USAGE CAUTIONS
INERT INGREDIENTS: ** 21.30%	
100.00% Sulfur fineness: 93% through 325 mesh *Technical Chlorinated Camphene 67–69% chlorine	This product is toxic to fish and wildlife. Keep out of any body of water. Birds feeding in treated areas may be killed. Apply this pro- duct only as specified on this label.
**Manganese expressed as metallic2.04%	This product is toxic to bees and should not be applied when bees are actively visiting the area.
CAUTION	
KEEP OUT OF REACH OF CHILDREN	Dispose of wastes by burying in non-crop lands away from water supplies. Dispose of containers by burying with wastes or burning. Keep out of smoke.
Harmful If Swallowed! Avoid prolonged breath- ing of dust. Avoid contact with eyes, skin and clothing. Wash thoroughly after using. Avoid storage near feed and foodstuffs.	1100.25,1970
NOTE TO PHYSICIAN: Toxaphene is a central nervous system stimulant and may cause convul- sions. There is no specific antidote. Barbituric acid derivatives may be used in treatment.	179-1095

NET WEIGHT 50 us.

Printed in U. S. A.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.





Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGH-BORING CROPS OR NON-CROP AREAS OR USE IN A MAN-NER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UN-DESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE On purchase or use of this product buyer and user agree to the following conditions

WARRANTY Niagara warrants that this product, as of the time of sale by Niagara. (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS NIAGARA MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED

DAMAGES Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to Niagara within 10 days of discovery by buyer or user, failing which huyer or user waives any claim for such damage

EXCUSE FROM NON-CONTROLLABLE CONDITIONS Be cause of certain critical conditions created or incurred by buyer or user or over which Niagara has no contrôl, buyer and user. as a condition of purchase or use, assume responsibility for and release Niagara from, liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

MANUFACTURED IN U.S.A. BY



NIAGARA CHEMICAL DIVISION FMC CORPORATION MIDDLEPORT, NEW YORK



.