

ACCEPTED

APR 7 1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 274-1191

Code: 30301

50

ZIRAM 10 DUST

FUNGICIDE

DIRECTIONS FOR USE

Use as a dust at 40-50 pounds per acre unless stated otherwise. Apply in pre-bloom, blossom, post-bloom and cover applications. Maintain adequate coverage during periods of infection.

ON APPLES (Pacific Northwest and California): for control of Scab.

ON PEARS (Pacific Northwest and California): for control of Scab.

ON ALMONDS (in California): for control of Shot Hole fungus, Brown Rot, Scab and Leaf Blight. May be used until 5 weeks after petal-fall.

ON APRICOTS (California and Oregon): for control of Brown Rot and Shot Hole fungus. May be used thru early cover sprays.

ON PEACHES for control of Brown Rot, Shot Hole fungus and Peach Blight. Remove excessive residue by wiping or brushing.

ON TOMATOES for use as a dust for the control of Anthrachnose and Early Blight use Ziram 10 Dust at the rate of 35 to 25 pounds per acre depending on the size of the plants. For the control of Anthrachnose and Late Blight use Ziram 10 Dust on tomato plants use alternate application of Ziram 10 Dust and C-O-C-S Copodust at intervals of 7 to 10 days during the period when these diseases are a problem.

ON STRAWBERRIES for control of Strawberry Fruit Rot use 35-40 pounds per acre. Make first application at petal-fall and a second application 7-10 days later. Repeat as necessary. Fruit must be protected throughout the growing season. Remove residue by washing if application is made 7 days of harvest.

ON BEANS for control of Rust, Anthrachnose and Sclerotinia use 30 pounds per acre. Apply at first appearance and repeat as necessary. Do not apply beans within 4 days of picking.

Active Ingredient:

†Ziram

By Wt.

10.00%

Inert Ingredients

90.00%

†Zinc Dimethyl dithiocarbamate

100.00%

Zinc expressed as metallic 2.14%

CAUTION

KEEP OUT OF REACH OF CHILDREN

May be Harmful if Swallowed or Inhaled. May cause irritation of eyes, nose, throat and skin. Avoid contact with eyes, skin or clothing. Avoid breathing dust. In case of contact, flush with plenty of water. Foreyes, get medical attention. Keep away from domestic animals, feed and foodstuffs.


USAGE CAUTION
neighboring
time other
injury, ex
occur.

Dealers S

Manufactured in U.S.A. by



AMERICAN CANTON & RICHMOND, CA.
KANSAS CITY, MO. 64108
New York Office: 100 West 40th St., N.Y.



50 Pounds Net Weight

ZIRAM 10 DUST

FUNGICIDE

OR USE

acre unless stated otherwise, post-bloom and cover coverage during periods of

ifornia): for control of Scab.
ifornia): for control of Scab.
control of Shot Hole fungus.
May be used until 5 weeks

for control of Brown Rot
d thru early cover sprays.
Rot, Shot Hole fungus and
side by wiping or brushing.

the control of Anthrachnose
st at the rate of 15 to 25
size of plants. For the
Blight on tomato
ion of Ziram 10 Dust and
f 7 to 10 days during the
a problem.

Strawberry Fruit Rot Use 35-
plication at petal fall and
ater. Repeat as necessary.
ut the growing season. Re-
ation is not required.

Anthrachnose on tomato
a re. Use 10 to 20
sary. Do not use

Active Ingredient:	By Wt.
†Ziram	10.00%
Inert Ingredients	90.00%
†Zinc Dimethyl dithiocarbamate	100.00%
Zinc expressed as metallic 2.14%	

CAUTION
KEEP OUT OF REACH OF CHILDREN.

May be Harmful if Swallowed or Inhaled. May cause irritation of eyes, nose, throat and skin. Avoid contact with eyes, skin or clothing. Avoid breathing dust. In case of contact, flush with plenty of water. For eyes, get medical attention. Keep away from domestic animals, feed and foodstuffs.

Manufactured in U.S.A. by



AMERICAN CANNON & RICHMOND, CALIF.
FARMERS' SUPPLY CO. NEW YORK, N.Y.
Home Office: New York, N.Y.



This product is toxic to fish. Keep out of water.
material. Apply this material to crops strictly in
accordance with directions.

DISPOSAL WARNING: Do not dispose of
this material in any way which may be
harmful to the environment.

Cameraman's Note

Poor Copy

USAGE CAUTION: Do not allow this material to drift onto neighboring crop or non-crop areas or use in a manner or at a time other than in accordance with directions, because plant injury, excessive residues or other undesirable results may occur.

Dealers should Sell in Original Packages Only.