

Code: 30260

Net Weight
50 Pounds

PCNB 40 DUST

DIRECTIONS FOR USE

BEANS: Root and Stem Rot (Rhizoctonia Solani): FURROW: Pole Beans -- Use 6 to 10 1/2 ounces per 1000 feet of row. Bush Beans -- Use 2 to 6 ounces per 1000 feet of row. Dust the planting furrow and covering soil at planting time. Avoid applying to bare seeds. Do not feed treated vines to livestock.

BEANS — SNAP AND DRY: White Mold (Sclerotinia Sclerotiorum): Apply 12 1/2 pounds per acre when disease first appears. Application may be made by air or conventional ground duster. Do not cultivate after application. Dust should cover soil surface as well as plants and soil should remain undisturbed after treatment. Do not apply after first bloom. Do not feed treated vines to livestock.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE AND CAULIFLOWER: Clubroot (Plasmodiophora brassicae) **ROW APPLICATION:** Apply 50 to 100 pounds per acre (3 3/4 to 7 1/2 pounds per 1000 feet of row; Cauliflower (only) — 4 1/2 to 7 pounds per 1000 feet of row) in a 12 to 15 inch band by means of a suitable row applicator, and rototill, to a depth of 4 to 6 inches immediately prior to transplanting. **NOTE:** For clubroot of Cabbage, Cauliflower, Broccoli and Brussels Sprouts, it is very important to thoroughly mix PCNB 40 Dust with the soil. The rate of application depends upon the soil type — the lowest rate is for light soils and the higher rate for heavy soils. **BROADCAST:** Apply 75 to 150 pounds per acre to the soil surface prior to transplanting, disc and cross disc to a depth of 4 to 6 inches.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE: Wire Stem or Black Root (Corticium Solani) — Apply 12 1/2 pounds per 13,100 feet of row in an 8-inch band, centered on the row, immediately prior to seeding. Mix into the top 2 inches of soil.

CAULIFLOWER: Wire Stem or Black Root (Corticium Solani) — Apply 12 1/2 pounds per 13,100 feet of row in an 8-inch band, centered on the row, immediately prior to seeding.

COTTON: Damping-Off, Seed and Seedling Rots (Rhizoctonia): **FURROW APPLICATION:** Apply 5 1/4 pounds per 13,100 ft. of row to the planting furrow. Dust over seed and covering soil at planting time using suitable application equipment. Do not allow the feeding or grazing of cotton foliage by livestock. Do not plant any root crop not registered for PCNB in rotation on PCNB treated soil.

PLANTER BOX APPLICATION — COTTON SEED: Apply to seed through planter box at planting time at the rate of 1 1/2 to 2 1/2 pounds per 100 pounds of seed. Thoroughly mix this product with seeds before planting. This is a supplementary treatment which can be used in addition to regular seed treatment. To insure correct planting rate calibrate planter for proper seeding rate after adding the suggested dosage of this product.

GARLIC: White Rot (Sclerotium Cepivorum) Apply 1 3/4 to 2 1/2 pounds per 1000 feet of to the furrow at the time of planting. A suitable duster should be mounted on the planter and the duster tubes and attachments directed to treat the bottom of the furrow and the "Seed" as it is dropped.

FUNGICIDE

EPA REGISTRATION NO. 279-1250 AA

ACTIVE INGREDIENT:	BY WT.
Pentachloronitrobenzene	40.00%
INERT INGREDIENTS:	60.00%
	100.00%

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if Swallowed! Avoid inhaling dust. Avoid skin contact. This material may cause skin irritation. Wash hands and face thoroughly after using. Keep away from domestic animals, feed or foodstuffs.

It is advisable to wear gloves and a respirator when loading the duster.

For Emergency Assistance Call
716-735-3765

POTATOES (Rhizoctonia): BROADCAST TREATMENT: Use 60 pounds per acre prior to planting. Disc and cross disc to a depth of 4 to 6 inches. **ROW TREATMENT:** Prior to planting time apply 25 pounds per acre, (apply 1 3/4 pounds per 1000 feet of row) by means of suitable row applicator equipment. The fungicide should be thoroughly worked into the top 4 to 6 inches of soil. It is desirable to make treatments when the soil is slightly moist.

NOTE: The rate of application depends upon the type of soil. Light soils should receive the lowest rate and heavy soils the higher rate. It is very important to thoroughly mix PCNB into the tuber-forming zone of the soil.

LAWNS — non-grazed areas: Brown Patch (Rhizoctonia Solani): For Southern or Warm Season Lawns, such as St. Augustine, Bermuda and Similar Grasses. Make one application in the fall or spring at first indication of infection. Apply 1/2 pound per 1000 square feet following application, the treated area should be lightly watered to insure movement through the grass to soil level. If the treated area is subjected to unusually heavy rainfall, or flooded, or if the disease is severe or reappears, the area should be treated 3 to 4 weeks later.


For Northern or Cool Season Lawns Such As Rye Grass, Fescue, Bluegrass and Similar Grasses. Apply 1/2 pound per 1000 square feet at first sign of disease and repeat every 7 to 10 days during periods of hot wet weather. After application, trench or sprinkle lawn to wash chemical down to the soil surface.

NOTE: Under certain growing conditions a temporary discoloration of the grass occasionally may occur. This causes no harm and will disappear in a short time.


ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish. Keep out of lakes, ponds or streams. Do not apply where run-off is likely to occur. Do not apply when weather conditions favor drift from the areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label.

DISPOSE OF WASTES by burying in noncrop land away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.



ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT INTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:
WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.
DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user. Where FMC or user waives any claim for such damages, EXCEPT FOR A NON-CONTROLLABLE CONDITION. Because of certain uncontrollable conditions created or incident to the use of this product, buyer and user, as well as FMC, shall, at purchase or use, assume responsibility for any damage, FMC is not liable for any loss, out of the handling or use of this product attributable to such causes. Buyer and user shall not be held to waive any and all rights contained herein.

3 APR 1974

REGISTRATION DIVISION
PESTICIDE EPA

