

Code: 1047

Gallons Net Contents

TOXAKIL 8.0 MISCIBLE

DIRECTIONS FOR USE

Apply the following recommended rate in sufficient water to thoroughly cover one acre. (By ground apply a minimum of 10 gallons; by air apply sufficient water for thorough coverage but not less than 1 gallon of finished spray per acre). Do not use in undiluted form. Apply when insects first appear and repeat as necessary. May be applied with ground or airplane application equipment. Observe days interval between last application and harvest indicated by number in () following crop.

BEANS (8): Leafhopper, Bean Leafroller, Cowpea Curculio, Thrips, Armyworm, Mexican Bean Beetle, Corn Earworm, Climbing Cutworms, Serpentine Leafminer, Lygus Bug, Grasshopper — Use 2-6 pints per acre. Bean Lycaenid — Use 2 1/2-6 pints per acre. Bean Leaf Skeletonizer, Darkling Ground Beetle, Garden Webworm, Pea Weevil — Use 3-6 pints per acre. No limitation on shelled beans. Do not apply to green or snap beans after pods begin to form. Do not feed treated vines to dairy animals or animals being finished for slaughter.

BROCCOLI (30), BRUSSELS SPROUTS (30), CAULIFLOWER (30), KOHLRABI (30): Imported Cabbageworm, Corn Earworm, Diamondback Moth, Fall Armyworm, Serpentine Leafminer, Thrips — Use 1 1/2-8 pints per acre. Climbing Cutworms — Use 2-8 pints per acre. Flea Beetle — Use 3-8 pints per acre. Do not apply after edible parts begin to form.

CABBAGE (8): Imported Cabbageworm, Corn Earworm, Diamondback Moth Larvae, Fall Armyworm, Serpentine Leafminer, Thrips — Use 1 1/2-4 pints per acre. Climbing Cutworm — Use 2-4 pints per acre. Flea Beetle — Use 2 1/2-4 pints per acre. Do not apply after plants start to form.

CARROTS (8): Climbing Cutworm, Grasshopper — Use 2-5 pints per acre.

CELERY (8): Imported Cabbageworm, Armyworm, Thrips, Serpentine Leafminer — Use 1 1/2-5 pints per acre. Climbing Cutworms — Use 2-5 pints per acre. Celery Leafminer, Vegetable Weevil — Use 4-5 pints per acre. Do not apply after plants start to bunch or after plants are half mature.

CORN (8): Armyworm, Chinch Bug, Webworm, Climbing Cutworm — Use 2 1/2-6 pints per acre. Flea Beetle, Grasshopper — Use 2-6 pints per acre. Do not feed treated forage to dairy animals or animals being finished for slaughter. No limitations on the use of grain.

COTTON (8): Armyworm, Boll Weevil, Bollworm, Lygus Bugs, Cotton Leafworm, Leaf Tier, Darkling Ground Beetle, Climbing Cutworm, Grasshoppers, Plant Bugs — Use 2-8 pints per acre. Stink Bug — Use 4-8 pints per acre. Garden Webworm — Use 3-8 pints per acre. Thrips, Fleahopper, Flea Beetle — Use 3-4-8 pints per acre. If 4 pints per acre or less are used do not graze dairy animals or animals being finished for slaughter in fields treated late in the season. At a rate over 4 pints but not more than 8 pints per acre do not apply after bolls open. Do not feed gin waste to livestock.

EGGPLANT: Armyworm, Blister Beetle, Colorado Potato Beetle, Climbing Cutworms, Flea Beetles, Leafhopper, Serpentine Leafminer — Use 2-6 pints per acre. Cucumber Beetle — Use 3-6 pints per acre. At the 3 pints per acre rate do not apply within 5 days of harvest. At a rate over 3 pints but not more than 6 pints, do not apply after edible parts start to form.

MANUFACTURED IN U.S.A. BY
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
MIDDLEPORT, N. Y. 14105

074 -S

INSECTICIDE

EPA REGISTRATION NO. 279-1390 AA EPA EST 279-TX-1

ACTIVE INGREDIENTS:	BY WT.
*Toxaphene	70.25%
Xylene base aromatic petroleum solvent	22.00%
INERT INGREDIENTS:	7.75%
	100.00%

*Technical Chlorinated Camphene 67% to 68% Chlorine
Contains 8 pounds Toxaphene per gallon.

WARNING

KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed! Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Do not store near feed and food products.

Toxaphene is a central nervous system stimulant. There is no specific antidote. Barbituric acid derivatives may be used for convulsions. Treat symptomatically.

USAGE CAUTIONS


This product is toxic to fish, birds and other wildlife. Keep out of lakes, ponds and streams. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from the areas treated. Do not apply when run-off is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes and containers. Apply this product only as specified on the label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not use or store near heat or open flame. Do not store below 0° F.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROP OR NON-CROPPED AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

Dealers Should Sell in Original Packages Only.



279-1390

LETTUCE (8): Armyworm, Imported Cabbageworm, Cucumber Beetle, Climbing Cutworm, Flea Beetle, Grasshopper. — Use 2-5 pints per acre. Field Crickets — Use 3-5 pints per acre. Do not apply after seedling stage on leaf lettuce. Do not apply after heads begin to form on head lettuce.

ONIONS (8): Climbing Cutworms — Use 2-5 pints per acre. Thrips — Use 1 1/2-5 pints per acre. Do not apply to green or spring onions.

PEPPERS: Armyworm, Blister Beetle, Colorado Potato Beetle, Climbing Cutworm, Hornworm, Serpentine Leafminer — Use 2-6 pints per acre. Flea Beetles — Use 1 1/2-6 pints per acre. Pepper Weevil — Use 3-6 pints per acre. At the 3 pint per acre rate do not apply within 5 days of harvest. At a rate over 3 pints but not more than 6 pints per acre do not apply after peppers begin to form.

SORGHUM: Armyworm, Chinch Bug, Corn Earworm, Climbing Cutworm, Flea Beetle, Webworm — Use 2 1/2-3 pints per acre. Grasshoppers — Use 2-3 pints per acre. When 2 pints rate is used do not apply within 28 days of harvest. When a rate over 2 pints is used do not apply more than once after heads start to form nor within 40 days of harvest. Do not graze dairy animals or animals being finished for slaughter on treated fields. Do not ensile treated forage.

TOMATOES: Armyworm, Blister Beetle, Colorado Potato Beetle, Climbing Cutworm, Flea Beetle, Pinworm, Russet Mite, Serpentine Leafminer, Tomato Fruitworm, Thrips, Hornworm — Use 2-5 pints per acre. Grasshoppers — Use 1 1/2-5 pints per acre. When 1 1/2-2 pints are used do not apply within 1 day of harvest. At a rate over 2 pints per acre do not apply within 3 days of harvest.

For Emergency Assistance Call
716-735-3765

Do not re-use empty 1 and 5 gallon containers. Destroy them by perforating or crushing. Return drum to reconditioner or destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from, liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

FMC Chemicals

Printed in U S A