

Code 30176

Net Contents 28 Gallons

Ethion Summer Oil

Light-Medium

Insecticide-Miticide

For Agricultural or Commercial Use Only

EPA Reg. No. 279-1406 EPA Est.,

Active Ingredients:	By Wt.
*Ethion	2.20%
Petroleum Oil	95.00%
Inert Ingredients:	2.80%
	100.00%

*O,O'-tetraethyl S,S'-methylene bisphosphorodithioate
Contains 0.16 pound Ethion per gallon
Classification—Light Medium
Unadulterated Residue—92.0% Minimum

KEEP OUT OF REACH OF CHILDREN WARNING

For Emergency Assistance Call 716-736-3766.

See Other Panels for Additional Precautionary Information

PRECAUTIONARY STATEMENTS


Hazards to Humans (& Domestic Animals)

Warning

May be Fatal if Swallowed, Inhaled, or Absorbed Through Skin! Do not breathe vapor or spray mist. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Do not contaminate or store near feed and foodstuffs.



FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103



Environmental Hazards

This product is poisonous to fish and wildlife. Keep out of lakes, ponds, and streams. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not apply to or allow this material to drift onto feed or forage crops. Avoid application of oil sprays just prior to or during hot or freezing weather or during rainy weather or dry, hot winds. Avoid double spraying. Plants receiving oil sprays should be in healthy condition, and proper moisture should be present. Use efficient equipment. Do not make an oil spray application within 2 weeks prior to or after an application of sulphur.

Do not allow this material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinse that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. To dispose of container, triple rinse (or equivalent) and offer for recycling, reconditioning or disposal in approved landfill, or bury in a safe place. Do not cut or weld this container. Do not reuse container. Consult Federal, State or local disposal authorities for approved alternative procedures.

Apply when pests first appear and repeat as necessary

Directions for Mixing

With enough water in tank to cover agitator shaft and with shaft in motion, add desired amount of this material for full tank. Pump this mixture through the overflow and back into tank for about two minutes to ensure proper emulsification. Fill with water, adding other desired materials as tank is filling. Keep agitator running at all times.

Almonds: To control Brown Almond Mite, European Red Mite, Two-spotted Spider Mite. Use at the rate of 1 to 2 gallons per 100 gallons of water. If 12 gallons or less of Ethion Summer Oil are applied per acre do not make more than 1 application during the fruiting period. Do not apply within 30 days of harvest. If a rate exceeding 12 gallons but not over 15 gallons per acre is used, do not apply after husk split.

Apples: To control Brown Almond Mite, European Red Mite, McDaniel Mite, Two-spotted Spider Mite. Use 1 to 2 gallons per 100 gallons of water. Do not apply more than 4 times during the fruiting period nor

within 40 days of harvest. If no more than 2 applications are made apply within 20 days of harvest.

Ornamentals, Non-bearing Deck Nursery Stock: For the control of insects. Use at a rate of 1 to 2 gallons per acre coverage applications when insect control is desired.

Peaches: To control European Red Mite, Brown Almond Mite. Use 1 to 2 gallons per acre. Do not apply more than twice during fruit harvest.

Pears: To control Brown Almond Mite, Spotted Spider Mite, Pear Psylla. Use 1 to 2 gallons per acre. Do not apply more than 3 applications during the fruiting period. Do not apply more than 3 treatments during the fruiting period. Do not apply more than 3 treatments during the fruiting period. Do not apply more than 3 treatments during the fruiting period.

Plums and Prunes: To control European Red Mite, Brown Almond Mite, European Red Mite, European Red Mite. Use 1 to 2 gallons per 100 gallons of water. Do not apply more than 3 treatments during the fruiting period. Do not apply more than 3 treatments during the fruiting period. Do not apply more than 3 treatments during the fruiting period.

Dealers Should Sell in

Terms of Sale or Use: On purchase agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, that the use of this product other than as directed on the label will result in the control of the pest. The product is sold as is, with all faults, and use and/or handling and/or storage and/or handling and/or storage is at the user's risk.

Directions and Recommendations: Directions and Recommendations: Use of this product is based upon tests believed to be representative of the pest and the method of application, weather, and other factors in the use of this product. The user is assumed to be familiar with the use of this product and is assumed to be responsible for the results obtained.

Use of Product: FMC's recommendations are based upon tests believed to be representative of the pest and the method of application, weather, and other factors in the use of this product. The user is assumed to be familiar with the use of this product and is assumed to be responsible for the results obtained.

Damages: Buyer's or user's exclusive of warranty or negligence shall be deemed the purchase price paid, less consequential damages.

