

Code: 30416

50 Pounds Net Weight

SEVIN* 5 DUST

DIRECTIONS FOR USE

Apply when insects first appear and repeat as necessary. May be applied with ground or airplane application equipment. Observe days interval between last application and harvest indicated by number in () following the crop.

ALFALFA (0), CLOVER (0), PEANUTS (0): Blister Beetle, Mexican Bean Beetle, Alfalfa Caterpillar, Bean Leaf Beetle, Cucumber Beetle, Leafhoppers, Three-Cornered Alfalfa Hopper, Thrips, Armyworms, Corn Earworm, Grasshoppers, Stink Bugs, Webworms — Use 20 pounds per acre.

APRICOTS (3), PEACHES (1), NECTARINES (3): Oriental Fruit Moth, Peach Twig Borer, Periodical Cicada, Codling Moth, Tussock Moth — Use 60-80 pounds per acre.

ASPARAGUS (1): Asparagus Beetle on Seedlings or Spears — Use 20-40 pounds per acre. Apply during cutting season. Do not repeat application within 3 days. Asparagus Beetle, Apache Cicada on ferns or brush growth in the post harvest period — Use 40-80 pounds per acre.

BEANS (0): Mexican Bean Beetle — Use 10 pounds per acre. Bean Leaf Beetle, Cucumber Beetles, Flea Beetles, Leafhoppers, Western Bean Cutworm — Use 20 pounds per acre. Armyworms, Corn Earworm, Stink Bugs, Tarnished Plant Bug — Use 20-30 pounds per acre. California: Lygus Bug, Stink Bug, Lima Bean Pod Borer — Use 40 pounds per acre.

BLACKBERRIES (7), BOYSENBERRIES (7), DEWBERRIES (7), LOGANBERRIES (7), RASPBERRIES (7): Leafrollers, including Omnivorcus Leafroller, Sawfly, Climbing Cutworms, Snowy Tree Cricket (California), Orange Tortrix (Oregon), Leafhoppers, Raspberry Aphids — Use 40 pounds per acre.

CABBAGE (3), BROCCOLI (3), BRUSSELS SPROUTS (3), CAULIFLOWER (3), KOHLRABI (3): Flea Beetles, Harlequin Bugs — Use 10-20 pounds per acre. Armyworm, Imported Cabbageworm, Corn Earworm, Cabbage Looper (small) — Use 20-40 pounds per acre. For small Cabbage Looper control apply in a 5-7 day application schedule.

CARROTS (0), CHINESE CABBAGE (14), COLLARDS (14), DANDELION (14), ENDIVE (14), BEETS (Roots 3) (Tops 14), HORSERADISH (3), KALE (14), LETTUCE (Head 3) (Leaf 14), MUSTARD GREENS (14), PARSLEY (14), PARSNIPS (3), RADISHES (3), RUTABAGAS (3), SALSIFY (Roots 3) (Tops 14), SPINACH (14), SWISS CHARD (14), TURNIPS (Roots 3) (Tops 14): Flea Beetle, Harlequin Bugs, Leafhoppers — Use 10-20 pounds per acre. Six-Spotted Leafhopper, Armyworm, Corn Earworm, Tarnished Plant Bug, Stink Bugs, Cabbage Looper (small) — Use 20 pounds per acre. For small Cabbage Looper control apply in a 5-7 day application schedule.

CHERRIES (1), PLUMS (1), PRUNES (1): Cherry Fruit Fly, Peach Twig Borer, Leafhopper, Codling Moth, Orange Tortrix, Western Tussock Moth — Use 60-80 pounds per acre. For Codling Moth, apply at petal fall and every 10 to 14 days thereafter until control is achieved.

063W

INSECTICIDE

EPA REGISTRATION NO. 279-1418 AA

ACTIVE INGREDIENT:	BY WT.
Carbaryl (1-naphthyl N-methylcarbamate)	5.00%
INERT INGREDIENTS:	95.00%
	100.00%

*Trade-mark Union Carbide Corporation for the active ingredient Carbaryl.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Skin contact may be harmful. Avoid contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

ACCEPTED

USAGE CAUTIONS

DEC 5 1973

Keep out of lakes, ponds and streams.

This product is highly toxic to bees exposed on crops. Protective information may be obtained from Agricultural Extension Service.

Some phytotoxicity may occur on tender foliage in presence of rain or high humidity of several days duration following application.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

For Emergency Assistance Call
716-735-3765

CORN (0): Corn Earworm — Use 40 pounds per acre. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when silks appear and continuing until silks begin to dry. Three or more applications will be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. Application of this product to the tassal region of corn during the pollen shed period will seriously reduce bee populations.

COTTON (0): Early Season: Thrips, Flea Beetles, Cotton Fleahopper, Cotton Leafworm, Striped Blister Beetle — Use 10-20 pounds per acre. Mid and Late Season: Boll Weevil, Bollworm, Fall Armyworm, Cotton Leaf Perforator, Leafrollers, Leafhoppers, Tarnished Plant Bug, Lygus Bug, Aphids (suppression) — Use 20-40 pounds per acre. Pink Bollworm — Use 35-50 pounds per acre. Stink Bug, Salt Marsh Caterpillar — Use 40-50 pounds per acre. Grasshoppers — Use 10-30 pounds per acre.

FORAGE GRASSES, PASTURE: Armyworms, Grasshoppers, Thrips (seed crop) — Use 20-30 pounds per acre.

GRAPES (0): Grape Leafroller, Grape Leafhoppers, Western Grape Leaf Skeletonizer — Use 20-40 pounds per acre. Apply just before first brood Leafroller Larvae emerge from rolls and as needed for Leafhoppers. This product may be used on wild or cultivated grapes. Ground application equipment preferred.

OKRA (0): Corn Earworm, Stink Bugs — Use 20-40 pounds per acre.

PEAS (0): Armyworms, Grasshoppers — Use 20-30 pounds per acre. Leafhoppers, Colorado Potato Beetle — Use 20 pounds per acre.

RICE (14): Leafhoppers, Armyworms — Use 30 pounds per acre. Tadpole Shrimp (California) — Use 40 pounds per acre. Do not apply within 14 days before or after applications of the herbicide propanil (Stam F-34 or Rogue) as injury may result.

SORGHUM (MILO AND GRAIN SORGHUM) (Grain 21) (Forage 0): Armyworm, Corn Earworm, Stink Bugs, Webworm — Use 20-40 pounds per acre. Sorghum Midge — Use 30 pounds per acre. Treatment for Sorghum Midge control should be made 3 to 4 days after heads have emerged from boot.

STRAWBERRIES (1): Meadow Spittlebug, Strawberry Leafroller, Strawberry Weevil — Use 20-40 pounds per acre.

TOMATO (0), EGGPLANT (0), PEPPERS (0): Colorado Potato Beetle, Flea Beetles, Leafhoppers — Use 10-20 pounds per acre. Fall Armyworm, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bugs, Stink Bugs — Use 20-40 pounds per acre.

Printed in U.S.A.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED**

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from, liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

279-1418
ACCEPTED
DEC 5 1973
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
OF 1947 THIS PRODUCT IS REGISTERED
EPA REG. NO. 100-100000
SUBJECT TO APPROVED COMMENTS

Manufactured in U.S.A. by:
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
Middleport, N.Y. 14105

FMC
Chemicals