

Code: 30428

50 Pounds Net Weight

# TRITHION<sup>®</sup> SULPHUR 3-50 DUST

## DIRECTIONS FOR USE

Apply when pests or disease first appear and repeat as necessary. For control beyond the usual limitations of this product use a suitable material. May be applied with ground or aerial application equipment. Observe days interval between last application and harvest as listed by the manufacturer following the crop.

**APPLES (30), CRABAPPLES (30), PEAR (30), QUINCE (30):** Scab, Pear and Crab Apple Aphid, European Red Mite, Rosy Apple Aphid, Woolly Apple Aphid, Two Spotted Mites, Curling Mites. Use 40 pounds per acre.

**ALMONDS, PLUMS (30), PRUNES (30):** Brown Rat Aphid, Orchard Mites. Use 40 to 60 pounds per acre. If almond hulls are used as feed do not apply within 7 days of harvest.

**BEEF TOPS (21), SPINACH (21):** Spider mite, Leaf miner, Two Spotted Mite, Bean Aphid. Use 15 to 40 pounds per acre.

**CHERRIES (30), NECTARINES (30), PEACHES (30):** Brown Rat Aphids, Orchard Mites, Flower Mite. Use 40 to 60 pounds per acre.

**CITRUS (14): (Grapefruit, Lemons, Limes, Oranges, Tangelos, Tangerines):** Citrus Red Mite, Citrus Rust Mite, Six Spotted Mite, Texas Citrus Mite, Yuma Mite. Use 35 to 100 pounds per acre. Allow 30 days between applications. Do not apply to Kumquats or Citron.

**COTTON (0):** Cotton Aphid, Cotton Leaf Perforator, Cotton Leafhopper, Cotton Leaf-worm, Curculionid Mite, Tripping Mite, Two Spotted Mite. Use 20 to 30 pounds per acre. Aphids, Cotton Leaf Perforator, and the 30 pound per acre. Do not graze dairy animals or meat animals treated with it.

**EGGPLANTS (7):** Spider mite, Leaf miner, Bean Leafhopper, Potato Flea Beetle, Purple Mite, Tobacco Mite. Use 20 to 40 pounds per acre.

**FIGS (7):** Two Spotted Mite. Use 40 to 60 pounds per acre.

**GARDEN BEETS:** Bean Aphid, Leafhoppers, Serpentine Leafminer, Two Spotted Mites, Flea Beetle. Use 10 to 40 pounds per acre. Do not apply within 21 days of harvest if the plants are used as feed for livestock.

**GRAPES (30):** Flower Mite, Grape Leafhopper, Two Spotted Mite. Use 20 to 40 pounds per acre. Aerial application equipment preferred.

**ORNAMENTALS (Camelias, Carnations, Gladiolus, Violets, Evergreens, Deciduous Shrubs):** Red Spider Mite, Flower Mites, Aphids, and Mites. Use 20 pounds per acre. Do not graze dairy animals or meat animals treated with it.

**PEAS (7):** Bean Aphid, Serpentine Leafminer, Two Spotted Mite. Use 20 to 40 pounds per acre. Do not graze dairy animals or meat animals treated with it.

**PEPPERS (7), PIMENTO (7):** Serpentine Leafminer, Flea Beetle, Potato Leafhopper, Tobacco Mite, Flower Mite. Use 20 to 40 pounds per acre.

## FUNGICIDE-INSECTICIDE-MITICIDE

EPA REGISTRATION NO. 279-1437-AA

ACTIVE INGREDIENTS	BY WT.
Carbophenothion, Sulfophenothion, Lithio-methyl, 0,0-Diethyl phosphorothioate	3.00%
Sulphur	50.00%
INERT INGREDIENTS	47.00%
	100.00%

Trithion is a registered trademark of Stauffer Chemical Company for its insecticide/miticide.

## WARNING

### KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed, inhaled, or absorbed through skin. Do not get in eyes, on skin, or on clothes. Do not breathe dust. In case of contact, wash immediately with copious water. Wear clean clothing. Wash clothing in hot water with soap and hot water before reuse. Do not store near feed and food products.

Trithion may cause drowsiness, dizziness, Atropine is antidote. 2 PAM is antidote for atropine. Do not store in combination with atropine.

**ACCEPTED**

NOV 19 1973

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND MITICIDE ACT  
FOR ECONOMIC PEST CONTROL  
REGISTERED UNDER NO. 279-1437-AA  
SUBJECT TO ATTACHED COMMENTS.

**SNAP BEANS, LIMA BEANS, SOYBEANS (IN POD):** Powder, Mite, Leafhopper, Bean Aphid, Mexican Bean Beetle, Serpentine Leafminer, Potato Leafhopper, Two Spotted Mites. Use 20 to 40 pounds per acre. On Snap beans do not apply within 7 days of harvest. Do not feed treated forage to livestock. On beans at the 20 pounds per acre rate do not apply within 7 days of harvest. Do not apply more than twice per season. Do not graze or cut vetch for hay within 21 days after application. At a rate above 20 pounds per acre do not apply within 7 days of harvest. Do not feed treated forage to livestock. Sulphur may injure certain varieties, check tolerance to sulphur prior to use.

**STRAWBERRIES (3):** Flower Mite, Two Spotted Mites, Strawberry Aphid. Use 20 to 40 pounds per acre. Do not apply while fruit is present.

**SUGAR BEETS (14):** Bean Aphid, Serpentine Leafminer, Potato Flea Beetle, Two Spotted Mite. Use 20 to 30 pounds per acre.

**TOMATOES (7):** Serpentine Leafminer, Potato Psyllid, Potato Flea Beetle, Potato Leafhopper, Tomato Russet Mite, Tripping Mite, Two Spotted Mites. Use 20 to 40 pounds per acre.

## USAGE CAUTIONS

This product is toxic to fish and wildlife. Keep out of lakes, ponds, and streams. Do not feed to animals. Do not use in areas where it may be hazardous. Do not contaminate water, or cleaning equipment, or disposal of wastes. Apply this product only as directed on the label.

This product is toxic to birds and should not be applied when birds are present in the treated area.


Applications of this product containing Sulphur should not be made during or immediately after periods of excessive high temperatures, or with a relative humidity of 80% or higher.

Do not apply when weather conditions favor drift from the application area. Do not apply when wind is blowing toward you.

DO NOT FEED WASTES. Equipment used in application should be washed with water before being used for any other purpose.

For Emergency Assistance Call  
716-735-3765

**ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.**



**Dealers Should Sell in Original Packages Only.**

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

**TERMS OF SALE OR USE:** On purchase or use of this product buyer and user agree to the following conditions:

**WARRANTY:** FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

**DAMAGES:** Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

**EXCUSE FROM NON-CONTROLLABLE CONDITIONS:** Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured In U.S.A. by:  
FMC CORPORATION  
AGRICULTURAL CHEMICAL DIVISION  
Middleport, N.Y. 14105

**FMC**  
Chemicals