



Code 30428

Pounds Net Weight

Trithion[®] Sulphur 3-50 Dust

Fungicide-Insecticide-Miticide

For Agricultural or Commercial Use Only

EPA Reg. No. 279-1437AA

EPA Est., 279-

Active Ingredients:	By Wt.
Carbophenothion: S-[(p-Chlorophenylthio) methyl] 0,0-diethyl phosphorodithioate . . .	3.00%
Sulphur	50.00%
Inert Ingredients:	47.00%
	100.00%

Trithion is a registered trademark of Stauffer Chemical Company for an insecticide-miticide

KEEP OUT OF REACH OF CHILDREN WARNING

Note to Physician

Trithion may cause cholinesterase inhibition. Atropine is antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine.

For Emergency Assistance Call 716-735-3765.

See Other Panels for Additional Precautionary Information.

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)


Warning

May be fatal if swallowed, inhaled, or absorbed through skin. Do not get in eyes, on skin or on clothing. Do not breathe dust. In case of contact, wash immediately with soap and water. Wear clean clothing. Wash all contaminated clothing with soap and hot water before reuse.

FMC

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

PA 8/80



Environmental Hazards

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Applications of materials containing Sulphur should not be made just prior to or during periods of excessively high temperatures or within 10 days to two weeks of an oil spray.

Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide spray mixture or rinseate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of container in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Consult Federal, State, or local disposal authorities for approved alternative procedures such as limited open burning.

Apply when pests or disease first appear and repeat as necessary. For control beyond the usage limitations of this product use a suitable material. May be applied with ground or airplane application equipment. Observe days interval between last application and harvest indicated by the number in () following the crop.

Apples (30), Crabapples (30), Pears (30), Quince (30): Scab, Powdery Mildew, Apple Aphid, European Red Mite, Rosy Apple Aphid, Woolly Apple Aphid, Twospotted Spider Mite, Codling Moth—Use 40 pounds per acre.

Almonds, Plums (30), Prunes (30): Brown Rot, Aphid, Orchard Mites—Use 40 to 60 pounds per acre. If almond hulls are used as test do not apply within 7 days of harvest.

Beet Tops (21), Spinach (21): Serpentine Leafminer, Twospotted Spider Mite, Bean Aphid—Use 15 to 40 pounds per acre.

Cherries (30), Nectarines (30), Peaches (30): Brown Rot, Aphid, Orchard Mites, Powdery Mildew—Use 40 to 60 pounds per acre.

Citrus (Grapefruit, Lemons, Limes, Oranges, Tangerines, Tangerines) (14): Citrus Red Mite, Citrus Rust Mite, Sixspotted Mite, Texas Citrus Mite, Yuma Mite—Use 35 to 100 pounds per acre. Allow 30 days between applications. Do not apply to Kumquats or Citron.

Cotton (0): Cotton Aphid, Cotton Leafhopper, Cotton Leafworm, Cinnabar Mite, Tropical Mite, Twospotted Spider Mite—Use 20 to 30 pounds per acre. Aphid, Cotton Leafperforator—Use 30 pounds per acre. Do not graze dairy animals or meat animals in treated areas.

279-1437

Eggplants (7): Serpentine Leafminer, Potato Leafhopper, Potato Flea Beetle, Tropical Mite, Twospotted Spider Mite—Use 20 to 40 pounds per acre

Figs (7): Twospotted Spider Mite—Use 40 to 80 pounds per acre

Garden Beets: Bean Aphid, Leafhopper, Serpentine Leafminer, Twospotted Spider Mite, Potato Flea Beetle—Use 15 to 40 pounds per acre. Do not apply within 21 days of harvest if tops are to be used as food or feed

Grapes (30): Powdery Mildew, Grape Leafhopper, Twospotted Spider Mite—Use 20 to 33 pounds per acre. Ground application equipment preferred

Ornamentals (Camellias, Carnations, Gladiolus, Violets, Evergreens, Deciduous Shrubs): Rust, Powdery Mildew, Spider Mite, Aphid—Use 15 to 25 pounds per acre, or dust to cover

Peas (7): Pea Aphid, Serpentine Leafminer, Twospotted Spider Mite—Use 20 to 40 pounds per acre. Do not feed treated forage to livestock

Pecans, Walnuts: Aphid, Orchard Mite—Use 40 to 60 pounds per acre. Do not use after husks open

Peppers (7), Pimientos (7): Serpentine Leafminer, Potato Flea Beetle, Potato Leafhopper, Tropical Mite, Twospotted Spider Mite—Use 20 to 40 pounds per acre.

Snap Beans (7), Lima Beans (7), Soybeans (In Pod) (7): Powdery Mildew, Rust, Bean Aphid, Mexican Bean Beetle, Serpentine Leafminer, Potato Leafhopper, Twospotted Spider Mite—Use 20 to 40 pounds per acre. On Soybeans do not feed treated forage to livestock. On Beans at the 20 pound per acre rate do not apply more than twice per season. Do not graze or cut vines for hay within 21 days after application. At a rate above 20 pounds per acre do not feed treated forage to livestock. Sulphur may injure certain varieties, check tolerance to sulphur prior to use.

Beans (Shelled): Powdery Mildew, Rust, Bean Aphid, Mexican Bean Beetle, Serpentine Leafminer, Twospotted Spider Mite, Potato Leafhopper—Use 20-40 pounds per acre. At the 20 pound per acre rate do not apply more than twice per season. Do not graze or cut vines for hay within 21 days after application. Above the 20 pound per acre rate do not feed treated vines or hulls to livestock. Sulphur may injure certain varieties, check tolerance to sulphur prior to use

Strawberries (3): Powdery Mildew, Twospotted Spider Mite, Strawberry Aphid—Use 20 to 40 pounds per acre. Do not apply while fruit is present.

Sugar Beets (14): Bean Aphid, Serpentine Leafminer, Potato Flea Beetle, Twospotted Spider Mite—Use 20 to 30 pounds per acre

Tomatoes (7): Serpentine Leafminer, Potato Psyllid, Potato Flea Beetle, Potato Leafhopper, Tomato Russet Mite, Tropical Mite, Twospotted Spider Mite—Use 20 to 40 pounds per acre.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages

OF - DOC - NOT AVAILABLE

