

Code: 30451

Net Weight
50 Pounds

KOLODUST® CAPTAN 50-10 DUST

DIRECTIONS FOR USE

May be applied by ground or air.

CHERRIES: Brown Rot, Cherry Leaf Spot — Use 30 to 50 pounds per acre. Apply at first sign of disease. Repeat at regular intervals as necessary to maintain control.

GRAPES: Powdery Mildew — Use 20 to 30 pounds per acre. Treatment should be started after growth starts in the spring and be repeated as necessary. Ground application preferred. Botrytis Rot — Use 40 to 50 pounds per acre. Apply twice before bloom and once just after bloom. Repeat periodically to make 3 cover applications before bunches close. Apply to every row and direct toward the clusters. To be effective, dusts must be applied before rains. Whenever rain threatens apply an additional dust. Applications made following rainfall have been of very little benefit in controlling slip skin.

PEACHES, NECTARINES: Brown Rot, Powdery Mildew, Peach Scab — Use 50 pounds per acre. Apply at first sign of disease. Repeat at regular intervals as necessary to maintain control.

PLUMS, PRUNES: Brown Rot, Prune Rust — Use 40 to 50 pounds per acre. Apply at first sign of disease. Repeat at regular intervals as necessary to maintain control.

STRAWBERRIES: Grey Mold, Powdery Mildew — Use 20 to 30 pounds per acre. Apply at first sign of disease. Repeat at regular intervals as necessary to maintain control.

POTATOES: Early and Late Blight — Use 25 to 30 pounds per acre. Begin applications when plants are 2 to 4 inches high and repeat at 5 to 7 day intervals.

TOMATOES: Early Blight, Late Blight, Anthracnose, Septoria Leaf Spot, Russet Mite — **Young Plants** — Use 25 to 30 pounds per acre. **Mature Plants** — Use 35 to 40 pounds per acre. Begin applications when disease or mites appear and repeat as necessary.

ORNAMENTALS: **Roses** — Black Spot, Powdery Mildew and Rust. **Carnations** — Powdery Mildew, Rust and Leaf Spot. **Chrysanthemums** — Flower Blight, Powdery Mildew and Rust; Apply at first sign of disease and repeat as necessary. Dust for complete coverage.

FUNGICIDE

E.P.A. REGISTRATION NO. 279-1445 AA

ACTIVE INGREDIENTS:	BY WT.
*Captan	10.00%
Sulphur	42.00%
INERT INGREDIENTS:	48.00%
	100.00%

*N-[(Trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide

U.S. Pat. Nos. 2,553,770 - 2,553,776 - 2,553,771

USAGE CAUTIONS

Applications of materials containing Sulfur should not be made just prior to or during periods of excessively high temperatures or within 10 days to two weeks of an oil spray.

Do not allow this material to drift onto sulfur sensitive crops.

This product contains Bentonite Sulfur which is made by the adsorption of fused sulfur into Bentonite in the proportions of thirty parts sulfur and seventy parts Bentonite. A portion of the sulfur particles are very minute, some of them colloidal. The process of manufacture and the character of the sulfur particles impart to the product distinctive properties with respect to toxicity to certain fungi, persistence on plants, suspensibility in spray tank and safety to host plants.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid inhalation of dust. Avoid contact with skin. Avoid contamination of feed or foodstuffs.


ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment, or disposal of wastes and containers. Do not apply when weather conditions favor drift from the areas treated. Do not apply when run-off is likely to occur. Apply this product only as specified on the label.

Do not use this product in a manner or at a time other than in accordance with label directions because animal, plant, or crop injury, illegal residues or other undesirable results may occur.

DISPOSE OF WASTES by burying in non-crop land away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

For Emergency Assistance Call
716-735-3765



ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



279-1445

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured in U.S.A. by:
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
Middleport, N.Y. 14105

FMC
Chemicals