

CODE

20102

NET CONTENTS

MISCIBLE CITRUS OIL

5 Gallons

INSECTICIDE - MITICIDE - SCALACIDE

DIRECTIONS FOR USE

Niagara Miscible Citrus Oil is a miscible spray oil intended for use on citrus for the control of scale insects (snow, soft brown, purple and Florida red), spider mites and whiteflies. MISCIBLE CITRUS OIL is suggested for use at a concentration of 0.7% to 0.9% (3.5 to 4.5 gallons per 500 gallons water) actual oil. For use in combination with parathion for control of scale insects a concentration of 0.4% to 0.6% (2.0 to 3.0 gallons per 500 gallons water) actual oil plus 5 pounds parathion 15% wettable powder (or equivalent) or 2.5 - 3.75 pts. Ethion 4 Miscible or 5 - 7.5 pounds Ethion 25 Wettable Powder per 500 gallons of water is suggested. The lower rates are for light to moderate scale, spider mite and whitefly infestations while the higher rates are for moderate to heavy scale infestations. All dosage rates given are for 500 gallons finished spray. Note: Do not apply Parathion plus oil mixtures within 14 days of harvest nor more than 3 applications of such mixtures to any one crop.

U.S.D.A. Reg. No. 279-1448

ACTIVE INGREDIENTS:

Petroleum Oil ----- 99.2%

INERT INGREDIENTS: ----- 0.8%

100.0%

UNSULFONATED RESIDUE - 92.0%
MINIMUM

DO NOT USE THIS MATERIAL IN SPRAY EQUIPMENT WITH POOR AGITATION.

Do not use this material if it does not mix with water. Do not use in undiluted form.

CAUTION

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed.

To mix, fill the tank 1/4 to 1/3 full with water and with agitator in motion add the required amount of this material. (If other materials are to be used in the same tank, it is suggested that they be added first, for ease in mixing.) Finish filling tank with water and keep agitated. If an injector is used, first fill tank 1/4 to 1/3 full with water with engine running to provide good pressure. Then place suction hose in measured amount of spray oil and suck into tank. Finish filling tank with water.

For best results with this product the following precautions should be observed: Do not pour this material directly into empty tank; avoid spraying during or immediately before periods of hot or near freezing weather, hot dry winds, periods of drought or when trees are wilting or near wilting; avoid over spraying or double spraying; do not apply within 30 days before or after an application of sulphur. Trees should be sprayed with oil only when in a vigorous condition when humidity is high, and when moisture is plentiful.

MANUFACTURED IN U.S.A. BY



Jacksonville, Florida