

Code: 30465

Net Weight
50 Pounds

CAPTAN 10 DUST



DIRECTIONS FOR USE

Apply to give thorough coverage to upper and lower leaf surfaces. May be applied by ground or air. Observe days interval between application and harvest indicated by number in () following the crop.

ALMONDS (12): Brown Rot, Shot Hole, Scab, Leaf Blight — Use 25 to 50 pounds per acre. Apply at popcorn, bloom, petal fall stages and up to 5 weeks after petal fall. When applied after petal fall, do not feed almond hulls to dairy animals or animals being finished for slaughter.

APPLES (0): Apple Scab — Use 20 to 80 pounds per acre. Apply in pre-bloom, blossom, petal fall, and first cover periods. For secondary infection use 11 to 25 pounds per acre and apply in cover periods. Brooks Fruit Spot, Flyspeck — Use 11 to 37½ pounds per acre and apply in cover periods. Bitter Rot, Black Rot, Black Pox, Botryosphaeria — Use 20 to 37½ pounds per acre. Apply in late cover and preharvest periods.

APRICOTS (0): Brown Rot, Jacket Rot — Use 50 pounds per acre. Apply at red bud stage and repeat at 75 percent petal fall.

CHERRIES (0): Cherry Leaf Spot — Use 30 to 40 pounds per acre. Apply in pink, full bloom, petal fall and first cover periods. Brown Rot — Use 11 to 50 pounds per acre. Apply in pink, full bloom, petal fall, shuck shed and cover periods.

GRAPES (0): Botrytis Rot (Bunch Rot) — Use 40 to 50 pounds per acre. Apply twice before bloom and once just after bloom. Repeat periodically to make 3 cover applications before bunches close. Apply to every row and direct dust toward the clusters. To be effective, dusts must be applied before rains. Whenever rain threatens apply an additional dust. Applications made following rainfall have been of very little benefit in controlling slip skin. Ground application equipment preferred.

GRAPES: Mold control for fruit on drying trays in fields *Penicillium* spp., *Aspergillus* spp., *Botrytis Cinerea*, *Rhizopus* spp. — Use 10 pounds per acre. Apply to fruit on trays for protection of fruit and prevention of above molds. Apply to give even coverage. During damp weather make a second application after trays are turned. Apply this dust by hand or ground dusters only.

MANGOS (0): Cercospora Spot or Blotch — Use 50 to 70 pounds per acre. Begin applications at first sign of disease. Repeat at regular intervals to maintain control.

PEACHES (0), NECTARINES (0): Brown Rot, Peach Scab — Use 50 pounds per acre. Make 2 to 3 prebloom applications. Repeat at petal fall making 3 to 4 cover sprays as necessary. Make pre-harvest application where necessary.

PEARS (0): Pear Scab (except Pacific Northwest) — Use 20 to 30 pounds per acre. Make first applications during early finger and petal fall stages. Repeat applications for each infection period or at weekly intervals as long as necessary. Russeting may be produced on Bosc Pears. Do not use on D'Anjou Pears.

PLUMS (0), PRUNES (0): Brown Rot — Use 20 to 50 pounds per acre. Apply at redbud and 75 percent bloom, full bloom, petal-fall and in cover sprays as necessary. Thorough coverage is essential for maximum control.


FUNGICIDE

EPA REGISTRATION NO. 279-1449 AA

ACTIVE INGREDIENT: *†*Captan 10.0%
INERT INGREDIENTS: 90.0%

*†*N-[(Trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide

U.S. Pat. Nos. 2,553,770 — 2,553,771 — 2,553,776



CAUTION

KEEP OUT OF REACH OF CHILDREN

Avoid inhalation of dust. Avoid contact with skin. Avoid contamination of feed or foodstuffs.

For Emergency Assistance Call
716-735-3765

ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply when runoff is likely to occur. Apply this product only as specified on the label.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Empty bags should be buried with wastes or burned. Keep out of smoke.

STRAWBERRIES (0): Botrytis Rot — Use 20 to 40 pounds per acre. Make first application when new growth starts in spring before fruit starts to form. Repeat applications weekly. **Severe Infection Conditions:** Continue applications through harvest period, treating immediately after each picking.

BEANS, PEAS (0): Damping-off, Root Rot — Use 60 pounds per acre. Work into top 3 or 4 inches of soil before planting.

CANTALOUPE (0), CUCUMBERS (0), PUMPKINS (0), SQUASH (Summer - Winter) (0), WATERMELON (0): Downy Mildew, Angular Leaf Spot, Anthracnose — Use 20 to 25 pounds per acre on young plants, increasing to 30 to 35 pounds per acre on mature plants. Apply at first sign of disease. Repeat at 5 to 7 day intervals until danger of infection is past.

CELERY (Field and Seed Bed only): Pink Rot, Damping-Off — Use 35 to 65 pounds per acre. Apply when all seeds have germinated. Repeat as necessary.

POTATOES: Early and Late Blight — Use 17½ to 30 pounds per acre. Begin applications when plants are 2 to 4 inches high and repeat at 5 to 7 day intervals as long as disease threatens.

POTATOES: (Seed Pieces): Seed Rot and Damping-Off — Dust at 7½ ounces per bushel cut seed pieces (12 ozs or ¾ pound to 100 pounds cut seed pieces. Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days the treated pieces should be stored for 2 days in open crates before bagging.

TOMATOES: Early Blight, Late Blight, Anthracnose, Septoria Leaf Spot — Young Plants — Use 25 to 30 pounds per acre. Mature Plants — Use 35 to 50 pounds per acre. Begin applications at first sign of disease. Repeat at 5 to 7 day intervals.

ORNAMENTALS: ROSES — Black Spot, CARNATIONS — Rust and Leaf Spot — CHRYSANTHEMUMS — Flower Blight — Use 22½ to 30 pounds per acre. Apply at first sign of disease to give even coverage of upper and lower leaf surfaces. Repeat at 7 to 10 day intervals as necessary.

SOIL AND GREENHOUSE BENCH TREATMENT: Pre-planting for Damping-Off and Root Rot Diseases on Seedling or Transplants of Roses (and other shrubs and trees), many Flowers and Vegetables (such as Beans, Peas, Cucurbits) and Lawn Seedbeds — Use 1½ to 2 pounds per 1000 sq. ft. (50 to 60 pounds per acre). Work into upper 3 to 4 inches of soil before planting.

STOCK: Botrytis Rot — Use 40 to 50 pounds per acre. Apply when disease appears. Repeat at weekly intervals as needed.

Turf and Lawn: Brown Patch, Leaf Spot, Damping-Off, Root Rot — Use 1 to 2 pounds per 1000 sq. ft. Begin treatment before infection occurs. Repeat at 7 to 10 day intervals throughout summer. Do not graze treated areas. Do not feed clippings to livestock.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



279-1449

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.



Camerman's Note
Poor Copy

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured in U.S.A. by:
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
Middleport, N.Y. 14105

FMC
Chemicals