

Code: 30465

Net Weight

Captan 10 Dust

Fungicide

For Agricultural or Commercial Use Only

EPA Reg. No. 279-1449 ZA

EPA Est., 279-

Active Ingredient:	By Wt.
Captan: N-[(Trichloromethyl) thio]-4-cyclohexene-1,2-dicarboximide	9.50%
Related Derivatives	0.50%
Inert Ingredients:	90.00%
	100.00%

**KEEP OUT OF REACH OF CHILDREN
DANGER**

PELIGRO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

Statement of Practical Treatment


In case of eye contact: immediately flush eyes with fresh water for 15 minutes and see a doctor.

If swallowed: give a large amount of water to drink, make person vomit, and call a doctor.

If on skin: wash with soap and water.

For Emergency Assistance Call 716-735-3765

See other panels for additional precautionary information.



FMC

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

PA 683-R
PA 384-R

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

Danger

Causes irreversible eye damage. May cause allergic skin reactions. Harmful if swallowed or inhaled. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing and breathing of dust. Remove and wash contaminated clothing before reuse.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from the areas treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Do not enter treated areas without protective clothing until dusts have settled.

Protective clothing means, at least, a hat or other suitable head covering, a long sleeved shirt and long legged trousers or a coverall type garment (all of closely woven fabric covering the body, including the arms and legs), shoes and socks.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Inform workers of areas or fields that may not be entered without specific protective clothing, period of time field must be vacated and appropriate actions to take in case of accidental exposure. An example of such information is given under written warnings. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information:

DANGER

Area treated with Captan 10 Dust on (insert date). Do not enter without appropriate protective clothing until dusts have settled (insert State Department of Agriculture's re-entry interval, if more restrictive).

In case of accidental exposure to pesticide spray or dust, wash the skin thoroughly with soap and water. Remove contaminated clothing and wash before reuse. If in eyes, flush with plenty of water. If inhaled, go to an area where the pesticide has not been applied. Get medical attention if needed.

STORAGE AND DISPOSAL

Pesticide Storage

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, fold and roll back bags, clamp and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. Call FMC collect. (716) 735-3765.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal

Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

(3)

219-1447

Apply to give thorough coverage to upper and lower leaf surfaces. May be applied by ground or air. Observe days interval between application and harvest indicated by number in () following the crop.

Almonds (12): Brown Rot, Shot Hole, Scab, Leaf Blight—Use 25 to 50 pounds per acre. Apply at popcorn, bloom, petal fall stages and up to 5 weeks after petal fall. When applied after petal fall, do not feed almond hulls to dairy animals or animals being finished for slaughter.

Apples (0): Apple Scab—Use 20 to 80 pounds per acre. Apply in pre-bloom, blossom, petal fall, and first cover periods. For secondary infection use 11 to 25 pounds per acre and apply in cover periods. Brooks Fruit Spot, Flyspeck—Use 11 to 37½ pounds per acre and apply in cover periods. Bitter Rot, Black Rot, Black Pox, Botryospheria—Use 20 to 37½ pounds per acre. Apply in late cover and preharvest periods.

Apricots (0): Brown Rot, Jacket Rot—Use 50 pounds per acre. Apply at red bud stage and repeat at 75 percent petal fall.

Cherries (0): Cherry Leaf Spot—Use 30 to 40 pounds per acre. Apply in pink, full bloom, petal fall and first cover periods. Brown Rot—Use 11 to 50 pounds per acre. Apply in pink, full bloom, petal fall, shuck shed and cover periods.

Grapes (0): Botrytis Rot (Bunch Rot)—Use 40 to 50 pounds per acre. Apply twice before bloom, and once just after bloom. Repeat periodically to give 3 cover applications before bunches close. Apply to every row and direct dust toward the clusters. To be effective, dust must be applied before rains. Whenever rain threatens apply an additional dust. Applications made following rainfall have been of very little benefit in controlling slip skin. Ground application equipment preferred.

Grapes: Mold control for fruit on drying trays in fields—*Penicillium* spp., *Aspergillus* spp., *Botrytis Cinerea*, *Rhizopus* spp.—Use 10 pounds per acre. Apply to fruit on trays for protection of fruit and prevention of above molds. Apply to give even coverage. During damp weather make a second application after trays are turned. Apply this dust by hand or ground dusters only. Do not apply within 5 days of a preharvest application of Captan.

Mangos (0): *Cercospora* Spot or Blotch—Use 50 to 70 pounds per acre. Begin applications at first sign of disease. Repeat at regular intervals to maintain control.

Peaches (0), Nectarines (0): Brown Rot, Peach Scab—Use 50 pounds per acre. Make 2 to 3 prebloom applications. Repeat at petal fall making 3 to 4 cover sprays as necessary. Make preharvest application where necessary.

Pears (0): Pear Scab (except Pacific Northwest)—Use 20 to 30 pounds per acre. Make first applications during early finger and petal fall stages. Repeat applications for each infection period or at weekly intervals as necessary. Russeting may be reduced on Boac Pears. Do not use D'Anjou Pears.

Plums (0), Prunes (0): Brown Rot—Use 20 to 50 pounds per acre. Apply at redbud and 75 percent bloom, full bloom, petal fall and in cover sprays as necessary. Thorough coverage is essential for maximum control.

Strawberries (0): Botrytis Rot—Use 20 to 40 pounds per acre. Make first application when new growth starts in spring before fruit starts to form. Repeat applications weekly. Severe Infection Conditions: Continue applications through harvest period, treating immediately after each picking.

Beans, Peas: Damping-Off, Root Rot—Use 60 pounds per acre. Work into top 3 or 4 inches of soil before planting.

Cantaloupes (0), Cucumbers (0), Pumpkins (0), Squash (Summer-Winter) (0), Watermelon (0): Downy Mildew, Angular Leaf Spot, Anthracnose—Use 20 to 25 pounds per acre on young plants, increasing to 30 to 35 pounds per acre on mature plants. Apply at first sign of disease. Repeat at 5 to 7 day intervals until danger of infection is past.

Celery (Field and Seed Bed only): Pink Rot, Damping-Off—Use 35 to 65 pounds per acre. Apply when all seeds have germinated. Repeat as necessary.

Potatoes: Early and Late Blight—Use 17½ to 30 pounds per acre. Begin applications when plants are 2 to 4 inches high and repeat at 5 to 7 day intervals as long as disease threatens.

Potatoes: (Seed Pieces): Seed Rot and Damping-Off—Dust at 7½ ounces per bushel cut seed pieces (12 ozs. or ¾ pound to 100 pounds cut seed pieces). Apply in convenient container or by duster attachment

over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 or 2 days the treated pieces should be stored for 2 days in open crates before bagging.

Tomatoes: Early Blight, Late Blight, Anthracnose, Septoria Leaf Spot—Young Plants—Use 25 to 30 pounds per acre. Mature Plants—Use 35 to 50 pounds per acre. Begin applications at first sign of disease. Repeat at 5 to 7 day intervals.

Ornamentals: Roses—Black Spot, Carnations—Rust and Leaf Spot—Chrysanthemums—Flower Blight—Use 22½ to 30 pounds per acre. Apply at first sign of disease to give even coverage of upper and lower leaf surfaces. Repeat at 7 to 10 day intervals as necessary.

Soil and Greenhouse Bench Treatment: Pre-planting for Damping-Off and Root Rot Diseases on Seedling or Transplants of Roses (and other shrubs and trees), many Flowers and Vegetables (such as Beans, Peas, Cucurbits) and Lawn Seedbeds—Use 1¼ to 2 pounds per 1000 square feet (50 to 60 pounds per acre). Work into upper 3 to 4 inches of soil before planting.

Stock: Botrytis Rot—Use 40 to 50 pounds per acre. Apply when disease appears. Repeat at weekly intervals as needed.

Turf and Lawn: Brown Patch, Leaf Spot, Damping-Off, Root Rot—Use 1 to 2 pounds per 1000 square feet. Begin treatment before infection occurs. Repeat at 7 to 10 day intervals throughout summer. Do not graze treated areas. Do not feed clippings to livestock.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

BEST DOCUMENT AVAILABLE

FMC—FMC Trademark