

Code: 30476

Net Weight

Captan Sulphur 0.25 Dust Fungicide

EPA REG NO 279-1459 AA

ACTIVE INGREDIENTS:

**Captan	BY WT
Sulphur	10.00
	15.00
INERT INGREDIENTS:	65.00%
	100.00%

**N-(2-Trichloromethyl-4-cyclohexenyl)-2-methylimidazole

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Avoid inhalation of dust. Avoid contact with skin. Avoid contact with feed or foodstuffs.

For Emergency Assistance Call 716-735-3765

See other cans for additional precautionary information.

Usage Caution

Use only as directed. Do not use on crops or plants not listed on the label. Do not use on crops or plants if the label does not specifically mention the crop or plant. Do not use on crops or plants if the label does not specifically mention the disease or pest.

DIRECTIONS FOR USE

Apply to crops and plants as directed on the label. Do not apply to crops or plants if the label does not specifically mention the crop or plant. Do not apply to crops or plants if the label does not specifically mention the disease or pest. Use a long-sleeved shirt and long-legged trousers or a coverall type garment to protect your body from contact with the dust. Do not use if you are pregnant or nursing.

For more information, contact your local Extension Office or the Agricultural Department.

Written by the Agricultural Department. All rights reserved. This product is registered with the Agricultural Department. Do not use if you are pregnant or nursing. Do not use if you are pregnant or nursing. Do not use if you are pregnant or nursing.

FMC Corporate
Agricultural Chemical Group
2000 Market Street
Philadelphia, PA



Cherries—Powdery Mildew, Rust, Leaf Spot, Anthracnose, Fruit Rot, Bacterial Spot, Blight, Fire Blight, and Cherry Leaf Miner. Apply at first sign of disease. Repeat at regular intervals as necessary to maintain control.

Grapes—Powdery Mildew, Black Rot, Leaf Spot, Anthracnose, and Grapevine Leafhopper. Apply 20-30 pounds per acre. Equipment for use of this product should be started after leaves start to show signs of disease. It is necessary to use ground application equipment, preferably a sprayer. Begin spraying July 1 to August 15, apply 4 to 5 times at 1 week intervals. Apply every row and direct toward the cluster. Treat the grapes that should be made while the clusters are still loose. The fungicide dust must be applied before rains. Whenever rain falls, apply an additional dust. Applications made following rains should be made at 7 to 10 day intervals to maintain control.

Nectarines, Peaches—Brown Rot, Powdery Mildew, Peach Scab. Use 40-50 pounds per acre. Apply at first sign of disease. Repeat at regular intervals as necessary to maintain control.

Ornamentals—Roses—Black Spot, Powdery Mildew, and Rust. **Carnations**—Powdery Mildew, Rust, and Leaf Spot. **Chrysanthemums**—Flower Blight, Powdery Mildew, and Rust. Dust for complete coverage. Apply at first sign of disease and repeat as necessary.

Potatoes—Early and Late Blight. Use 25 to 30 pounds per acre. Apply applications when plants are 2 to 4 inches high and repeat as necessary.

Plums, Prunes—Brown Rot, Plum Rust. Use 40-50 pounds per acre. Apply at first sign of disease. Repeat at regular intervals as necessary to maintain control.

Strawberries—Gray Mold, Botrytis, Powdery Mildew. Use 20-40 pounds per acre. Apply at first sign of disease. Repeat at regular intervals as necessary to maintain control.

Tomatoes—Early Blight, Late Blight, Anthracnose, Septoria Leaf Spot, Russet Mite—Young Plants. Use 20-40 pounds per acre. **Mature Plants**—Use 35-50 pounds per acre. Begin dusting when disease or mites first appear and repeat as necessary.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use—This product is sold on the basis of an agreement between the user and the manufacturer. The user agrees to use the product in accordance with the directions and recommendations on the label and to accept full responsibility for the results obtained. The manufacturer is not responsible for the results obtained from the use of this product if the user does not follow the directions and recommendations on the label.

Warranty—The manufacturer warrants that the product is free from defects in material and workmanship at the time of shipment. The manufacturer does not warrant the results obtained from the use of this product.

Directions and Recommendations—This product is to be used as a fungicide and insecticide. It is effective against a wide range of diseases and insects. The user should read the label carefully and follow the directions and recommendations on the label. The user should also read the literature accompanying the product for more detailed information.

Use of Product—This product is to be used as a fungicide and insecticide. It is effective against a wide range of diseases and insects. The user should read the label carefully and follow the directions and recommendations on the label. The user should also read the literature accompanying the product for more detailed information.

Damages—The manufacturer is not responsible for any damages or losses resulting from the use of this product. The user should read the label carefully and follow the directions and recommendations on the label. The user should also read the literature accompanying the product for more detailed information.