

CODE 543

FOLPET 50 WETTABLE

NET WGT.
25 lbs.

FUNGICIDE

DIRECTIONS FOR USE DO NOT USE IN UNDILUTED FORM

FOLPET 50 Wettable should not be used in combination with or closely following oil sprays. Do not use with TEPP formulations when a concentrate mixture is being applied.

OBSERVE DAYS INTERVAL BETWEEN LAST APPLICATION AND HARVEST INDICATED BY NUMBER IN () FOLLOWING CROP.

APPLES (0): Bitter Rot, Botryosphaeria, Alternaria Spot, Sooty Blotch, Brooks Fruit Spot, Black Rot, Fly Speck, Scab - 2 pounds per 100 gallons of water. Apply up to 800 gallons of spray per acre. Make first application at delayed dormant and repeat at 7 day intervals as disease threatens up to bloom. Avoid applications between bloom and July 1 - may cause fruit russetting. Use other suitable fungicides during this period. Begin applications of FOLPET after July 1 and repeat on 10 to 14 day schedule.

CHERRIES (RED TART) (0): Leaf Spot, Brown Rot - 2 pounds per 100 gallons of water. Apply at pre-bloom and popcorn stage. Repeat in bloom and at 7 to 14 day intervals.

GRAPEFRUIT (0), LEMONS (0), LIMES (0), ORANGES (0), TANGELOS (0), TANGERINES (0): Greasy Spot, Melanose, Scab - 2 pounds per 100 gallons of water. Use up to 1000 gals. water per acre. Apply at dormant, 2-3 petal fall, 2 weeks later and on fall flush in August and September.

GRAPES (0): 2 pounds per 100 gallons of water. Apply 250 to 300 gallons of spray per acre. Dead Arm - make first application when new growth is 4 to 6 inches long. Repeat when growth is 8 to 10 inches long. Black Rot, Downy Mildew - make first application just before bloom and repeat immediately after bloom. Depending upon the susceptibility of the grape variety, from 1 to 3 more sprays may be needed at 7 to 10 day intervals. Powdery Mildew - make first application 2 to 3 weeks after bloom and repeat 2 weeks later. NOTE: During extended periods of hot dry weather, foliage injury on several grape varieties may occur.

BLACKBERRIES (0), BOYSENBERRIES (0), DEWBERRIES (0), LOGANBERRIES (0), RASPBERRIES (0): Fruit Rot, Gray Mold, Spur Blight - 2 lbs. per 100 gals. water. Apply 250 gals. per acre. Apply when berries have set and repeat at 7 day intervals until harvest.

EPA REG. NO. 279-1508

ACTIVE INGREDIENT:

*Folpet 50.0%

INERT INGREDIENTS: 50.0%

100.0%

*N-(trichloromethylthio)phthalimide

ACCEPTED

CAUTION

KEEP OUT OF REACH OF CHILDREN.

For Emergency Assistance Call
716-735-3765

Avoid inhalation of dust or spray mist. Avoid contact with skin. Avoid storage near feed and food products.

ENVIRONMENTAL AND DISPOSAL CAUTIONS

This product is toxic to fish. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

STORE IN COOL, DRY PLACE. PROTECT FROM EXCESSIVE HEAT.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

BLUEBERRIES (0), HUCKLEBERRIES (0): Gray Mold, Powdery Mildew, Leaf Spot Gloeosporium - 2 lbs. per 100 gals. water. Apply 250 gals. per acre. Apply before first infection. Repeat at 7 day intervals until harvest.

CRANBERRIES (30): Fruit Rot - 9 pounds per acre in 200 to 300 gallons of water. Apply during bloom and repeat at 10 to 14 day intervals.

GOOSEBERRIES (0), CURRANTS (0): Leaf Spot - 4 lbs. per 100 gals. water. Apply 150 gals. per acre. Apply just before bloom period. Repeat at 7 to 10 day intervals in the post-bloom period.

STRAWBERRIES (0): Leaf Spot, Fruit Rot - 2 pounds per 100 gallons of water. Apply up to 200 gallons of spray per acre. Make first application before bloom and repeat at 7 day intervals until after harvest.


CELERY (7): Early Blight, Late Blight, Cercospora Leaf Spot - 2 pounds per 100 gallons of water. Apply after plants are set in field and established. Repeat at 7 day intervals as necessary to maintain control.

CUCUMBERS (0), PUMPKINS (0), SQUASH (0), MELONS (Cantaloupes, Watermelon, Honeydew) (0): Anthracnose, Downy Mildew, Powdery Mildew - 2 to 4 pounds per 100 gallons of water. Apply up to 200 gallons of spray per acre depending on size of plants. Make first application when first true leaves appear and repeat at weekly intervals up to harvest.

LETTUCE (0): Downy Mildew - 2 lbs. per 100 gals. water. Apply up to 200 gals. per acre. Apply when plants are established in field. Repeat at 7 day intervals until harvest.

ONIONS (0), GARLIC (0), LEEKS (0), SHALLOTS (0): Purple Blotch, Downy Mildew - Use 4 lbs. per 100 gals. water. Apply 200 gals. per acre. Apply before first infection. Repeat at weekly intervals until tops begin to dry before harvest.

TOMATOES (0): Anthracnose - 4 pounds per 100 gallons of water. Apply up to 200 gallons of spray per acre. Make first application during first bloom and repeat at 7 to 10 day intervals up to harvest.



ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



279-1508

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product, buyer and user agree to the following conditions.

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of claim must be reported to FMC within 90 days of discovery by buyer or user, failing which the user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain natural conditions created or incurred by buyer or user, FMC has no control, buyer and user, as a condition of purchase, shall assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

FMC
Chemicals