ACCEPTED 279-1**5**29

DIBROM 8 MISCIBLE - Additional Uses EPA Reg. No. 279-1529

UNDER THE PEDERAL BALLCTICIDE FUNGICIDE AND HODENTICIDE ACT FOR ECONOMIC POSSON REGISTER. ED UNDER NO. SUBJECT TO ATTACHED COMMENTS.

OCT 31 1974

ALFALFA - Field: To control Grasshoppers use 3/4 pint per acre before crop is 12 inches high. To control Aphids, Loopers (including Cabbage Loopers), Leafhoppers, Lygus Bugs and Armyworms use 1 pint per acre. Seed: To control Lygus Bugs and Thrips use 1/2 pint per Combine with recommended chemicals for residual control. Mites use 1/2 pint per acre. Apply once for suppression of populations, or twice 7 days apart for control. Apply in a minimum of 20 gallons of water by ground or 5 gallons of water by air (10 gallons of water is preferred on seed alfalfa). Do not apply within 4 days of harvest.

STRAWBERRIES: To control Leafroller, Spider Mites, Aphids and Omnivorous Leaftier use 1 pint per acre. Do not apply within 1 day of harvest. Dormant: To control Twospotted Mites and Aphids, 1 gallon of Supreme Oil may be added per 100 gallons of spray mix.

CONCENTRATE SPRAY EQUIPMENT - Vegetable and Field Crops - Unless otherwise noted, use recommended amount in 20 to 30 gallons of water per acre.

CELERY: To control Aphids, Loopers (including Cabbage Loopers), Armyworms, Saltmarsh Caterpillars, and Leafminers use 1 pint per acre. Do not use within 4 days of harvest.

LETTUCE, SPINACH, SWISS CHARD, TURNIP GREENS: To control Aphids, Loopers (including Cabbage Loopers), Armyworms, Saltmarsh Caterpillars and Leafminers use 1 pint per acre. Do not use within 1 day of harvest.

CANTALOUPES, HONEYDEW MELONS, MUSKMELONS, WATERMELONS, PUMPKINS, WINTER SQUASH: To control Loopers (including Cabbage Loopers), Armyworms, Leafhoppers, Spider Mites and Leafminers use 1 pint per acre. By Air use 1 to 2 pints in a minimum of 5 gallons of water per acre. Do not apply when temperature is over 90 degrees F. Do not apply within 1 day of harvest.

TOMATOES: To control Drosophila use 1 pint per acre. Apply 5 to 7 days before first picking and repeat at 5 to 7 day intervals until the day before final harvest. To control Flea Beetles, Hornworms, Leafminers, Mites and Tomato Fruitworm use 1 pint per 100 gallons of water. Apply up to 200 gallons of finished spray per acre.