


30497



28

SUPREME OIL

FOR WESTERN AGRICULTURE

AMOUNTS GIVEN ARE FOR 100 GALLONS OF DILUTED SPRAY.

ALMONDS, APRICOTS, CHERRIES, NECTARINES, PEACHES, PLUMS, PRUNES: Dormant or Delayed Dormant: Brown Apricot Scale, Black Scale, San Jose Scale, Olive Scale (Parlatoria Scale), Oleander Scale, Frosted Scale, Italian Pear Scale, European Fruit Lecanium, Putnam Scale, Parlatoria, Soft Brown Scale, Certain Other Scale Insects, Eggs of Aphids, Red Spider Mites — 1½ to 2 gallons. Summer Use: Red Spider Mites — 3 to 4 quarts; Summer Stages of Brown Apricot Scale, San Jose Scale — 1 gallon.

APPLES: Rust Mite — Delayed Dormant — 1½ to 2 gallons. Summer Use — 1 to 1½ gallons.

APPLES, PEARS: Dormant or Delayed Dormant: San Jose Scale, Italian Pear Scale, Eggs of Aphids, Red Spider Mites — 1½ to 2 gallons; Summer Use: Red Spider Mites — 3 to 4 quarts; Summer Forms of San Jose and Italian Pear Scale — 1 to 1½ gallons.

FIGS: Fig Scale — Dormant — Use 3 gallons Supreme Oil or 2½ gallons Supreme Oil plus ½ pint Ethion 4.0 Miscible. Severe infestation — Use 4 gallons Supreme Oil plus 1 pint Phoskil 4.0 Miscible.

GRAPES: Mealybug — Dormant, 2 gallons per 2½ quarts Parathion 4.0 Emulsifiable in 200 gallons spray per acre depending upon size of vines.

LEMONS, GRAPEFRUIT: Red Scale, Yellow Scale, Purple Scale, Black Scale, Red Spider Mites — 1-1½ gallons Supreme Oil plus 1 pound Malathion 25 Wettable. For best results, spray during winter and spring months when Red Scale are more easily controlled and trees generally less reactive to oil sprays. Do not use within 8 weeks of harvest on lemons.

OLIVES: Parlatoria Scale, Spider Mites, Oleander Scale, Black Scale — Use 1½ gallons Supreme Oil or combine 1½ gallons Supreme Oil with ½ pint Phoskil 4.0 Miscible or 12 pounds Phoskil 25 Spray per 100 gallons water. Apply in the pre-bloom or in post-bloom stage up to August 1st.

NAVEL ORANGES: Red Scale, Black Scale, Yellow Scale, Purple Scale, Red Spider Mites — 3 quarts Supreme Oil plus 1 pound Malathion 25 Wettable. Do not use on Navel Oranges after September 1st.

VALENCIA ORANGES: Red Scale, Black Scale, Yellow Scale, Purple Scale, Red Spider Mites 1 gallon. Apply no later than September 1st; or use 3 quarts Supreme Oil plus

INSECTICIDE

EPA REGISTRATION NO 279-1539 AA

ACTIVE INGREDIENT:	BY WT.
Petroleum Oils	98.00%
INERT INGREDIENTS	2.00%
	100.00%

Petroleum Oils Min. Unsulfonated Residue92.00%

Petroleum Oils Classification Unclassified

Supreme Oil is an especially refined Phytonomic oil contact Insecticide, and a "sticking agent" for various insecticides and fungicides. It is manufactured to give maximum insecticide efficiency. Supreme Oil is for use on plants in foliage or the dormant stages. The high insecticide efficiency of Supreme Oil makes it possible to use much smaller amounts than normally used of petroleum oils for the summer and dormant spray control of various insects. Read Entire Label. Use Strictly In Accordance With Label Cautions, Warnings and Directions.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Do not induce vomiting. Call a Physician Immediately.

NOTE: If other products are combined with Supreme Oil observe the precautions on that product label.

ORNAMENTALS and SHADE TREES (Conifer Hawthorne, Flowering Peach, Flowering Oak, Privet, Birch): Soft Brown Scale, Oyst Scale, San Jose Scale, Red Spider Mites and Spray on Deciduous Varieties — 2 gallons per Varieties — 1 gallon per 100 gallons water spruce. Do not use on conifers or ferns no **CORN:** Supreme Oil may be used in combination 1 gallon in 20-40 gallons of water per than 5 gallons of water per acre by air. For use given on an Aatrex 80 WP label

MIXING DIRECTIONS: With agitator running combining with Wettable Powder insecticide when tank is 1/3 filled. Add Supreme Oil. is sufficient agitation of the spray to insure

CAUTION: Before using make certain spray sulfur within 30 days or Captan or Phalton Supreme Oil. (If in doubt, spray a portion Captan or Phalton using 1 quart to 25 gallons sufficient time has not elapsed). Avoid application hot weather (90° F.), rainy weather or above 40° F. Avoid over spraying or do healthy condition and when moisture conditions disease, drought or high nitrogen application. Rinse equipment and **DISPOSE OF WASTE** water supplies. Punch holes in containers and supplies.

Do not apply when weather conditions do not apply when runoff is likely to occur. Do not in accordance with label directions. See residues or other undesirable results may

Dealer Should Sell in **TERMS OF SALE OR USE** On purchase or use following conditions: **WARRANTY:** Niagara warrants that this product conforms to the ingredient statement on the

ACCEPTED
 G1772
 COMMENTS



SUPREME OIL

28 Gallons Net Contents

IN AGRICULTURE
 OF DILUTED SPRAY.
APPLES, PEACHES, PLUMS, PRUNES: Dormant or Black Scale, San Jose Scale, Olive Scale, Black Scale, Italian Pear Scale, European Fruit Brown Scale, Certain Other Scale Insects, 2 gallons. Summer Use: Red Spider Mites — Apricot Scale, San Jose Scale — 1 gallon.
 1 1/2 to 2 gallons. Summer Use — 1 to 1 1/2
 dormant: San Jose Scale, Italian Pear Scale. Eggs 2 gallons; Summer Use: Red Spider Mites — and Italian Pear Scale — 1 to 1 1/2 gallons
 2 gallons Supreme Oil or 2 1/2 gallons Supreme Oil infestation — Use 4 gallons Supreme Oil plus
 per 2 1/2 quarts Parathion 40 Emulsifiable in upon size of vines.
 Scale, Purple Scale, Black Scale, Red Spider 1 pound Malathion 25 Wettable. For best with when Red Scale are more easily controlled trays. Do not use within 8 weeks of harvest
 canker Scale, Black Scale — Use 1 1/2 gallons Supreme Oil with 1/2 pint Phoskol 40 Miscible or 2 gallons water. Apply in the pre-bloom or in
 Yellow Scale, Purple Scale, Red Spider Mites — Malathion 25 Wettable. Do not use on Navel
 Scale, Yellow Scale, Purple Scale, Red Spider Mites — 1st, or use 3 quarts Supreme Oil plus

INSECTICIDE

EPA REGISTRATION NO. 279-1539 AA

ACTIVE INGREDIENT:	BY WT.
Petroleum Oils	98.00%
INERT INGREDIENTS	2.00%
	100.00%
Petroleum Oils Min. Unsulfonated Residue	92.00%
Petroleum Oils Classification	Unclassified

Supreme Oil is an especially refined Phytonomic oil contact Insecticide, and a "sticking agent" for various insecticides and fungicides. It is manufactured to give maximum insecticide efficiency. Supreme Oil is for use on plants in foliage or the dormant stages. The high insecticide efficiency of Supreme Oil makes it possible to use much smaller amounts than normally used of petroleum oils for the summer and dormant spray control of various insects. Read Entire Label. Use Strictly In Accordance With Label Cautions, Warnings and Directions.

CAUTION
KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Do not induce vomiting. Call a Physician Immediately.
 NOTE: If other products are combined with Supreme Oil observe the precautions on that product label.

ORNAMENTALS and SHADE TREES (Conifers (See Note): Holly, Pyracantha, Sycamores, Hawthorne, Flowering Peach, Flowering Cherry, Apple, Magnolia, Maple, Dogwood, Oak, Privet, Birch): Soft Brown Scale, Oyster Shell Scale, Brown Apricot Scale, Black Scale, San Jose Scale, Red Spider Mites and Eggs 1 gallon to 100 gallons water. Dormant Spray on Deciduous Varieties — 2 gallons per 100 gallons water. Dormant Spray on Other Varieties — 1 gallon per 100 gallons water. NOTE: May cause discoloration of blue spruce. Do not use on conifers or ferns not tolerant to oil sprays.
CORN: Supreme Oil may be used in combination with Aatrex 80 WP at the rate of 1 gallon in 20-40 gallons of water per acre by ground and 1/2 gallon in not less than 5 gallons of water per acre by air. Follow the cautions, limitations and directions for use given on an Aatrex 80 WP label
MIXING DIRECTIONS: With agitator running start filling spray tank with water. When combining with Wettable Powder insecticide or fungicide products, add these first when tank is 1/3 filled. Add Supreme Oil when tank is about 1/2 full and while there is sufficient agitation of the spray to insure complete emulsification.
CAUTION: Before using make certain spray tank is free of sulfur residues. Do not apply sulfur within 30 days or Captan or Phaltan within 90 days after or 60 days before Supreme Oil. (If in doubt, spray a portion of one tree previously treated with Sulfur, Captan or Phaltan using 1 quart to 25 gallons water. Burn will show in several days if sufficient time has not elapsed). Avoid application of oil sprays just prior to or during hot weather (90° F.), rainy weather or drying winds. Apply when temperature is above 40° F. Avoid over spraying or double spraying. Use only when trees are in healthy condition and when moisture conditions are sufficient. Trees weakened by disease, drought or high nitrogen applications may be injured by oil.
 Rinse equipment and DISPOSE OF WASTES by burying in non-crop land away from water supplies. Punch holes in containers and bury in non-crop land away from water supplies.
 Do not apply when weather conditions favor drift from the areas treated. Do not apply when runoff is likely to occur. Do not use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.
Dealer Should Sell in Original Packages Only.
TERMS OF SALE OR USE On purchase or use of this product buyer and user agree to the following conditions
WARRANTY: Niagara warrants that this product, as of the time of sale by Niagara, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the pur-