

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER-ED UNDER NO.274-155 BUBIECT

NET CONTENTS Gallons

CODE SUPERIOR 847 USDA REG. NO. 279-1551

DIRECTIONS FOR MIXING

Fill the spray tank about one-third full of water, start the agitator and pour in the required amounts of Superior, 70 Oil. If other spray materials are to be used with the oil, first mix the other materials thoroughly with the water in the spray tank, and then add the oil last as the filling of the tank is being completed.

DIRECTIONS FOR USE

TO CONTROL	Amount Pe 100 Gallon	
Apple Red Bug European Fruit Le- canium Scale Fruit Tree Leaf Roller	2 gallons 2 gallons 3 gallons	Delayed Dormant
European Red Mite San Jose Scale Scurfy Scale	2 gallons 2 gallons 3 gallons	Green Tip to Delayed Dormant Delayed Dormant Delayed Dormant
Pear Leaf Blister Mite Pear Psylla	3 gallons 2 gallons	Dormant Late Dormant
European Red Mite San Jose Scale European Fruit Le- canium	2 gallons 2 gallons 2 gallons	Delayed Dormant
	Apple Red Bug European Fruit Le- canium Scale Fruit Tree Leaf Roller European Red Mite San Jose Scale Scurfy Scale Pear Leaf Blister Mite Pear Psylla European Red Mite San Jose Scale European Fruit Le-	Apple Red Bug European Fruit Le- canium Scale Fruit Tree Leaf Roller European Red Mite San Jose Scale Scurfy Scale Pear Leaf Blister Mite Pear Psylla European Red Mite San Jose Scale Scurfy Scale European Red Mite San Jose Scale Sear Jose Scale Sear Jose Scale San Jose Scale

ACTIVE INGREDIENT	By Wt
Petroleum Oil (Superior Type)	96.0%
INERT INGREDIENTS:	4.0%

100.0%

Petroleum Oil Specifications:

Unsulpho	nated Residue	. 92 - 94%
Viscosity		70 Second

KEEP OUT OF REACH OF CHILDREN.

MANUFACTURED IN U.S.A. BY

NIAGARA CHEMICAL DIVISION FMC CORPORATION

AYER, MASSACHUSETTS GREELEY, COLORADO MALAGA, NEW JERSEY MIDDLEPORT, NEW YORK SOUTH HAVEN, MICHIGAN WYOMING, ILLINOIS

Home Office - Middleport, N.Y.



When a gun has been shut off or the outfit stopped for any length of time, spray on the ground for a few moments until the spray appears creamy white. This is desirable also where there are dead ends in the pipe lines of stationary outfits. All the material in stationary outfits should be sprayed out each day.

Spray on warm days in late Winter or early Spring, and before the buds separate into the clusters. Do not spray in freezing weather, or apply strong dosages after the buds have burst to any extent. Do not apply when buds have separated in the clusters.

Do not apply within 2 weeks of a sulfur spray or within 7 days of Captan.

DO NOT STORE AT TEMPERATURES BELOW 0° F.

Do not use if this spray oil does not emulsify. Correct emulsification is indicated if, when this product is added to the water in the spray tank, a uniform milk white spray mixture results, provided the spray mixture is being agitated. If the spray mixture, after adding the oil to the spray tank with the agitation running, is not milk-white, do not use the resulting spray or the remainder of this product.

Dealers Should Sell in Original Packages Only.

USAGE CAUTION: USE OF THIS MATERIAL IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH THE DIRECTIONS MAY PRODUCE PLANT INJURY, EXCESSIVE RESIDUES OR OTHER UNDESIRABLE RESULTS.

NOTICE: Niagara and Seller warrant, to the extent of the purchase price, that the product will conform to the ingredient statement set forth on the label but they make no other warranty, express or im-plied covering the product. Neither Niagara nor Seller shall be held responsible in any manner for personal injury, property damage or other loss of any kind resulting from the handling, storage or use of the product, whether or not in accordance with directions, regardless of the negligence or fault of Niagara or Seller. Buyer assumes all risk and liability therefrom and accepts and uses the product on these conditions. No person is authorized to vary or waive anything contained in this notice.

Printed in U. S.

ζ.