HINDER THE FEDERAL INSECTICIDE FUNCICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER ED UNDER NO. 279-153

## Insecticide

EPA Reg. No. 279-1551 EPA Est., 279-IL-1 Active ingredient: Petroleum Oil Specifications: 

# CAUTION KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed.

For Emergency Assistance Call 716 - 735-3765

Do not store at temperatures below 32°F

## Directions for Mixing

Fill the spray tank about one-third full of water, start the agitator and pour in the required amounts of Superior 70 Oil. If other spray materials are to be used with the oil. mix the other materials thoroughly with the water in the spray tank first, and then add the Superior 70 Oil as the filling of the tank is being completed. Do not use in undiluted form.

### **Usage Cautions**

Soray on warm days in late Winter or early Spring, and before the buds separate into the clusters. Do not spray in freezing weather, or apply strong dosages after the buds have burst to any extent. Do not apply when buds have separated in the clusters.

Do not apply within 2 weeks of sulfur spray or within 7 days of Captan.

Do not use if this spray oil does not emulsify. Correct emulsification is indicated if, when this product is added to the water in the soray tank, a uniform milk white spray mixture results, provided the spray mixture is being agitated. If the spray mixture, after adding the oil to the spray tank with the agitator running, is not milk-white, do not use the resulting spray or the remainder of this product.

When a gun has been shut off or the outfit stopped for any length of time, spray on the ground for a few moments until the spray appears creamy white. This is desirable also where there are dead ends in the pipe lines of stationary outfits. All the material in stationary outfits should be sprayed out each day.

#### Directions for Use

Apples, Pears: Apple Red Bug, European Fruit Lecanium Scale, San Jose Scale - Apply 2 gallons per 100 gallons of water in the delayed dormant stage. Fruit Tree Leaf Roller. Scurfy Scale - Apply 3 gallons per 100 gallons of water in the delayed dormant stage, European Red Mite - Apply 2 gallons per 100 gallons of water in the green tip to delayed dormant stage. Pear Leaf Blister Mite-Apply 3 gallons per 100 gallons of water in the domant stage. Pear Psylla-Apply 2 gallons per 100 gallons of water in the late dormant stage.

Fruit Lecanium -Use 2 gallons per 100 gallons of water in the delayed dormant stage. Terrapin Scale-Use 3 gallons per 100 gailons of water in the delayed domant stage.

Peaches: European Red Mite, San Jose Scale, European

### Environmental and disposal cautions

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not allow this material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

Do not reuse empty containers. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

#### Dealers Should Sell in Original Packages Only.

Terms of sale or use: On purchase or use of this product buyer and user agree to the following conditions:

Warranty: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably lit for the purposes set forth in the Directions for Use. Except as so warranted the product is sold as is. FMC makes no other warranty express or implied.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user walves any claim for such

Excuse from non-controllable conditions: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from, liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.





**FMC Corporation Agricultural Chemical Division** Middleport New York 14105