

Code 506

Sevin® 10 Dust

50 Lbs. Net Weight

Insecticide

EPA Reg No. 279-1569 EPA Est., 279-NJ-1

Active Ingredient

*Carbaryl (1-Naphthyl N-methylcarbamate) 10.0%

Inert Ingredients 90.0%

100.0%

*Sevin, U.S. Pat. Nos. 2,903,478; 3,009,855

Trademark of Union Carbide Corporation for the active ingredient carbaryl.

CAUTION

KEEP OUT OF REACH OF CHILDREN.

Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Skin contact may be harmful. Avoid contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. For Emergency Assistance Call 716-735-3765.

Note for Physician: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

Usage Caution

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application.

Directions for Use

Apples, Pears Codling Moth, Green Apple Aphid, Red-banded Leaf Roller, Pear Psylla, Apple Maggot, Periodical Cicada - Use 40 to 60 pounds per acre. To avoid possible fruit thinning do not apply until 30 days after full bloom. To avoid possible injury to McIntosh and York apples, do not apply before the second cover. Repeat at 10 day intervals if needed. Do not apply within 1 day of harvest.

Corn Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles, Japanese Beetles - Use 15 to 20 pounds per acre. For larvae in whorl and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks begin to dry. Three or more applications may be required. For European Corn Borer control make one or more applications beginning when borer damage threatens, depending on the extent of infestation and the degree of protection desired. Consult your state agricultural experiment station for proper timing of application. Application to the tassel region of sweet corn during the pollen shed period will seriously reduce bee populations.

Cotton Boll Weevil, Bollworm, Cotton Fleahopper, Lygus Bugs, Thrips, Pink Bollworm - Use 15 to 20 pounds of dust per acre. Apply when insects first threaten damage to crop, and repeat at 5 to 7 day intervals as required to maintain control.

Rice, Forage Grass and Pasture Thrips, Armyworms, Stink Bugs, Grasshoppers - Use 10 to 15 pounds of dust per acre. Apply when insects first threaten damage to crop. Repeat as necessary. Allow 14 days between last application and rice harvest.

Sorghum (Grain and Forage) Armyworms, Webworms, Corn Earworm, Stink Bugs, Grasshoppers, Sorghum Midge - Use 15 to 20 pounds per acre. For optimum control on grain sorghum direct dust into forming heads. For sorghum midge control, make applications 3 to 4 days after heads have emerged from boot. Do not apply within 21 days of harvest of grain. May be applied to day of harvest for sorghum forage.

Peaches Oriental Fruit Moth, Peach Twig Borer, Periodical Cicada - Use 20 to 40 pounds per acre. Apply when insects first appear and repeat every 10 days if needed. For Oriental fruit moth, apply at petal fall and every 10 to 14 days thereafter until control is achieved. A minimum of 30 pounds of dust is required to insure adequate coverage. Do not apply within 1 day of harvest.

Tobacco Hornworms, Budworms, Tobacco Flea Beetles, Grasshoppers, Japanese Beetles - Use at the rate of 15 to 20 pounds of dust per acre in broadcast treatment. Full coverage is essential. Begin treatments when worms are small. Repeat as needed. Use the highest dosage on large plants. For budworms when hand dust applications are directed to the bud and top leaves only, apply 4 to 6 pounds per acre. Avoid heavy applications with cloth bags as plant injury may result. Plant injury may occur when applied in presence of rain. If late applications are necessary, treat immediately after priming rather than before priming. Treated fields may be entered immediately after dust has settled. Do not apply within 3 days of priming or cutting.

Environmental and Disposal Cautions

This product is toxic to fish, shrimp, crabs, and other wildlife. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Dispose of wastes by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

LABEL

Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.**

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured in U.S.A. by:
FMC CORPORATION
AGRICULTURAL CHEMICALS DIVISION
Middlebury, N.Y. 14101

FMC
Chemicals