

CODE
257



SPECIMEN LABEL

NET WEIGHT
50 LBS.

KOLODUST EXTRADICHLONE 1 DUST

FUNGICIDE

FPA REG. NO. 279-1700

ACTIVE INGREDIENTS:

Sulfur	71.00%
*Dichlone	1.00%
INERT INGREDIENTS:	28.00%
<hr/>	
*2,3-dichloro-1,4-naphthoquinone	100.00%

Sulfur Fineness 99.7% thru 325 mesh.

CAUTION KEEP OUT OF REACH OF CHILDREN.


For Emergency Assistance Call 716-735-3765

May cause skin irritation. Avoid contact with skin, eyes or clothing. Avoid inhalation of dust. Wear protective clothing. Wash with soap and water after using. Avoid storage near feed and food products

This product is toxic to fish. Keep out of lakes, ponds or streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.



NIAGARA CHEMICAL DIVISION
FMC CORPORATION
MIDDLEPORT, N. Y. 14105



This product contains 29.3% bentonite sulfur.

Bentonite sulfur is made by the adsorption of fused sulfur into bentonite in the proportions of thirty parts sulfur and seventy parts bentonite. A portion of the sulfur particles are very minute, some of them colloidal. The process of manufacture and the character of the sulfur particles impart to the product distinctive properties with respect to toxicity to certain fungi, persistence on plants, suspensibility in spray tank and safety to host plants.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

DIRECTIONS FOR USE

APPLE: Powdery Mildew, Scab - Apply at the rate of 1/2 to 1 1/2 pound of dust per tree using ground equipment, or by airplane apply 30 to 35 pounds of dust per acre. Apply in the pink, petal-fall and calyx periods. In the event of rain, more applications may be required to maintain effective control. Use of this dust during the bloom period may result in blackening of the blossoms. During this period the use of a Niagara Niacide dust formulation is recommended if necessary. Do not apply later than 10 days after petal-fall.

PEACHES: Brown Rot (Blossom Blight) - Apply at the rate of 1/4 to 1/2 pound per tree not to exceed 40 pounds per acre. For blossom blight control, start applications during the pink stage of development. Repeat at 3 to 4 day intervals during bloom in order to protect newly opened blossoms. Do not apply after petal-fall.

On apples and peaches, do not apply within 10 days to 2 weeks of an oil spray. Do not apply prior to or during periods of excessively high temperatures.