



Code 1160

Lbs. Net Weight

Captan 7.5 Dust

Fungicide

EPA Reg. No. 279-1649

EPA Est.

Active Ingredients:	Wt.
*Captan.....	7.5%
Inert Ingredients:	92.5%
	100.0%

*N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide

KEEP OUT OF REACH OF CHILDREN

CAUTION

For Emergency Assistance Call 716-735-3785.
See Other Panels for Additional Precautionary Information.

FMC

FMC Corporation
Agricultural Chemical Group
2000 Market Street
Philadelphia Pennsylvania 19103

PA 5/80

PRECAUTIONARY STATEMENTS

Hazards to Humans (& Domestic Animals)

Caution

Avoid inhalation of dust. Avoid contact with skin. Avoid contamination of feed and foodstuffs.

Environmental Hazards

This product is toxic to fish. Keep out of lakes, ponds or streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not allow this material to drift onto neighboring crops or non-crop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues or other undesirable results may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Pesticide, spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of container in an incinerator or landfill approved for pesticide containers, or bury in a safe place. Consult Federal, State or local disposal authorities for approved alternative procedures such as limited open burning.

Apples: Apple Scab, Brooks Fruit Spot, Flyspecking, Bitter Rot, Black Pox, Botryosphaeria—Use 25 to 40 lbs. per acre. Apply at delayed dormant period. Repeat at weekly intervals as required to maintain control.

Apricots: Brown Rot, Jocke Rot—Use 40 to 60 lbs. per acre. Apply at red-bud stage and repeat at 75% petal-fall.

Cherries: Cherry Leaf Spot, Brown Rot—Use 50 to 60 lbs. per acre. Apply at popcorn, blossom and petal-fall stages. Continue at 10 to 14 day intervals as required.

Peaches: Brown Rot, Peach Scab—Use 50 to 60 lbs. per acre. Make 2 to 3 pre-bloom applications. Repeat at petal-fall and 3 to 4 covers as necessary. Make preharvest application where necessary.

Pears: Pear Scab (except Pacific Northwest)—Use 25 to 40 lbs. per acre. Apply during early finger and petal-fall stages. Repeat at weekly intervals as necessary to maintain control. Russeting may be reduced on Bosc pears. Do not use on D'Anjou pears.

Raspberries: Fruit Rot—Use 30 lbs. per acre making applications 3 to 5 days before harvest starts, at midharvest and 8 to 10 days later.

Strawberries: Botrytis Rot—Use 40 lbs. per acre. Make the first application when new growth starts in the spring and before the fruit starts to form. Repeat at weekly intervals. During severe infection periods continue through harvest, treating immediately after each picking.

Tomatoes: Anthracnose, Early Blight, Late Blight, Septoria Leaf Spot—On young plants use 25 to 30 lbs. per acre. On mature plants use 40 to 50 lbs. per acre. Begin applications at first sign of disease. Repeat at 7 to 10 day intervals.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: FMC makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted, the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by buyer at his own risk.

Use of Product: FMC's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied, is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

(1160-5/80-PA)