

Code 1160

50 Lbs. Net Weight

Captan 7.5 Dust

Fungicide

EPA Reg. No. 279-1649 EPA Est., 279-NJ-1

Directions for Use

Active Ingredients	
* Captan	7.5%
Inert Ingredients	92.5%
	100.0%

*N- [(trichloromethyl)thio] -4-cyclohexene-1,2-dicarboximide

CAUTION KEEP OUT OF REACH OF CHILDREN.

Avoid inhalation of dust. Avoid contact with skin. Avoid contamination of feed and foodstuffs.

For Emergency Assistance Call 716 - 735 - 3765

Apples Apple Scab, Brooks Fruit Spot, Flyspecking, Bitter Rot, Black Pox, Botryosphaeria - Use 25 to 40 lbs. per acre. Apply at delayed dormant period. Repeat at weekly intervals as required to maintain control.

Apricots Brown Rot, Jacket Rot - Use 50 to 60 lbs. per acre. Apply at red bud stage and repeat at 75% petalfall.

Cherries Cherry Leaf Spot, Brown Rot - Use 50 to 60 lbs. per acre. Apply at popcorn, blossom and petal fall stages. Continue at 10 to 14 day intervals as required.

Peaches Brown Rot, Peach Scab - Use 50 to 60 lbs. per acre. Make 2 to 3 pre-bloom applications. Repeat at petal-fall and 3 to 4 covers as necessary. Make preharvest application where necessary.

Pears Pear Scab (except Pacific Northwest) - Use 25 to 40 lbs. per acre. Apply during early finger and petalfall stages. Repeat at weekly intervals as necessary to maintain control. Russetting may be reduced on Bosc pears. Do not use on D'Anjou pears.

Raspberries Fruit Rot - Use 30 lbs. per acre making applications 3 to 5 days before harvest starts, at midharvest and 8 to 10 days later.

Strawberries Botrytis Rot - Use 40 lbs. per acre. Make the first application when new growth starts in the spring and before the fruit starts to form. Repeat at weekly intervals. During severe infection periods continue through harvest, treating immediately after each picking.

Tomatoes Anthracnose, Early Blight, Late Blight, Septoria Leaf Spot - On Young plants use 25 to 30 lbs. per acre. On mature plants use 40 to 50 lbs. per acre. Begin applications at first sign of disease. Repeat at 7 to 10 day intervals.

ACCEPTED

18 DEC 1975

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 279-1649

Potatoes (Seed Piece Treatment) Seed Rot, Damping Off, Verticillium Wilt, Brown Eye Disease - Dust at the rate of 10 ounces per bushel of cut seed pieces (1 lb. per 100 lbs. cut seed pieces). Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 to 2 days the treated pieces should be stored for 2 to 3 days in open crates before bagging. Do not use treated seed for food or feed purposes.

Potatoes (Young Plants) Early Blight, Late Blight - Use 30 to 35 lbs. per acre. Begin applications when plants are 2 to 4 inches high. Repeat at 7 to 10 day intervals as long as disease threatens.

Potatoes (Mature Plants) Early blight, Late Blight - Use 40 lbs. per acre. Begin applications at first sign of disease. Repeat at 7 to 10 day intervals.

Ornamentals (Roses, Carnations, Chrysanthemums) Black Spot, Leaf Spot, Rust Flower Blight - Apply at first sign of disease. Repeat at 7 to 10 day intervals as necessary. Dust plants thoroughly.

Environmental and Disposal Cautions
This product is toxic to fish. Keep out of lakes, ponds or streams. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Dispose of wastes by burying in non-croplands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED.



Dealers Should Sell in Original Packages Only.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS. DAMAGE TO ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase or use of this product buyer and user agree to the following conditions:

WARRANTY: FMC warrants that this product, as of the time of sale by FMC, (1) conforms to the ingredient statement on the label, and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED**

DAMAGES: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages. Notice of any claim must be reported to FMC within 10 days of discovery by buyer or user, failing which buyer or user waives any claim for such damage.

EXCUSE FROM NON-CONTROLLABLE CONDITIONS: Because of certain critical conditions created or incurred by buyer or user or over which FMC has no control, buyer and user, as a condition of purchase or use, assume responsibility for and release FMC from liability arising out of the handling or use of the product attributable to such causes. No person is authorized to vary or waive any statement contained herein.

Manufactured in U.S.A. by:
FMC CORPORATION
AGRICULTURAL CHEMICAL DIVISION
Middleport, N.Y. 14105

FMC
Chemicals