

Code
30888

MALATHION SEVIN 4-5 DUST

50 Pounds
Net Weight

[279-1707]

INSECTICIDE

Apply to plants before insects appear. For control of insects, Malathion and Sevin are effective against many insects. For control of insects, Malathion and Sevin are effective against many insects. For control of insects, Malathion and Sevin are effective against many insects.

BEANS (1) To control Leafhoppers, Whitefly, Aphids, Cucumber Beetle, Western Bean Cutworm, Corn Earworm and Armyworm use 40 to 40 pounds per acre.

CABBAGE (7), **CAULIFLOWER** (2), **BROCCOLI** (3), **BRUSSELS SPROUTS** (7), **KOHLRABI** (7), **COLLARDS** (14), **BEETS** (14), **TURNIPS** (14) and **MUSTARD** (14) To control Flea Beetles, Leafhoppers, Corn Earworm, Cabbage Caterpillars, Armyworms and Aphids use at the rate of 30 to 40 pounds per acre. May be applied to within 3 days of harvest on turnips and to within 7 days of harvest on beets if tops are not to be used as food or feed.

CUCUMBERS (1), **MELONS** (1), **PUMPKIN** (3), and **SQUASH** (1) To control Pineappleworm, Melonworm, Cucumber Beetle, Flea Beetle, Leafhoppers, and Squash Bug use 20 pounds per acre. Do not apply unless plants are dry.

LETTUCE (Head) (14) To control Flea beetles, Leafhoppers, Corn Earworm, Armyworm, Cabbage Caterpillars and Aphids use at the rate of 30 to 40 pounds per acre.

PEAS (1) To control Leafhoppers, Colorado Potato Beetle, Armyworms, Bean Hoppers, Pea Aphid and Pea Weevil use 30 to 40 pounds per acre.

POTATOES (10), **TOMATOES** (1), **PEPPERS** (1) and **SWEET POTATOS** (3) To control Colorado Potato Beetle, Flea Beetle, Leafhoppers, Flea, Armyworm, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bug and Stink Bug use 20 to 25 pounds per acre. On Potatoes apply as a foliage application.

RASPBERRIES (7), **BLYTHEBERRIES** (7), **BLACKBERRIES** (7), **GOOSEBERRIES** (7) and **PAIRBERRIES** (7) To control Leafhoppers, including Greenhopper, Flea Beetle, Ground Squirrel, Tree Cricket, Orange Tortrix, and Blueberry Aphids use 20 to 25 pounds per acre.

STRAWBERRIES (7) To control Leafhoppers, Flea Beetle, and Strawberry Root Weevil use 20 to 25 pounds per acre.

WATERMELONS (1) To control Pineappleworm, Melonworm, Cucumber Beetle, Flea Beetle, Leafhoppers, and Squash Bug use 20 pounds per acre. Do not apply unless plants are dry.

Apply to plants before insects appear. For control of insects, Malathion and Sevin are effective against many insects. For control of insects, Malathion and Sevin are effective against many insects. For control of insects, Malathion and Sevin are effective against many insects.

BEANS (1) To control Leafhoppers, Whitefly, Aphids, Cucumber Beetle, Western Bean Cutworm, Corn Earworm and Armyworm use 40 to 40 pounds per acre.

CABBAGE (7), **CAULIFLOWER** (2), **BROCCOLI** (3), **BRUSSELS SPROUTS** (7), **KOHLRABI** (7), **COLLARDS** (14), **BEETS** (14), **TURNIPS** (14) and **MUSTARD** (14) To control Flea Beetles, Leafhoppers, Corn Earworm, Cabbage Caterpillars, Armyworms and Aphids use at the rate of 30 to 40 pounds per acre. May be applied to within 3 days of harvest on turnips and to within 7 days of harvest on beets if tops are not to be used as food or feed.

CUCUMBERS (1), **MELONS** (1), **PUMPKIN** (3), and **SQUASH** (1) To control Pineappleworm, Melonworm, Cucumber Beetle, Flea Beetle, Leafhoppers, and Squash Bug use 20 pounds per acre. Do not apply unless plants are dry.

LETTUCE (Head) (14) To control Flea beetles, Leafhoppers, Corn Earworm, Armyworm, Cabbage Caterpillars and Aphids use at the rate of 30 to 40 pounds per acre.

PEAS (1) To control Leafhoppers, Colorado Potato Beetle, Armyworms, Bean Hoppers, Pea Aphid and Pea Weevil use 30 to 40 pounds per acre.

POTATOES (10), **TOMATOES** (1), **PEPPERS** (1) and **SWEET POTATOS** (3) To control Colorado Potato Beetle, Flea Beetle, Leafhoppers, Flea, Armyworm, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bug and Stink Bug use 20 to 25 pounds per acre. On Potatoes apply as a foliage application.

RASPBERRIES (7), **BLYTHEBERRIES** (7), **BLACKBERRIES** (7), **GOOSEBERRIES** (7) and **PAIRBERRIES** (7) To control Leafhoppers, including Greenhopper, Flea Beetle, Ground Squirrel, Tree Cricket, Orange Tortrix, and Blueberry Aphids use 20 to 25 pounds per acre.

STRAWBERRIES (7) To control Leafhoppers, Flea Beetle, and Strawberry Root Weevil use 20 to 25 pounds per acre.

WATERMELONS (1) To control Pineappleworm, Melonworm, Cucumber Beetle, Flea Beetle, Leafhoppers, and Squash Bug use 20 pounds per acre. Do not apply unless plants are dry.

ACCEPTED

CAUTION

KEEP OUT OF REACH OF CHILDREN

For Emergency Assistance Call
716-735-3765

Harmful if inhaled or swallowed. Avoid breathing of dust. Do not take internally. Skin contact may be harmful. Avoid contact with skin. Wash hands and face before eating. Take shower or bath after work. Wear goggles, dustproof work clothes, and clean clothes daily.

NOTE TO PHYSICIANS: Malathion and Malathion Sevin are contraindicated with Atropine and similar drugs.

AVOID CONTAMINATION OF FOOD, FEEDS, DRINKING WATER AND WATERING EQUIPMENT.

Malathion and Malathion Sevin are contraindicated with Atropine and similar drugs.

Apply to plants before insects appear. For control of insects, Malathion and Sevin are effective against many insects. For control of insects, Malathion and Sevin are effective against many insects. For control of insects, Malathion and Sevin are effective against many insects.

BEANS (1) To control Leafhoppers, Whitefly, Aphids, Cucumber Beetle, Western Bean Cutworm, Corn Earworm and Armyworm use 40 to 40 pounds per acre.

SUGAR BEETS (14) To control Flea beetles, Leafhoppers, Corn Earworm, Armyworm, Cabbage Caterpillars and Aphids use at the rate of 30 to 40 pounds per acre.


ORNAMENTALS: To control Aphids, Flea Beetles, Box Elder Bug, Leaf Hopper, Leaf Hoppers, Plant Bugs, Thrips, exposed Psyllids and Tent Caterpillars. Apply for full coverage. Injury may occur on certain ferns and some species of grasses. Do not use on *Baccharis* and *Conium* species.

TOBACCO: To control Aphids, Flea Beetles, Flea Beetle, and Beet. For 100 sq. yds. of plants, apply 2 to 3 pounds dust depending upon size of plants. Start applications when insects first appear in the plants. Apply at 4 day intervals until transplanting time.

Flea beetle use 15 to 25 pounds per acre. Full coverage is essential. Reapplications needed. Do not apply to plants near water. Reapply immediately after pruning rather than after pruning. Do not apply to plants near water.

Apply as a foliage application.

ADDITIONAL INFORMATION APPEARING ON BAGS TO WHICH THIS LABEL WILL BE ATTACHED



Dealers Should Sell in Original Packages Only

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

TERMS OF SALE OR USE: On purchase of this product, the user agrees to the following conditions:
WARRANTY: FMC warrants that this product, as at the time of sale by FMC, conforms to the ingredient statement on the label and is its responsibility for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED THE PRODUCT IS SOLD AS IS. FMC MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.
DAMAGES: Buyer's or user's recovery for damages for each of warrants or negligence shall be limited to the amount of the purchase price and shall not include consequential or incidental damages. Recovery by buyer or user shall be limited to the amount of the purchase price. FMC shall not be liable for such damage or loss FROM NON-CONTROLLED USE CONDITIONS. Because certain uses are optional, damage incurred by buyer or user when FMC has no control over use and user assumes liability for purchase shall assume responsibility for and release FMC from liability existing in the use of the product and from any such release. No party shall be held liable for any damage or loss resulting from such use.

